



Topological Charge and Cooling Scales in Pure $SU(2)$ Lattice Gauge Theory

David A. Clarke

Florida State University

HEP Seminar, March 21, 2018

B. A. Berg and D. A. Clarke, Phys. Rev. D **97** 054506 (2018)

- ▶ To reach the continuum limit, send the lattice spacing to zero (compared to a physical length)
- ▶ In the continuum limit, one can predict dimensionless mass ratios $c_k = m_k/m_0$ (m_0^{-1} can serve as a length scale)
- ▶ Choosing the reference m_0 is called **scale setting**
- ▶ Lüscher introduces the gradient flow¹, suggests gradient scale as a new reference scale; scale setting gains renewed interest
- ▶ Bonati and D'Elia² suggest standard cooling³ can be used similarly for scale setting; advantage in computational efficiency
- ▶ We verify⁴ this can be done in pure SU(2)
- ▶ This project: Investigate whether there is any dependence of the cooling scales on topological charge
- ▶ This project: Obtain a new estimate for topological susceptibility

¹M. Lüscher, J. High Energy Phys. 2010 (2010).

²C. Bonati and M. D'Elia, Phys. Rev. D, 89 (2014).

³B. A. Berg, Phys. Lett. B, 104, 475 (1981).

⁴B. A. Berg and D. A. Clarke, Phys. Rev. D, 95 (2017).

Outline



1. Topological charge
2. Brief overview of lattice
3. Results
4. Conclusion



1. Topological charge
2. Brief overview of lattice
3. Results
4. Conclusion



Consider classical, euclidean $SU(2)$ gauge theory

$$\mathcal{L} = -\frac{1}{2g^2} \text{tr} F_{\mu\nu} F_{\mu\nu} = \frac{1}{4} F_{\mu\nu}^a F_{\mu\nu}^a.$$

The quantum theory has infinitely many vacuum states.

1. Classical vacuum state: $F_{\mu\nu}^a = 0 \Rightarrow A_\mu = U \partial_\mu U^\dagger$
2. If you can smoothly deform U into U' , a **topological charge** Q is conserved
3. If you can't smoothly deform U into U' , there is an energy barrier between A_μ and A'_μ (**topological sectors**)
4. There are infinitely many U all having different Q
5. This corresponds to different vacuum states in quantum theory

⁵M. Srednicki *Quantum Field Theory* Cambridge: Cambridge, 2007.

Topological charge



We can use **topological winding number**

$$Q \equiv -\frac{1}{24\pi^2} \int d^3x \epsilon_{ijk} \text{tr} U \partial_i U^\dagger U \partial_j U^\dagger U \partial_k U^\dagger$$

- ▶ Invariant under coordinate changes
- ▶ Invariant under smooth deformations of U
- ▶ Counts number of times U covers spatial three-sphere
- ▶ We can rewrite Q as

$$Q = \frac{1}{32\pi^2} \int d^4x \epsilon_{\mu\nu\rho\sigma} \text{tr} F_{\mu\nu} F_{\rho\sigma},$$

which will be useful on the lattice




1. Topological charge
2. Brief overview of lattice
3. Results
4. Conclusion

Pocket dictionary for pure SU(2) LGT

- ▶ 4D space-time with Euclidean metric and periodic BCs
- ▶ **Sites** $x = (an_1, an_2, an_3, an_4)$ with n_4 in time direction
- ▶ Regularization through **lattice spacing** a
- ▶ **Link variables** $U_\mu(x) = e^{-aA_\mu(x)} \in \text{SU}(2)$ on links
- ▶ **Plaquette** $U_{\mu\nu}^\square(x)$
- ▶ Hypercube of volume $(aN)^4$
- ▶ Discretized calculus

$$\int d^4x \leftrightarrow a^4 \sum_x$$

$$\partial_\mu f(x) \leftrightarrow \frac{f(x + a\hat{\mu}) - f(x)}{a}$$





- Bare coupling controls lattice spacing

$$a\Lambda_L = \exp\left(-\frac{1}{2b_0g^2}\right) (b_0g^2)^{-b_1/2b_0^2} \left(1 + \mathcal{O}(g^2)\right)$$

- **Continuum limit** $a \rightarrow 0$, *with care*
- **Wilson action:**

$$S = \beta \sum_{x, \mu < \nu} \left(1 - \frac{1}{2} \text{tr } U_{\mu\nu}^{\square}(x)\right) \approx -\frac{\beta}{8} \sum_x a^4 \text{tr } F_{\mu\nu}(x) F_{\mu\nu}(x)$$

- Identify $\beta = 4/g^2$ so that in continuum limit

$$S \rightarrow -\frac{1}{2g^2} \int d^4x \text{tr } F_{\mu\nu} F_{\mu\nu}$$

SU(2) LGT simulations



Calculate expected value of operator O

- ▶ In QFT expectation values given by

$$\langle O \rangle = \frac{1}{\mathcal{Z}} \int \mathcal{D}\phi O(\phi) e^{-S(\phi)}, \quad \mathcal{Z} \equiv \int \mathcal{D}\phi e^{-S(\phi)}$$

- ▶ **Markov Chain Monte Carlo** basic idea:
 - ▶ Each configuration generated depending on last one only
 - ▶ Accept new configuration with probability $\min\{1, e^{-\Delta S}\}$
 - ▶ Create a **time series** of measurements O_n of O
 - ▶ Multiple **sweeps** between measurements to reduce correlation
- ▶ The estimator for $\langle O \rangle$ on the lattice is

$$\bar{O} = \frac{1}{N_{\text{conf}}} \sum_{n=1}^{N_{\text{conf}}} O_n$$

Q on the lattice

Naive discretization using pocket dictionary:

$$Q = \frac{1}{32\pi^2} \int d^4x \epsilon_{\mu\nu\rho\sigma} \text{tr} F_{\mu\nu} F_{\rho\sigma}$$

becomes

$$Q = \frac{1}{2^9\pi^2} a^4 \sum_x \sum_{\mu\nu\rho\sigma=\pm 1}^{\pm 4} \epsilon_{\mu\nu\rho\sigma} \text{tr} U_{\mu\nu}^\square(x) U_{\rho\sigma}^\square(x)$$

- ▶ Sum over backward directions to symmetrize
- ▶ Suffers from short-ranged renormalization effects

Topological susceptibility

$$\chi = \frac{1}{V} \langle Q^2 \rangle = \frac{1}{N^4} \frac{1}{N_{\text{conf}}} \sum_{i=1}^{N_{\text{conf}}} Q_i^2$$

- ▶ Provides information about distribution of Q
- ▶ Phenomenological interest

We use “smoother” using **standard cooling**

$$V_{\mu}(x, n_c) = \frac{V_{\mu}^{\sqcup}(x, n_c - 1)}{\sqrt{\det V_{\mu}^{\sqcup}(x, n_c - 1)}}.$$

- ▶ Locally minimizes action
- ▶ Suppresses local fluctuations without changing global charge
- ▶ Extract Q and χ from cooled configurations, after these quantities become “metastable”
- ▶ Q will “freeze out” at large enough β

Given **target value** y , a **cooling scale** $s(\beta)$ is defined by

$$y(t) = t^2 \langle E_t \rangle \quad \text{with} \quad s(\beta) = \sqrt{t(\beta)}$$



1. Topological charge
2. Brief overview of lattice
3. Results
4. Conclusion


Table: Lattices used for both χ and $L(Q)$ analysis.

| β | Measurements \times sweeps | N |
|---------|--|---------------------------|
| 2.710 | 128×2^{13} | 16, 28, 40 |
| 2.751 | 128×2^{13} | 16, 28, 40 |
| 2.816 | 128×2^{13} | 28, 40, 44 |
| 2.875 | 128×2^{13} | 28, 40, 44, 52 |
| 2.928 | $128 \times 2^{13} \quad (\times 1.5)$ | 28, 40, 44, 52, 60 |


- ▶ Multiple N to extrapolate $N \rightarrow \infty$
- ▶ Multiple β to extrapolate $a \rightarrow 0$
- ▶ Generate 128 configurations by MCMC, then cool them
- ▶ Largest in study of χ in pure SU(2)

"Freezing out" of cooling trajectories


$$\beta = 2.300, N^4 = 16^4$$




$$\beta = 2.510, N^4 = 28^4$$




$$\beta = 2.751, N^4 = 40^4$$



$$\beta = 2.928, N^4 = 60^4$$



Stabilization of χ




χ and Q stable for $\beta \gtrsim 2.751$

Systematics: Finite size extrapolation




$$L_{10}\chi^{1/4}(\beta, N) = L_{10}\chi^{1/4}(\beta) + \frac{c}{N^4}$$



Systematics: Continuum limit extrapolation



$$L_{10}\chi^{1/4}(\beta) \approx L_{10}\chi^{1/4} + k a^2 \Lambda_L^2 = L_{10}\chi^{1/4} + c \left(\frac{1}{L_{10}(\beta)} \right)^2$$



Prediction for topological susceptibility



$$\chi^{1/4}/\sqrt{\sigma} = 0.4446(82) \text{ for } n_c = 1000$$

$$\chi^{1/4}/\sqrt{\sigma} = 0.4655(87) \text{ for } n_c = 100$$

Table: Comparisons with past predictions using Gaussian difference tests.

| $\chi^{1/4}/\sqrt{\sigma}$ | q_{1000} | q_{100} |
|----------------------------|------------|-----------|
| 0.501(45) ⁶ | 0.31 | 0.44 |
| 0.528(21) ⁷ | 0.00 | 0.01 |
| 0.480(23) ⁸ | 0.32 | 0.56 |
| 0.4831(56) ⁹ | 0.01 | 0.09 |
| 0.4745(63) ⁹ | 0.07 | 0.40 |
| 0.4742(56) ⁹ | 0.06 | 0.40 |

⁶P. De Forcrand, M. G. Perez, and I.-O. Stamatescu, Nucl. Phys. B, 499, 409 (1997).

⁷T. DeGrand, A. Hasenfratz, and T. G. Kovacs, Nuclear Physics B, 505, 417 (1997).

⁸B. Allés, M. D'Elia, and A. Di Giacomo, Phys. Lett. B, 412, 119 (1997).

⁹B. Lucini and M. Teper, J. of High Energy Phys. 2001, 050 (2001).

Dependence of cooling scale on Q

- ▶ Compared scales using Student difference tests
- ▶ Found $L(Q)$ statistically compatible with $L(-Q)$
- ▶ Found scales with $|Q| > 1$ to be statistically compatible
- ▶ Therefore combine into three bins


$$|Q| = 0, \quad |Q| = 1, \quad \text{and} \quad |Q| \geq 2.$$

- ▶ L_{10} highly correlated with L_{11} ; average them
- ▶ Statistical analysis on remaining $4 \times 15 = 60$ scales

Table: Example cooling scales on largest lattice, by sector, at $n_c = 1000$.

| β | $ Q $ | n | L_{10} | L_{11} | L_{12} |
|---------|----------|-----|-----------|-----------|-----------|
| 2.928 | 0 | 26 | 12.61(23) | 12.55(23) | 11.66(21) |
| | 1 | 49 | 12.74(18) | 12.68(17) | 11.66(18) |
| | ≥ 2 | 53 | 12.39(14) | 12.34(14) | 11.64(14) |

Dependence of cooling scale on Q





1. Topological charge
2. Brief overview of lattice
3. Results
4. Conclusion

- ▶ For large enough β and N , standard cooling can be used to obtain stable topological sectors
- ▶ Best estimate for topological susceptibility:

$$\chi^{1/4}/\sqrt{\sigma} = 0.4446(82),$$

which is surprisingly close to past predictions

- ▶ Within our statistics, we find no evidence of correlations between cooling scales due to topological charges

Thanks for listening