

A Comparison of Scales in Pure $SU(2)$ LGT

David A. Clarke

Florida State University

HEP Seminar, February 24, 2017



Outline

1. Motivation
2. Introduction
3. Meet the Scales
4. Scaling Analysis
5. Conclusion







1. Motivation
2. Introduction
3. Meet the Scales
4. Scaling Analysis
5. Conclusion



Lots of cool experiments

- ▶ Hadronic structure
- ▶ Hadron spectroscopy
- ▶ Studying confinement
- ▶ New physics searches
- ▶ ...and more



Lattice calculations are useful

- ▶ Lattice spacing provides a regularization of QFT
- ▶ Allows one to calculate non-perturbative quantities
- ▶ In principle arbitrary precision can be achieved provided enough computing power is available
- ▶ That can help guide experimental searches
- ▶ Can help confirm Standard Model with high precision, or otherwise hint at new physics



1. Motivation
2. Introduction
3. Meet the Scales
4. Scaling Analysis
5. Conclusion



Scale setting

- ▶ To reach the continuum limit, send the lattice spacing to zero (compared to a physical length)
- ▶ In the continuum limit, one can predict dimensionless mass ratios $c_k = m_k/m_0$ (m_0^{-1} can serve as a length scale)
- ▶ Choosing the reference m_0 is called **scale setting**
- ▶ Many challenging calculations rely on scales of previous work; in some cases inaccuracy of reference scale greatly influenced result
 - ▶ 2008 and subsequent 2010 HPQCD calculation of f_{D_s} [1, 2]
- ▶ Ideally, reference scale can be calculated with high precision using modest computational resources



Gauge symmetries and SU(2): A quick review

- ▶ Noether's Theorem: each continuous symmetry of the Lagrangian implies a conserved current
- ▶ SU(N) group of $N \times N$ unitary matrices with determinant 1
 - ▶ $N^2 - 1$ generators
 - ▶ Corresponding to each generator T^a is a gauge field A_μ^a
 - ▶ Particles associated with these fields are gauge bosons, but in SU(N) LGT we just call them “gluons”
- ▶ SU(2) has $2^2 - 1 = 3$ generators, the Pauli matrices σ^a
- ▶ The kinetic term

$$\mathcal{L}_G = -\frac{1}{2g^2} \text{Tr } F_{\mu\nu} F^{\mu\nu}$$

is invariant under local SU(2) transformations




Why pure $SU(2)$?

- ▶ When the Lagrangian has the $SU(N)$ kinetic term and nothing else, we call the theory **pure $SU(N)$**
- ▶ $SU(2)$ is a gauge group of the Standard Model
- ▶ Computationally simple compared to complex systems like QCD
 - ▶ Accessible to our resources
 - ▶ No fermions, which are challenging to put on the lattice
 - ▶ Comparatively short simulations provide large statistics
 - ▶ Allows one to study a wider range of lattice size and bare coupling combinations
- ▶ General insight discovered in pure $SU(2)$ can perhaps be applied valuable for complex theories such as QCD




Pure SU(2) LGT

- ▶ Non-perturbative approach to QFT
- ▶ 4D space-time with Euclidean metric and periodic BCs
- ▶ **Sites** $x = (n_1, n_2, n_3, n_4)$ with n_4 in time direction
- ▶ Regularization through **lattice spacing** a
 - ▶ Finite volume $a^4 N_\tau N_s^3$
 - ▶ Ultraviolet cutoff $\sim a^{-1}$
 - ▶ Infrared cutoff $\sim (aN)^{-1}$
- ▶ **Continuum limit** $a \rightarrow 0$



Pure SU(2) LGT

- ▶ **Link variables** or “gluons” $U_\mu = e^{-aA_\mu(x)} \in \text{SU}(2)$ on links
- ▶ For a closed path \mathcal{C} , the corresponding **Wilson loop** is $\text{Tr } U(\mathcal{C})$
 - ▶ Gauge invariant
 - ▶ Used to construct observables
 - ▶ **Plaquette** \square is the smallest Wilson loop



Pure SU(2) LGT

- Wilson action:

$$S_W = \beta \sum_{\square} \left(1 - \frac{1}{2} \text{Re Tr } U_{\square} \right) \approx -\frac{\beta}{8} \sum_x a^4 \text{Tr } F_{\mu\nu}(x) F_{\mu\nu}(x)$$

- $\beta \equiv 4/g^2$
- In limit $a \rightarrow 0$ agrees with

$$S_G = -\frac{1}{2g^2} \int d^4x \text{Tr } F_{\mu\nu} F_{\mu\nu}$$

- Equivalence with 4D statistical mechanics
 - ★ $\beta \leftrightarrow$ “inverse temperature”
 - ★ **Physical temperature** defined by $T \equiv 1/aN_\tau$
- As we will see, there are other possible energy definitions



Taking the continuum limit

- ▶ How does a approach 0?
 - ▶ Renormalization group equation

$$B(g) = a \frac{dg}{da} = -b_0 g^3 - b_1 g^5 + \mathcal{O}(g^7)$$

- ▶ Solve the differential equation:

$$\begin{aligned} a\Lambda_L &= \exp\left(\int^g \frac{dg'}{B(g')}\right) \\ &= \exp\left(-\frac{1}{2b_0 g^2}\right) (b_0 g^2)^{-b_1/2b_0^2} \left(1 + \mathcal{O}(g^2)\right) \\ &\equiv f_{\text{as}}(g^2) \left(1 + \mathcal{O}(g^2)\right) \end{aligned}$$

- ▶ Λ_L integration constant, so this shows how $g \rightarrow 0$ drives $a \rightarrow 0$
- ▶ Hence $\beta \rightarrow \infty$ brings the system to its continuum limit



Scaling in LGT

- ▶ Lattice Λ -parameter

$$\Lambda_L = \lim_{g \rightarrow 0} \frac{1}{a} f_{\text{as}}(g^2)$$

- ▶ $m_k = c_k \Lambda_L$ in continuum limit
- ▶ Ratios of masses **scale** as

$$\frac{m_j}{m_i} = \frac{c_j}{c_i} \left[1 + \mathcal{O}(a^2 \Lambda_L^2) \right]; \quad \text{equivalently} \quad \frac{L_i}{L_j} \approx r_{ij} + k_{ij} a^2 \Lambda_L^2$$

- ▶ $\mathcal{O}(a^2 \Lambda_L^2)$ deviations from constant ratios are **scaling violations**
- ▶ Want reference scale to depend at most weakly on a



SU(2) LGT simulations

- ▶ Calculate expected value of operator O
 - ▶ In QFT expectation values given by

$$\langle O \rangle = \frac{1}{\mathcal{Z}} \int \mathcal{D}\phi O[\phi] e^{-S[\phi]}, \quad \mathcal{Z} \equiv \int \mathcal{D}\phi e^{-S[\phi]}$$

- ▶ An estimator for $\langle O \rangle$ on the lattice is

$$\bar{O} = \frac{1}{N} \sum_{n=1}^N O_n$$

- ▶ **Markov Chain Monte Carlo** basic idea:
 - ▶ Create a **time series** of measurements of O , each of which is calculated using the configuration
 - ▶ Configuration $n + 1$ generated based on configuration n ; usually sample from some distribution
 - ▶ Accept or reject new configuration with probability $e^{-\Delta S}$



SU(2) LGT simulations

- ▶ In practice, configurations n and $n + 1$ are highly correlated
- ▶ Solution:
 - ▶ Update the lattice multiple times between measurements, e.g. measurements on configurations n and $n + 5$
 - ▶ Divide measurements into N blocks, called **repetitions**, and compute average over data in the block
 - ▶ These averages are effectively statistically independent
- ▶ Updating every link on the lattice is called a **sweep**
- ▶ Length scale A is **more efficient** than length scale B if A requires fewer *total* sweeps to achieve the same fractional error bar

QUESTION: Which reference scale is most efficient?



1. Motivation
2. Introduction
3. Meet the Scales
4. Scaling Analysis
5. Conclusion



Deconfining phase transition temperature

- ▶ Remember physical temperature is $T = 1/aN_\tau$
- ▶ **Deconfining phase transition** at T_c in $N_s \rightarrow \infty$ limit
 - ▶ Polyakov loop susceptibility diverges at β_c
 - ▶ “glueballs” for $T < T_c$
 - ▶ “gluon plasma” for $T > T_c$
- ▶ On finite lattices, Polyakov loop susceptibility has maximum at the **pseudocritical coupling** $\beta_c(N_\tau, N_s)$
- ▶ Determine **critical coupling constant** $\beta_c(N_\tau)$ from $\beta_c(N_s, N_\tau)$ using three parameter fit

$$\beta_c(N_s, N_\tau) = \beta_c(N_\tau) + A N_s^{-B}$$

- ▶ Inverting the results defines the deconfining length scale $N_\tau(\beta)$
- ▶ Clear physical meaning, but computationally expensive






Table: Pseudocritical β values N_s : β_c . Error bars of β_c are in parentheses.

| $N_\tau = 4$ | $N_\tau = 6$ | $N_\tau = 8$ |
|--------------------------|-------------------------|-------------------------|
| 08: 2.30859(53) | 12: 2.43900(33) | 16: 2.52960(90) |
| 12: 2.30334(33) | 18: 2.43096(43) | 24: 2.51678(43) |
| 16: 2.30161(30) | 20: 2.42973(11) | 32: 2.51296(20) |
| 20: 2.30085(17) | 24: 2.42873(35) | 40: 2.51192(12) |
| 24: 2.30060(16) | 28: 2.427939(74) | 44: 2.51150(11) |
| 28: 2.30025(19) | 30: 2.427690(87) | 48: 2.51119(11) |
| 32: 2.299754(99) | 36: 2.427274(67) | 52: 2.51130(11) |
| 40: 2.299593(74) | 44: 2.426827(67) | 56: 2.511096(85) |
| 48: 2.299452(83) | 48: 2.426756(64) | 64: 2.510635(83) |
| 56: 2.299435(29) | 56: 2.426605(62) | 72: 2.510716(72) |
| | 60: 2.426596(55) | 80: 2.510517(79) |
| ∞ : 2.299188(61) | ∞ : 2.42636(52) | ∞ : 2.510363(71) |
| $q = 0.56$ | $q = 0.73$ | $q = 0.14$ |
| $N_\tau = 4 \pm 0.00063$ | $N_\tau = 6 \pm 0.0011$ | $N_\tau = 8 \pm 0.0019$ |

Lots of simulations needed for each point!




Energy operators

- ▶ Time dependent plaquettes with parameterization

$$\langle U_{\square}(t) \rangle = a_0(t) \mathbf{1} + i \sum_{i=1}^3 a_i(t) \sigma_i$$

- ▶ Multiple possible energy discretizations, must agree when $a \rightarrow 0$
- ▶ Examined three discretizations for this project:
 - ▶ $E_0(t) \equiv 2[1 - a_0(t)]$
 - ▶ $E_1(t) \equiv \sum_{i=1}^3 a_i(t)^2$
 - ▶ $E_4(t) \equiv \frac{1}{4} \sum_{i=1}^3 (a_i^{(1)} + a_i^{(2)} + a_i^{(3)} + a_i^{(4)})^2$
 - ▶ E_4 suggested by Lüscher [3]



QUESTION: Is there any scaling advantage to using E_4 ?



Gradient flow

- ▶ Lüscher's gradient flow equation [3]

$$\dot{V}_\mu(x, t) = -g^2 V_\mu(x, t) \partial_{x, \mu} S[V(t)]$$

- ▶ **Flow time** t and initial condition $V_\mu(x, t)|_{t=0} = U_\mu(x)$
 - ▶ Gradient flow equation drives action down
- ▶ Given **target value** y , length scale $s(\beta) = \sqrt{t_y(\beta)}$ defined implicitly via equation


$$t^2 \langle E_t \rangle |_{t=t_y} = y$$

- ▶ Requires no fits or extrapolations
- ▶ And at least 100 times more efficient than $N_\tau(\beta)$
- ▶ But no obvious physical meaning
- ▶ And some ambiguity in choosing y : $2 \times 3 = 6$ targets per scale

QUESTION: Does choosing one target over another result in seriously distinct scaling behavior?



Determination of gradient scale



Determination of target value




Table: Gradient length scales using energy operators E_0 , E_1 , E_4 and target values $y_0 = 0.0376$, $y_1 = 0.0370$, $y_4 = 0.0300$, respectively

| β | Lattice | $L_1 = s_0$ | $L_2 = s_1$ | $L_3 = s_4$ |
|---------|---------|-------------|-------------|-------------|
| 2.3 | 16^4 | 1.3593(28) | 1.3589(27) | 1.2756(75) |
| 2.43 | 28^4 | 2.1023(30) | 2.0911(30) | 1.9666(98) |
| 2.51 | 28^4 | 2.7590(73) | 2.7428(73) | 2.570(14) |
| 2.574 | 40^4 | 3.4103(72) | 3.3896(71) | 3.149(16) |
| 2.62 | 40^4 | 3.954(10) | 3.9293(99) | 3.672(19) |
| 2.67 | 40^4 | 4.622(17) | 4.593(17) | 4.297(24) |
| 2.71 | 40^4 | 5.199(22) | 5.167(22) | 4.817(27) |
| 2.751 | 40^4 | 5.909(34) | 5.872(34) | 5.457(41) |
| 2.816 | 44^4 | 7.092(48) | 7.049(47) | 6.530(54) |
| 2.875 | 52^4 | 8.510(64) | 8.456(65) | 7.883(68) |

Each scale needs only one lattice, access large β !



Cooling flow

- ▶ Introduced by Berg [4] for $O(3)$ topological charge and used by Bonati and D'Elia [5] for topological observables in $SU(3)$
- ▶ Iterative process
 - ▶ Instead of a differential equation, just replace link variable with one that locally minimizes action

$$V_\mu(x, n_c) = \frac{V_\mu^\sqcup(x, n_c - 1)}{|V_\mu^\sqcup(x, n_c - 1)|}$$

- ▶ n_c cooling sweeps corresponds to a gradient flow time [5]

$$t_c = n_c/3$$

- ▶ Length scale $x(\beta) = \sqrt{t_y(\beta)}$ defined implicitly like with the gradient flow

$$t^2 \langle E_t \rangle |_{t=t_y} = y$$

- ▶ Same advantages and disadvantages as the gradient flow
- ▶ And at least 34 times faster

QUESTION: Does the cooling flow experience significantly larger scaling violations than the gradient flow?



Table: Cooling length scales using energy operators E_0 , E_1 , E_4 and target values $y_0 = 0.044$, $y_1 = 0.043$, $y_4 = 0.035$, respectively

| β | Lattice | $L_7 = x_0$ | $L_8 = x_1$ | $L_9 = x_4$ |
|---------|---------|-------------|-------------|-------------|
| 2.3 | 16^4 | 1.3433(24) | 1.3385(23) | 1.2575(74) |
| 2.43 | 28^4 | 2.0892(28) | 2.0707(28) | 1.9446(95) |
| 2.51 | 28^4 | 2.7522(68) | 2.7267(66) | 2.548(15) |
| 2.574 | 40^4 | 3.4048(69) | 3.3730(67) | 3.137(17) |
| 2.62 | 40^4 | 3.9509(95) | 3.9137(93) | 3.645(22) |
| 2.67 | 40^4 | 4.618(17) | 4.574(16) | 4.298(26) |
| 2.71 | 40^4 | 5.203(21) | 5.154(21) | 4.794(28) |
| 2.751 | 40^4 | 5.913(32) | 5.857(32) | 5.434(40) |
| 2.816 | 44^4 | 7.105(45) | 7.039(45) | 6.511(55) |
| 2.875 | 52^4 | 8.514(60) | 8.433(59) | 7.825(68) |

Must check its scaling behavior



1. Motivation
2. Introduction
3. Meet the Scales
4. **Scaling Analysis**
5. Conclusion



Scaling analysis


- Recall scaling behavior given by

$$R_{ij} \equiv \frac{L_i}{L_j} \approx r_{ij} + k_{ij} a^2 \Lambda_L^2 = r_{ij} + c_{ij} \left(\frac{1}{L_j} \right)^2$$


- Collect all length scales from the project
 - $L_1 - L_3$: gradient with first set of target values
 - $L_4 - L_6$: gradient with second set of target values
 - $L_7 - L_9$: cooling with first set of target values
 - $L_{10} - L_{12}$: cooling with second set of target values
- In the following we fix $L_j = L_{10}$ and plot

$$\frac{R_{i,10}}{r_{i,10}} = 1 + c'_{i,10} \left(\frac{1}{L_{10}} \right)^2$$






Closer look at lengths using E_0 and E_1



Closer look at lengths using E_4



1. Motivation
2. Introduction
3. Meet the Scales
4. Scaling Analysis
5. Conclusion



Conclusions






- ▶ Does choosing different target values for the gradient and cooling flows lead to seriously distinct scaling behavior?
 - ▶ Not if using physical input to guide initial scaling values
 - ▶ Can use the deconfinement scale
- ▶ Is there any scaling advantage to using E_4 ?
 - ▶ None were observed
 - ▶ Estimates from the same statistics have smaller error bars using E_0 or E_1 than with E_4
- ▶ Does the cooling length experience significant scaling violations compared to the deconfining and gradient scales?
 - ▶ No noticeable loss of accuracy using cooling flow
- ▶ Which scale is most efficient?
 - ▶ Gradient scale at least 100 times more efficient than deconfining
 - ▶ Cooling at least 34 times more efficient than gradient



Thank you!



References

-  E. Follana *et al.*, “High-Precision Determination of the α_s , K , D , and D^* Decay Constants from Lattice QCD”, Physical review letters, **100** 062002 (2008).
-  C. T. H. Davies *et al.*, “Update: Precision D^* decay constant from full lattice QCD using very fine lattices”, Physical Review D, **82** (2010).
-  M. Lüscher, “Properties and uses of the Wilson flow in lattice QCD”, Journal of High Energy Physics, **2010** 1 (2010).
-  B. Berg, “Dislocations and topological background in the lattice $O(3)$ sigma model”, Physics Letters, **104B** 475 (1981).
-  C. Bonati and M. D’Elia, “Comparison of the gradient flow with cooling in $SU(3)$ pure gauge theory”, Physical Review D, **89** 105005 (2014).



Backup slides

- Deconfining scale error bars

$$\Delta N_\tau = \frac{N_\tau}{L_{10}^{1,3}(\beta_c)} \left[L_{10}^{1,3}(\beta_c) + L_{10}^{1,3}(\beta_c - \Delta\beta_c) \right]$$

- **Polyakov loop** definition $P(\vec{x}) = \prod U_\tau(\vec{x})$
- Polyakov loop susceptibility

$$\chi(\beta) = \langle P^2 \rangle - \langle |P| \rangle^2, \quad \text{where} \quad P = \frac{1}{N_s^3} \sum_{\vec{x}} P(\vec{x})$$

