

CS6033 Assign No.8

Minghe Yang

May 2021

1 Karger-Stein's Algorithm

1.1




Figure 1: Binary Tree

1.2

Firstly, $M_{0,j} = \prod_{l=0}^0 = m_{j+1} = m_j$.

Secondly, $M_{i+1,j} = \prod_{l=0}^{2^{i+1}-1} = m_{j2^{i+1}+l} = M_{i,2j}(\prod_{l=0}^{2^i-1} m_{(2j+1)2^i+l}) = M_{i,2j}M_{i,2j+1}$.

1.3

$M_{i,j}$ should be on i_{th} layer from the bottom, and j_{th} node from left.

1.4.a

Algorithm 1 Subproduct

Input: $n - 2^k, u_0, u_1, \dots, u_{n-1}$

Output: $M_{i,j}$

```

1: for  $i \in (0, n - 1)$  do
2:    $M_{0,i} \leftarrow X - u_i$ 
3: end for
4: for  $i \in (i, k)$  do
5:   for  $j \in (0, 2^{k-i} - 1)$  do
6:      $M_{i,j} \leftarrow M_{i-1,2j}M_{i-1,2j+1}$ 
7:   end for
8: end for
```

1.4.b

Algorithm 2 Fast Multipoint Evaluation

Input: $P, n - 2^k, u_0, u_1, \dots, u_{n-1}, M_{i,j}$

Output: Evaluation of $\{Pu_0, Pu_1, \dots, P(u_{n-1})\}$

```

1: function DD( $f, k, i$ )
2:   if  $n = 1$  then
3:     return  $f$ 
4:   end if
5:    $fleft \leftarrow f \bmod M_{k-1,2i}$ 
6:    $fright \leftarrow f \bmod M_{k-1,2i+1}$ 
7:    $lset \leftarrow DD(fleft, k - 1, 2i)$ 
8:    $rset \leftarrow DD(fright, k - 1, 2i + 1)$ 
9:   return  $lset, rset$ 
10: end function
11: return DD( $P, k, 0$ )
```

1.5.a

One layer means that f is just a constant, so returning f is true. Suppose that is correct in $k-1$ layer, here comes k th.

So $P = qlleft(u_i)M_{k-1,0} + rleft(u_i)$, $P = qright(u_i)M_{k-1,0} + rright(u_i)$. qM term will be 0, so the true value will be in the r term, which is evaluated in $k-1$ layer. So it is correct.

1.5.b

From solution above, we can get that $T(n) = 2T(n/2) + M(n)$, so the complexity should be $O(M(n) \log n)$.

2 Part Two

2.2

$m' = \sum_{i=0}^{n-1} (x - u_i)' \frac{m}{x - u_i} = \sum_{i=0}^{n-1} \frac{m}{X - u_i}$, Since all terms with $X - u_i$ is 0, only term remain will be $1/s_i$

2.3

Algorithm 3 Fast Evaluation

Input: $n - 2^k, u_0, u_1, \dots, u_{n-1}, \{P(u_0), P(u_1), \dots, P(u_{n-1})\}, M_{i,j}$

Output: P

```

1: function MULTUP( $f, k, i$ )
2:   if  $n = 1$  then
3:     return  $y$ 
4:   end if
5:    $fleft \leftarrow Multup(fleft, k - 1, 2i)$ 
6:    $fright \leftarrow Multup(frigh, k - 1, 2i + 1)$ 
7:   return  $fleft \times M_{k,1}, fright \times M_{k,0}$ 
8: end function
9: return  $Multup(P, k, 0)$ 

```

2.5

It might be correct, while needs much space for calculation.

3 Critical Thinking

2.2

From the first line we can get, $x \times y$ is not equal to two primes numbers, and have two or more complexities.

Form the second part we can get, $x + y$ is odd, and also have two or more combinations.

From Goldbach conjecture, all even numbers will be excluded from $x + y$, so all possible values will be 11, 17, 23, 29, 37, 41, 47, 51, 53, 57, 59, 65, 67, 71, 77, 79, 83, 87, 89, 93, 95, 97. Number like 27 and 35 should not be in the list, since it can be dived into to prime numbers. Also, if S2 know exact answer, the S1 should delete one wrong possibility (only one). The correspond will be : 18 - 11 24 - 11 28 - 11 50 - 27 52 - 17 54 - 29 76 - 23 92 - 27 96 - 35 98 - 51 100 - 29 112 - 23 124 - 35 140 - 27 144 - 51.

From 52 - 17, we can get the correct answer(like if your get 11, there are three combinations, you can't judge, so only one combination is preferred), means the number is 13 and 4.