

L12 Advanced DNA Assembly

Instructor: Prof. K. Solomon Ph.D.
Assistant Professor
Agricultural & Biological Engineering
Laboratory for Renewable
Resources Engineering

Fall 2018



Recall...

Required components for DNA delivery

4 mechanisms to assemble DNA

1. Chemical addition
2. Homologous recombination
3. Annealing & Polymerization
4. Digestion & Ligation

BioBrick™ Std

Advantages	Disadvantages
<ul style="list-style-type: none"> - idempotent/scalable - Easily automatable - Robust - Only 4 REs used 	<ul style="list-style-type: none"> - Need to remove internal Biobrick RE sites from all parts - Have a fixed scar (not good for protein fusions) - Can't add parts between parts

2


This lecture....

- Scarless digestion + ligation
- Anneal + polymerization methods
- Recombination-based methods
- Summary


3

Fusion proteins are common in biotech

Scars between parts may cause frameshift mutations or disrupt folding (function)




thermofisher.com



parts.igem.org

4

Type IIS restriction enzymes allow you to define overhang (and ligation scars)



Type IIS REs:

- Cleave outside their restriction site
- Are not palindromic
- Can give user-defined sticky ends

Cuts are directional


thermofisher.com

5

Golden Gate Cloning

GGTCTC NNNNNN
CCAGAG NNNNNN

Uses BsaI




Restriction site removed after digestion

Process is scalable – can be repeated on composite parts without digesting components

6

Golden Gate

Primers used to define overhangs for scarless fusions



7

Golden Gate

GGTCTC NNNNNN
CCAGAG NNNNNN

Cleavage site is user-defined (specified by position relative to recognition site)


- Can specify linker for fusion proteins
- Can do multi-part assembly with 1 enzyme simultaneously
 - Assembly specified with unique complementary overhangs for each junction
- Cloning is simple and **fast**
 - Can do digestion + ligation simultaneously (one-pot) ~30min
 - Re-ligation of backbone quickly digested because site is preserved

8

Golden Gate

Real Example: Modular Cloning (MoClo)

- Parts constructed in individual plasmids
- Parts flanked by BsaI site with complementary, user-defined overhangs




Weber, E. et al. (2011). *PLoS ONE*.

9

Golden Gate

Real Example: Modular Cloning (MoClo)

- Parts constructed in individual plasmids
- Parts flanked by BsaI site with complementary, user-defined overhangs
- Digestion and ligation result in assembly of multi-part constructs



Cloning is simple and fast


- Can do digestion + ligation simultaneously (one-pot) ~30min
- Re-ligation of backbone quickly digested because site is preserved

Weber, E. et al. (2011). *PLoS ONE*.

10

Golden Gate (MoClo)

Formatted part libraries



Lee, ME. et al. (2015). *ACS Synthetic Biology*.

11

Golden Gate

Advantages

- Simple and Fast
- Scarless (good for fusions)
- Good for libraries (efficient)
- Only one RE
- Up to 10 parts assembled in a single one pot reaction without purification

Disadvantages

- Parts need to be in appropriate format + no internal sites
- Overhangs must be unique to prevent misassembly

12

Sequence-independent cloning methods

RE-based assembly requires part formatting that may be difficult

- Overhangs, no internal restriction sites

POLYMERIZATION + ANNEAL

1. Gibson Chew Back Anneal (CBA)
2. Sequence + Ligase Independent Cloning (SLIC)
3. Splicing by Overlap Extension PCR (SOEing)

13

Gibson Assembly

Requires:

- Linear DNA with long regions of homology (~20 – 50 bp) @ junctions
- 3 enzymes:

1. T5 Exonuclease
Chews back 5' end of DNA to make 3' overhang
2. Phusion/Proof-reading DNA polymerase
Repairs the gaps in overhangs via PCR
3. Taq DNA Ligase
Forms covalent bonds between parts




Enzymes chosen to work together in one pot at one temperature

14

Gibson Assembly

Workflow




Gibson Assembly

Advantages	Disadvantages
<ul style="list-style-type: none"> • Parts don't need to be formatted (sequence independent) • Can make combinatorial libraries (up to 5 parts) • One-pot reaction (fast/simple) • Scarless 	<ul style="list-style-type: none"> • Requires long primers that may be challenging • Expensive

16

Sequence + Ligation Independent Cloning (SLIC)

- Like Gibson but no ligase or dNTPs to repair chew back
 - T4 DNA polymerase has 3' exonuclease activity in absence of dNTPs
 - dCTPs added to control amount of chew back
- Repair completed in host



Advantages	Disadvantages
• Cheaper	• Less efficient (bad for libraries)

17

Splicing by Overlap Extension PCR (SOEing)


Uses PCR to “stitch” together DNA

- PCR primers generate homology or adjacent parts
- Requires multiple reactions and only yield **linear** DNA

1 – Amplify parts separately



2 – Purify and then anneal products (melt + cool slowly)



3 – Products serve as “mega-primers” for each other


Extend/Amplify ~5 cycles to make complete product



18

SOEing → CPEC

- Can do similar procedure with a vector + parts (no ligase needed) = **Circular Polymerase Extension Cloning (CPEC)**
- Using just overlapping primers is how genes are synthesized



JBEI.org
19

SOEing Limitations

- Large stretches of ssDNA
 - Must avoid forming stable secondary structures and repeats
 - Can lead to assembly failures or mutations at junctions

But potentially, cheaper → only need primers and polymerase

20

Recombination-based methods: Gateway Cloning





Figure 1: Lambda phage integration and excision reactions. Recombination of attP and attB sites creates attL and attR sites.


Addgene.org
21

Recombination-based methods: Gateway Cloning

1 BP Reaction



2 LR Reaction



Addgene.org
22

Gateway Cloning

Advantages	Disadvantages
<ul style="list-style-type: none"> Fast Can make combinatorial libraries (up to 4 parts) Efficient 	<ul style="list-style-type: none"> Fixed vector choices

23

Next time....

- How to 'format' parts for assembly
- Practice problems

24