

L5 Part Libraries and Qualitative Properties

Instructor: Prof. K. Solomon Ph.D.
Assistant Professor
Agricultural & Biological Engineering
Laboratory for Renewable
Resources Engineering

Fall 2018



Recall...

Parts, devices and systems can be assembled in modular fashion with designed input/output for a function

Chassis → Systems → Devices → Parts → DNA

What properties must be true for a design to “work” as intended?

2

This lecture....

- Recap IO relationship and physical design
- Properties required for function
- Part libraries



3

Bacterial Photography




If no light, produce pigment

*What logic gates are involved?
How is this biochemically implemented?*




Tabor et al, Cell, 2009



ci added to tune system properties

- ci* repressor prevents **leaky** (non-zero background) expression, negates effects of low concentrations of AHL



- System is more sensitive to AHL (log vs linear scales)
- luxI* is catalytic, AHL will accumulate and ultimately overpower *ci*



6

To ensure proper functioning, parts must be

- Impedance matched
 - IO ranges of adjacent parts must be compatible for maximum signal transfer
- Independent
 - Parts that perform the same distinct function in isolation and in interaction with other systems. These are typically reliable and can be combined with other systems (composable)
- Orthogonal
 - Parts that do not cross talk or cross react

7

Impedance matching




8

Independence

Parts have distinct functions that allow use in multiple contexts without interaction

- E.g. LacO (O1), TetO (O2), and AraO (ara1/ara2) are distinct
- P_{lac} /lacI, P_{BAD} /AraC, P_{Tet} /TetR can be used simultaneously or combined to make chimeric promoters (**composable**)
- Components don't compete for binding across systems/don't titrate components (fixed IO, **reliable**)




Lutz and Bujard, NAR, 1997

Orthogonality

Parts do not cross-talk or cross react

T7 Promoter (~18 bps) – requires T7 RNAP

5' TAATACGACTCACTATAG 3'



10

Part libraries

We enforce these properties and create a variety of input/output relationships with engineered part libraries (see <http://parts.igem.org>)

- Promoter engineering
- Machinery from different organisms (e.g. T7, tetON/tetOFF)
- Directed evolution

Apart from canonical parts (promoters, etc), libraries come in other forms such as RNA, RNAP, ribosomes, etc

11



RNA Parts

are specific to individual genes (orthogonal) as they rely on RNA:RNA intxns

Operate in:

i) *Cis* – gene & regulatory RNA are on same transcript

e.g. B12 riboswitches
Transcriptional attenuation



Wikipedia.org


12

RNA parts

Operate in:


- ii) *Trans* - gene and regulatory RNA on separate transcripts

e.g. siRNA, trans activating RNA, antisense RNA



13

Selection & Directed Evolution can be used to make RNA parts that bind (SELEX – systematic evolution of ligands by exponential enrichment)




14

RNA interactions can be modeled

Computational design can be used to predict RNA properties via thermodynamics

- Vienna, NuPACK, Mfold
- Determine minimum free energy structure and/or likeliest structure

These tools only predict the final structure, which may or may not be relevant



15

Genome mining

Can look to nature for natural solutions



e.g. B12 switch – can reuse that aptamer in other contexts

All these approaches (selection and directed evolution, computational design, and mining) can be used for any type of part library

16

Orthogonal ribosomes


- Ribosomes that recognize non-canonical RBS
- +ve and -ve screens used to identify functional mutants via directed evolution
- O-ribosomes are independent, and can't be titrated away (fixed I/O)



Rackham &
Chin, *Nat Chem*
Biol. 2005

Orthogonal transcriptional machinery


- Use viral trxn machinery that is host independent (e.g. T7) – no titration
- Can use sigma factors from different organisms

Rhodijs et al. *MSB* 2013, **11**:12

1


CRISPRi/a

RNA guided technologies can be used to target individual genes for expression through presence/absence of guide (orthogonality)



CRISPRi
Qi et al, Cell, 2013

CRISPRa
Bikard et al, NAR, 2013



16

Next time....

- What are the quantitative properties that are needed for function?

3