
CHARACTER THEORY OF FINITE GROUPS — EXERCISE SHEET 4

JUN.-PROF. DR. CAROLINE LASSUEUR

Due date: Thursday 06.06.2017, 12:00

TU KAISERSLAUTERN

BERNHARD BÖHMLER

SS 2019

Throughout this exercise sheet all vector spaces are assumed to be finite-dimensional. Each Exercise is worth 4 points.

EXERCISE 13

Let G and H be finite groups. Let V and W be two \mathbb{C} -vector spaces.

- Assume V is a G -vector space and W is an H -vector space. Prove that $V \otimes_{\mathbb{C}} W$ is a $G \times H$ -vector space for the action defined by $(g, h).(v \otimes w) = g.v \otimes h.w \forall g \in G, h \in H, v \in V, w \in W$ with corresponding representation $\rho_{V \otimes_{\mathbb{C}} W} = \rho_V \otimes \rho_W$.
- Let $R : G \longrightarrow \mathrm{GL}_n(\mathbb{C})$, $S : H \longrightarrow \mathrm{GL}_m(\mathbb{C})$ be matrix representations. Prove that $R \otimes S$ is a matrix representation of $G \times H$.

EXERCISE 14

Let G and H be finite groups. Prove the following assertions:

- $\mathrm{Irr}(G \times H) = \{\chi \cdot \psi \mid \chi \in \mathrm{Irr}(G), \psi \in \mathrm{Irr}(H)\}$.
- $\{\chi \in \mathrm{Irr}(G) \mid \chi(1) = 1\}$ is a group for the multiplication of characters.
- If $\chi, \lambda \in \mathrm{Irr}(G)$ and λ is linear, then $\chi \cdot \lambda \in \mathrm{Irr}(G)$.

EXERCISE 15

Compute the complex character table of the alternating group \mathfrak{A}_4 through the following steps:

- Determine the conjugacy classes of \mathfrak{A}_4 (there are 4 of them) and the corresponding centraliser orders.
- Determine the degrees of the 4 irreducible characters of \mathfrak{A}_4 .
- Determine the linear characters of \mathfrak{A}_4 .
- Determine the non-linear character of \mathfrak{A}_4 using the 2nd Orthogonality Relations.

EXERCISE 16

Let G be a finite group. Prove the following assertions:

- If $N \trianglelefteq G$, then:

$$N = \bigcap_{\substack{\chi \in \mathrm{Irr}(G) \\ N \subseteq \ker(\chi)}} \ker(\chi)$$

- G is a simple group if and only if $\chi(g) \neq \chi(1) \forall \chi \in \mathrm{Irr}(G) \setminus \{1_G\}$, and $\forall g \in G \setminus \{1\}$.