

EXAMPLE

THE DEGREES OF THE IRREDUCIBLE
CHARACTERS OF $GL_3(2)$

EXAMPLE : THE DEGREES OF THE IRREDUCIBLE CHARACTERS OF $GL_3(2)$

$$G := GL_3(\mathbb{F}_2)$$

EXAMPLE : THE DEGREES OF THE IRREDUCIBLE CHARACTERS OF $GL_3(2)$

$G := GL_3(\mathbb{F}_2)$ simple group of order

$$\# \mathbb{F}_2\text{-bases of } \mathbb{F}_2^3 = (2^3 - 1)(2^3 - 2)(2^3 - 2^2) = 7 \cdot 6 \cdot 4 = 2^3 \cdot 3 \cdot 7 = 168 .$$

EXAMPLE : THE DEGREES OF THE IRREDUCIBLE CHARACTERS OF $GL_3(2)$

$G := GL_3(\mathbb{F}_2)$ simple group of order

$$\#\mathbb{F}_2\text{-bases of } \mathbb{F}_2^3 = (2^3 - 1)(2^3 - 2)(2^3 - 2^2) = 7 \cdot 6 \cdot 4 = 2^3 \cdot 3 \cdot 7 = 168 .$$

We accept here that :

- G is simple
- G has 6 conjugacy classes

EXAMPLE : THE DEGREES OF THE IRREDUCIBLE CHARACTERS OF $GL_3(2)$

$G := GL_3(\mathbb{F}_2)$ simple group of order

$$\#\mathbb{F}_2\text{-bases of } \mathbb{F}_2^3 = (2^3-1)(2^3-2)(2^3-2^2) = 7 \cdot 6 \cdot 4 = 2^3 \cdot 3 \cdot 7 = 168 .$$

We accept here that : • G is simple
• G has 6 conjugacy classes $\Rightarrow |\text{Irr}(G)| = 6$

EXAMPLE : THE DEGREES OF THE IRREDUCIBLE CHARACTERS OF $GL_3(2)$

$G := GL_3(\mathbb{F}_2)$ simple group of order

$$\#\mathbb{F}_2\text{-bases of } \mathbb{F}_2^3 = (2^3-1)(2^3-2)(2^3-2^2) = 7 \cdot 6 \cdot 4 = 2^3 \cdot 3 \cdot 7 = 168 .$$

We accept here that :

- G is simple
- G has 6 conjugacy classes $\Rightarrow |\text{Irr}(G)| = 6$

CLAIM : $x(1)$ is determined by this data $\forall x \in \text{Irr}(G)$.

EXAMPLE : THE DEGREES OF THE IRREDUCIBLE CHARACTERS OF $GL_3(2)$

$G := GL_3(\mathbb{F}_2)$ simple group of order

$$\#\mathbb{F}_2\text{-bases of } \mathbb{F}_2^3 = (2^3-1)(2^3-2)(2^3-2^2) = 7 \cdot 6 \cdot 4 = 2^3 \cdot 3 \cdot 7 = 168 .$$

We accept here that :

- G is simple
- G has 6 conjugacy classes $\Rightarrow |\text{Irr}(G)| = 6$

CLAIM : $x(1)$ is determined by this data $\forall x \in \text{Irr}(G)$.

STEP 1 : The degree formula yields

EXAMPLE : THE DEGREES OF THE IRREDUCIBLE CHARACTERS OF $GL_3(2)$

$G := GL_3(\mathbb{F}_2)$ simple group of order

$$\#\mathbb{F}_2\text{-bases of } \mathbb{F}_2^3 = (2^3 - 1)(2^3 - 2)(2^3 - 2^2) = 7 \cdot 6 \cdot 4 = 2^3 \cdot 3 \cdot 7 = 168 .$$

We accept here that :

- G is simple
- G has 6 conjugacy classes $\Rightarrow |\text{Irr}(G)| = 6$

CLAIM : $\chi(1)$ is determined by this data $\forall \chi \in \text{Irr}(G)$.

STEP 1 : The degree formula yields

$$168 = |G| = \sum_{i=1}^6 \chi_i(1)^2 = 1 + \sum_{i=2}^6 \chi_i(1)^2$$

STEP 2. Linear characters :

STEP 2. Linear characters: $G = G'$ as G is non-abelian simple

STEP 2. Linear characters: $G = G'$ as G is non-abelian simple
 \implies there are $|G : G'| = 1$ linear characters.

STEP 2. Linear characters: $G = G'$ as G is non-abelian simple
 \Rightarrow there are $|G : G'| = 1$ linear characters.

$$\Rightarrow \chi_i(1) \geq 2 \quad \forall 2 \leq i \leq 6$$

STEP 2. Linear characters: $G = G'$ as G is non-abelian simple
 \Rightarrow there are $|G : G'| = 1$ linear characters.

$$\Rightarrow \chi_i(1) \geq 2 \quad \forall 2 \leq i \leq 6$$

So, we have the following 5 possibilities for the degrees:

$\chi_1(1)$	$\chi_2(1)$	$\chi_3(1)$	$\chi_4(1)$	$\chi_5(1)$	$\chi_6(1)$
1	2	4	5	6	9
1	2	3	3	8	9
1	2	5	5	7	8
1	2	4	7	7	7
1	3	3	7	7	8

STEP 3. Use integrality! In part. $x_i(1) \mid |G| = 168 = 2^3 \cdot 3 \cdot 7 \quad \forall 1 \leq i \leq 6$:

STEP 3. Use integrality! In part. $\chi_i(1) \mid |G| = 168 = 2^3 \cdot 3 \cdot 7 \quad \forall 1 \leq i \leq 6$:

	$\chi_1(1)$	$\chi_2(1)$	$\chi_3(1)$	$\chi_4(1)$	$\chi_5(1)$	$\chi_6(1)$
1	2	4	5	6	8	8
1	2	3	3	8	8	8
1	2	5	5	7	8	7
1	2	4	7	7	7	7
1	3	3	7	7	8	

STEP 3. Use integrality! In part. $x_i(1) \mid |G| = 168 = 2^3 \cdot 3 \cdot 7 \quad \forall 1 \leq i \leq 6$:

	$\chi_1(1)$	$\chi_2(1)$	$\chi_3(1)$	$\chi_4(1)$	$\chi_5(1)$	$\chi_6(1)$
1	2	4	5	6	8	X
1	2	3	3	8	8	X
1	2	5	5	7	8	
1	2	4	7	7	7	
1	3	3	7	7	8	

} remains
2 possibilities

STEP 3. Use integrality! In part. $x_i(1) \mid |G| = 168 = 2^3 \cdot 3 \cdot 7 \quad \forall 1 \leq i \leq 6$:

	$\chi_1(1)$	$\chi_2(1)$	$\chi_3(1)$	$\chi_4(1)$	$\chi_5(1)$	$\chi_6(1)$
1	2	4	5	6	8	X
1	2	3	3	8	8	X
1	2	5	5	7	8	
1	2	4	7	7	7	
1	3	3	7	7	8	

{ remains
2 possibilities

STEP 4. Eliminate 4th possibility

STEP 3. Use integrality! In part. $\chi_i(1) \mid |G| = 168 = 2^3 \cdot 3 \cdot 7 \quad \forall 1 \leq i \leq 6$:

	$\chi_1(1)$	$\chi_2(1)$	$\chi_3(1)$	$\chi_4(1)$	$\chi_5(1)$	$\chi_6(1)$
1	2	4	5	6	8	X
1	2	3	3	8	8	X
1	2	X	X	7	8	
1	X	4	7	7	7	
1	3	3	7	7	8	

} remains 2 possibilities

STEP 4. Eliminate 4th possibility:

[Ex. 21, Sheet 6] $\Rightarrow \chi_2(1) \neq 2$ as G is simple.

STEP 3. Use integrality! In part. $x_i(1) \mid |G| = 168 = 2^3 \cdot 3 \cdot 7 \quad \forall 1 \leq i \leq 6$:

	$\chi_1(1)$	$\chi_2(1)$	$\chi_3(1)$	$\chi_4(1)$	$\chi_5(1)$	$\chi_6(1)$
1	2	4	5	6	8	X
1	2	3	3	8	8	X
1	2	X	X	7	8	
1	X	4	7	7	7	
1	3	3	7	7	8	{ remains 2 possibilities}

STEP 4. Eliminate 4th possibility:

[Ex. 21, Sheet 6] $\Rightarrow \chi_2(1) \neq 2$ as G is simple.

CONCLUSION: the degrees of the irreducible characters are

$$1 \quad 3 \quad 3 \quad 7 \quad 7 \quad 8$$