



Patient: **JENNIFER
HAO**
DOB: February 15, 1994
Sex: F
MRN: 0003231386


Order Number: **V8060181**
Reported: June 18, 2026
Received: June 06, 2026
Collected: June 04, 2026

UCSF Health
Max Brondfield

2200 GI Effects® Comprehensive Profile - Stool

Powered by **Genova AI**

Results Overview




Functional Imbalance Scores

Key **< 2** : Low Need for Support **2-3** : Optional Need for Support **4-6** : Moderate Need for Support **7-10** : High Need for Support

	Need for Digestive Support	Need for Inflammation Modulation	Need for Microbiome Support	Need for Prebiotic Support	Need for Antimicrobial Support
	MALDIGESTION	INFLAMMATION	DYSBIOSIS	METABOLIC IMBALANCE	INFECTION
	0	2	6	3	8
Biomarkers	Pancreatic Elastase Products of Protein Breakdown Fecal Fats	Eosinophil Protein X Secretory IgA Calprotectin Occult Blood	PP Bacteria/Yeast Reference Variance IAD/Methane Score Total Abundance	SCFA (%) Beta-glucuronidase n-Butyrate Conc. Total SCFA's	Pathogenic Bacteria PP Bacteria/Yeast Parasitic Infection Total Abundance
Therapeutic Support Options	Digestive Enzymes Betaine HCl Bile Salts Apple Cider Vinegar Mindful Eating Habits Digestive Bitters	Elimination Diet/ Food Sensitivity Testing Mucosa Support: Slippery Elm, Althea, Aloe, DGL, etc. Zinc Carnosine L-Glutamine Quercetin Turmeric Omega-3's GI Referral (If Calpro is Elevated)	Pre-/Probiotics Increase Dietary Fiber Intake Consider SIBO Testing Increase Resistant Starches Increase Fermented Foods Meal Timing	Pre-/Probiotics Increased Dietary Fiber Intake Increase Resistant Starches Increase Fermented Foods Calcium D-Glucarate (for high beta-glucuronidase)	Antibiotics (if warranted) Antimicrobial Herbal Therapy Antiparasitic Herbal Therapy (if warranted) <i>Saccharomyces boulardii</i>


Commensal Microbiome Analysis

Commensal Abundance



Total Commensal Abundance: The total commensal abundance is a sum-total of the reported commensal bacteria compared to a healthy cohort. Low levels of commensal bacteria are often observed after antimicrobial therapy, or in diets lacking fiber and/or prebiotic-rich foods and may indicate the need for microbiome support. Conversely, higher total commensal abundance may indicate potential bacteria overgrowth or probiotic supplementation.

Dysbiosis Patterns




Dysbiosis Patterns: Genova's data analysis has led to the development of unique dysbiosis patterns, related to key physiologic disruptions, such as immunosuppression and inflammation. These patterns may represent dysbiotic changes that could pose clinical significance. Please see Genova's published literature for more details: <https://rdcu.be/bRhzv>

Zone 1: The commensal profile in this zone does not align with profiles associated with intestinal inflammation or immunosuppression. If inflammatory biomarkers are present, other causes need to be excluded, such as infection, food allergy, or more serious pathology.

Zone 2: This pattern of bacteria is associated with impaired intestinal barrier function (low fecal sIgA and EPX). Patients in this zone have higher rates of opportunistic infections (e.g. *Blastocystis spp.* & *Dientamoeba fragilis*) as well as fecal fat malabsorption. Commensal abundance is higher in this group suggesting potential bacterial overgrowth.


Zone 3: Patients in this zone may have more inflammation compared to those in zone 4. However, commensal abundance is usually higher making use of antimicrobial therapy relatively safer. Patients in this zone may have higher rates of pathogenic infections.

Zone 4: This commensal profile is associated with increased intestinal inflammation. IBD patients are more likely to have this pattern of bacteria. Commensal abundance is lower in this zone; therefore, antibiotic use for GI potential pathogens should be used with caution. In addition to standard treatment for intestinal inflammation, modulation of the commensal gut profile is encouraged.




Commensal Microbiome Analysis

Commensal Balance



Relative Commensal Abundance



Relative Abundance: The relative abundance compares the quantity of each of 7 major bacterial phyla to a healthy cohort. This can indicate broader variances in the patient's gut microbiome profile. Certain interventions may promote or limit individual phyla when clinically appropriate. Please refer to Genova's Stool Testing Support Guide for more information on modulation of commensal bacteria through diet & nutrient interventions. ***Approximately 70% of the healthy cohort had below detectable levels of *Methanobrevibacter smithii*. Approximately 90% of the healthy cohort had below detectable levels of *Fusobacterium* spp.

Physician Notes/Recommendations

2200 GI Effects® Comprehensive Profile - Stool

Methodologies: GC-FID, Automated Chemistry, ELISA		QUINTILE DISTRIBUTION					Reference Range
Result		1st	2nd	3rd	4th	5th	
Digestion and Absorption							
Pancreatic Elastase 1 †	>500						>200 mcg/g
Products of Protein Breakdown (Total*) (Valerate, Isobutyrate, Isovalerate)	3.3						1.8-9.9 micromol/g
Fecal Fat (Total*)	10.5						3.2-38.6 mg/g
Triglycerides	0.4						0.3-2.8 mg/g
Long-Chain Fatty Acids	7.6						1.2-29.1 mg/g
Cholesterol	2.2						0.4-4.8 mg/g
Phospholipids	0.3						0.2-6.9 mg/g
Inflammation and Immunology							
Calprotectin † ♦	<27						<50 mcg/g
Eosinophil Protein X (EPX) †	1.4						<=2.7 mcg/g
Fecal secretory IgA	847						<=2,040 mcg/mL
Gut Microbiome Metabolites							
Metabolic							
Short-Chain Fatty Acids (SCFA) (Total*) (Acetate, n-Butyrate, Propionate)	45.2						>=23.3 micromol/g
n-Butyrate Concentration	5.3						>=3.6 micromol/g
n-Butyrate %	11.7 L						11.8-33.3 %
Acetate %	56.4						48.1-69.2 %
Propionate %	31.9 H						<=29.3 %
Beta-glucuronidase	<DL L						368-6,266 U/g

*Total value is equal to the sum of all measurable parts.

†These results are not represented by quintile values.


Tests were developed and their performance characteristics determined by Genova Diagnostics. Unless otherwise noted with ♦, the assays have not been cleared by the U.S. Food and Drug Administration.

Methodology: DNA by qPCR



Gastrointestinal Microbiome (PCR)

Commensal Bacteria (PCR)

Result
CFU/g stoolReference Range
CFU/g stool

Bacteroidetes Phylum

<i>Bacteroides uniformis</i>	4.1E8		<=9.5E8
<i>Phocaeicola vulgatus</i>	1.3E7		<=8.3E8
<i>Barnesiella spp.</i>	1.5E8		3.0E6-2.9E8
<i>Odoribacter spp.</i>	1.0E8 H		<=9.5E7
<i>Prevotella spp.</i>	1.0E9		6.6E7-3.8E9

Firmicutes Phylum

<i>Anaerotruncus colihominis/massiliensis</i>	<DL		<=2.0E7
<i>Butyrivibrio crossotus</i>	<DL		<=3.3E7
<i>Clostridium spp.</i>	3.4E5		<=1.5E7
<i>Coprococcus eutactus</i>	<DL		<=1.2E8
<i>Faecalibacterium prausnitzii</i>	<DL L		1.1E6-1.1E9
<i>Lactobacillus spp.</i>	5.6E3		<=1.6E6
<i>Pseudoflavonifractor spp.</i>	<DL L		1.3E4-2.9E7
<i>Roseburia spp.</i>	<DL L		3.6E5-4.6E8
<i>Ruminococcus bromii</i>	<DL		<=1.5E9
<i>Veillonella spp.</i>	2.7E6		<=4.1E6

Actinobacteria Phylum

<i>Bifidobacterium spp.</i>	4.0E6		4.6E5-2.6E8
<i>Bifidobacterium longum subsp. longum</i>	6.3E4		<=1.3E8
<i>Collinsella aerofaciens</i>	9.9E4		<=1.3E8

Proteobacteria Phylum

<i>Desulfovibrio piger</i>	<DL		<=5.4E7
<i>Escherichia coli</i>	1.9E7 H		<=7.5E6
<i>Oxalobacter formigenes</i>	<DL		<=1.1E7

Euryarchaeota Phylum

<i>Methanobrevibacter smithii</i>	<DL		<=2.0E7
-----------------------------------	-----	--	---------

Fusobacteria Phylum

<i>Fusobacterium spp.</i>	<DL		<=1.8E5
---------------------------	-----	--	---------

Verrucomicrobia Phylum

<i>Akkermansia muciniphila</i>	<DL L		>=8.5E3
--------------------------------	-------	--	---------

The gray-shaded portion of a quintile reporting bar represents the proportion of the reference population with results below detection limit.

Commensal results and reference range values are displayed in a computer version of scientific notation, where the capital letter "E" indicates the exponent value (e.g., 7.3E6 equates to 7.3×10^6 or 7,300,000).

Gastrointestinal Microbiome (Culture)

Human microflora is influenced by environmental factors and the competitive ecosystem of the organisms in the GI tract. Pathogenic significance should be based upon clinical symptoms.

Microbiology Legend			
NG	NP	PP	P
			
No Growth	Non-Pathogen	Potential Pathogen	Pathogen

Bacteriology (Culture)

Lactobacillus spp.

NG

Escherichia coli

4+ NP

Bifidobacterium (Anaerobic Culture)


4+ NP

Additional Bacteria

Non-Pathogen: Organisms that fall under this category are those that constitute normal, commensal flora, or have not been recognized as etiological agents of disease.

Potential Pathogen: Organisms that fall under this category are considered potential or opportunistic pathogens when present in heavy growth.

Pathogen: The organisms that fall under this category have a well-recognized mechanism of pathogenicity in clinical literature and are considered significant regardless of the quantity that appears in the culture.



Additional Bacteria

Salmonella spp.

NG

Shigella spp.

NG

Vibrio species

3+ P

Proteus vulgaris


4+ PP

Pseudomonas species

3+ NP


gamma haemolytic Streptococcus

4+ NP



Mycology (Culture)

NG





Parasitology

Microscopic O&P Results

Microscopic O&P is capable of detecting all described gastrointestinal parasites. The organisms listed in the box represent those commonly found in microscopic stool analysis. Should an organism be detected that is not included in the list below, it will be reported in the Additional Results section. These results were obtained using wet preparation(s) and trichrome stained smear. For an extensive reference of all potentially detectable organisms, please visit www.gdx.net/product/gi-effects-comprehensive-stool-test

Genus/species	Result
Nematodes - roundworms	
<i>Ancylostoma/Necator</i> (Hookworm)	Not Detected
<i>Ascaris lumbricoides</i>	Not Detected
<i>Capillaria philippinensis</i>	Not Detected
<i>Enterobius vermicularis</i>	Not Detected
<i>Strongyloides stercoralis</i>	Not Detected
<i>Trichuris trichiura</i>	Not Detected
Cestodes - tapeworms	
<i>Diphyllobothrium latum</i>	Not Detected
<i>Dipylidium caninum</i>	Not Detected
<i>Hymenolepis diminuta</i>	Not Detected
<i>Hymenolepis nana</i>	Not Detected
<i>Taenia</i> spp.	Not Detected
Trematodes - flukes	
<i>Clonorchis/Opisthorchis</i> spp.	Not Detected
<i>Fasciola</i> spp./ <i>Fasciolopsis buski</i>	Not Detected
<i>Heterophyes/Metagonimus</i>	Not Detected
<i>Paragonimus</i> spp.	Not Detected
<i>Schistosoma</i> spp.	Not Detected
Protozoa	
<i>Balantidium coli</i>	Not Detected
<i>Blastocystis</i> spp.	Not Detected
<i>Chilomastix mesnili</i>	Not Detected
<i>Cryptosporidium</i> spp.	Not Detected
<i>Cyclospora cayetanensis</i>	Not Detected
<i>Dientamoeba fragilis</i>	Not Detected
<i>Entamoeba coli</i>	Not Detected
<i>Entamoeba histolytica/dispar</i>	Not Detected
<i>Entamoeba hartmanii</i>	Not Detected
<i>Entamoeba polecki</i>	Not Detected
<i>Endolimax nana</i>	Not Detected
<i>Giardia</i>	Not Detected
<i>Iodamoeba buetschlii</i>	Not Detected
<i>Cystoisospora</i> spp.	Not Detected
<i>Trichomonads</i> (e.g. <i>Pentatrichomonas</i>)	Not Detected
Additional Findings	
White Blood Cells	Not Detected
Charcot-Leyden Crystals	Not Detected
Other Infectious Findings	

One negative specimen does not rule out the possibility of a parasitic infection.

Parasitology

PCR Parasitology - Protozoa

Methodologies: DNA by PCR

Organism	Quantity	Units	Result	Expected Result
<i>Blastocystis</i> spp.	<2.14e2	femtograms/microliter C&S stool	Not Detected	Not Detected
<i>Cryptosporidium parvum/hominis</i>	<1.76e2	genome copies/microliter C&S stool	Not Detected	Not Detected
<i>Cyclospora cayetanensis</i>	<2.65e2	genome copies/microliter C&S stool	Not Detected	Not Detected
<i>Dientamoeba fragilis</i>	<1.84e2	genome copies/microliter C&S stool	Not Detected	Not Detected
<i>Entamoeba histolytica</i>	<9.64e1	genome copies/microliter C&S stool	Not Detected	Not Detected
<i>Giardia</i>	<1.36e1	genome copies/microliter C&S stool	Not Detected	Not Detected

Additional Results

Methodology: Fecal Immunochemical Testing (FIT)		
	Result	Expected Value
Fecal Occult Blood♦	Negative	Negative
Color††	Brown	
Consistency††	Formed/Normal	

††Results provided from patient input.
Tests were developed and their performance characteristics determined by Genova Diagnostics. Unless otherwise noted with ♦, the assays have not been cleared by the U.S. Food and Drug Administration.



Commentary

Lab Comments

Ceftriaxone and Cefazolin are not reportable for Vibrio pharmaceutical sensitivity testing.

Acute Value of: Vibrio species reported to office via client services.

06/16/2026 DH

A modified version of the assay cleared by the US Food and Drug Administration is used to measure PE1. Performance characteristics were confirmed or determined by Genova Diagnostics, Inc. in a manner consistent with CLIA requirements.

Commentary is provided to the practitioner for educational purposes and should not be interpreted as diagnostic or as treatment recommendations. Diagnosis and treatment decisions are the practitioner's responsibility.

For more information regarding GI Effects clinical interpretation, please refer to the GI Effects Support Guide at www.gdx.net/gieffectsguide

Methodology: Vitek 2® System Microbial Antibiotic susceptibility, Manual Minimum Inhibition Concentration

Bacteria Sensitivity

Prescriptive Agents

<i>Proteus vulgaris</i>	R	I	S-DD	S	NI
Ampicillin	R				
Amox./Clavulanic Acid				S	
Cefazolin	R				
Ciprofloxacin				S	
Tetracycline	R				
Trimethoprim/Sulfa				S	

Natural Agents

<i>Proteus vulgaris</i>	LOW INHIBITION	HIGH INHIBITION
Berberine		
Oregano		
Uva-Ursi		

Prescriptive Agents:

The R (Resistant) category implies isolate is not inhibited by obtainable levels of pharmaceutical agent.

The I (Intermediate) category includes isolates for which the minimum inhibition concentration (MIC) values usually approach obtainable pharmaceutical agent levels and for which response rates may be lower than for susceptible isolates.

The S-DD (Susceptible-Dose Dependent) category implies clinical efficacy when higher than normal dosage of a drug can be used and maximal concentration achieved.

The S (Susceptible) column implies that isolates are inhibited by the usually achievable concentrations of the pharmaceutical agent.

NI (No Interpretive guidelines established) category is used for organisms that currently do not have established guidelines for MIC interpretation.

Refer to published pharmaceutical guidelines for appropriate dosage therapy.

Natural Agents:

In this assay, inhibition is defined as the reduction level on organism growth as a direct result of inhibition by a substance. The level of inhibition is an indicator of how effective the substance was at limiting the growth of an organism in an in vitro environment. High inhibition indicates a greater ability by the substance to limit growth, while Low Inhibition a lesser ability to limit growth. The designated natural products should be considered investigational in nature and not be viewed as standard clinical treatment substances.

Bacteria Sensitivity

Prescriptive Agents

<i>Vibrio species</i>	R	I	S-DD	S	NI
Trimethoprim/Sulfa				S	

Natural Agents

<i>Vibrio species</i>	LOW INHIBITION	HIGH INHIBITION
Berberine		
Oregano		
Uva-Ursi		

Prescriptive Agents:

The R (Resistant) category implies isolate is not inhibited by obtainable levels of pharmaceutical agent.

The I (Intermediate) category includes isolates for which the minimum inhibition concentration (MIC) values usually approach obtainable pharmaceutical agent levels and for which response rates may be lower than for susceptible isolates.

The S-DD (Susceptible-Dose Dependent) category implies clinical efficacy when higher than normal dosage of a drug can be used and maximal concentration achieved.

The S (Susceptible) column implies that isolates are inhibited by the usually achievable concentrations of the pharmaceutical agent.

NI (No Interpretive guidelines established) category is used for organisms that currently do not have established guidelines for MIC interpretation.

Refer to published pharmaceutical guidelines for appropriate dosage therapy.

Natural Agents:

In this assay, inhibition is defined as the reduction level on organism growth as a direct result of inhibition by a substance. The level of inhibition is an indicator of how effective the substance was at limiting the growth of an organism in an in vitro environment. High inhibition indicates a greater ability by the substance to limit growth, while Low Inhibition a lesser ability to limit growth. The designated natural products should be considered investigational in nature and not be viewed as standard clinical treatment substances.