

# **Predicting virus-receptor mutant binding by molecular dynam**

Austin G. Meyer<sup>1,2,3\*</sup>, Sara L. Sawyer<sup>2</sup>, Andrew D. Ellington<sup>2</sup>,  
and Claus O. Wilke<sup>1</sup>

Address:

<sup>1</sup>Department of Integrative Biology, Institute for Cellular and Molecular Biology, and Center for Computational Biology and Bioinformatics. The University of Texas at Austin, Austin, TX 78712, USA.

<sup>2</sup>Department of Molecular Biosciences, Institute for Cellular and Molecular Biology, The University of Texas at Austin, Austin, TX 78712, USA.

<sup>3</sup>School of Medicine, Texas Tech University Health Sciences Center, Lubbock, TX 79430, USA.

\*Corresponding author

Email: [austin.meyer@utexas.edu](mailto:austin.meyer@utexas.edu)

Phone: +1 512 971 0123

Manuscript type: research article

Keywords: protein–protein interaction, molecular dynamics, arenavirus, machupo

# 1 Introduction

The computational prediction of mutational effects on protein–protein interactions remains a challenging problem. Several methods are available to perform an energy difference calculation from an experimentally determined co-crystal structure. For example, ~~computational alanine-scanning~~ end point methods can be performed rapidly, with relatively low computational cost (??). However, such methods ~~calculate the relative contribution to overall energy for each amino acid in a protein. They do not probe the interface directly, but rather calculate the total energy with each substitution~~ can suffer from various simplifying assumptions. For example, they generally use an implicit solvent approximation and assume the end state difference with minimal structural rearrangement is sufficient to discriminate energetic differences (??). Alternative approaches have been developed using machine learning, training coefficients in a weighted equation containing geometric and energetic parameters (????). Unfortunately, such machine-learning approaches often suffer in novel applications, for which available training sets are small or non-existent. As such, these methods are poorly suited for most host-virus protein–protein systems. By contrast, first principles methods ~~offer the chance to forgo training. However, the~~ can forgo training, but currently available methods such as free energy perturbation (FEP) and thermodynamic integration (TI) rely on a ~~two-state transitional~~ model (where one state may be wild-type and the other may be a mutant) ~~that transitions directly from one three-dimensional model to another with no intermediate steps to make rigorous free energy calculations~~ (????). ~~As a result, the accuracy of the measurement suffers~~ While these may be considered two of the gold standard techniques for calculating affinity differences, there are a huge number theoretical and technical complexities that must all be properly managed to ensure a converged solution (?). Such considerations quickly come to dominate the protocol, and the necessary book keeping introduces the possibility of human error (?). Moreover, as the two ~~proteins become increasingly dissimilar ; therefore~~ ending states look ever more dissimilar the chances of convergence falls rapidly. To ensure convergence, these techniques are ~~largely limited to~~ typically limited to small differences (such as point mutant comparisons) with a few, very impressive exceptions (???). For most, larger differences quickly become intractable as the number of intermediate steps required to compute a converged solution grows or the complexity of adding restraining potentials and computing approximations expands (???).

Here we propose ~~a new method~~ applying an existing method to a new test case to probe mutational effects in protein–protein interactions ~~directly~~ based on first principles simulation, ~~by pulling two proteins apart that are initially bound in a complex~~. We impart a pulling force within an all-atom molecular dynamics simulation on one member of the complex while the other is held in place. Then, we measure the force required for dissociation (????). As a proxy for relative binding affinity, we use the maximum applied force required for separation and the area under the force-versus-distance curve (AUC). Although such biasing techniques are common in protein-ligand binding problems, they are less commonly applied to protein–protein interactions, and almost never to mutational analysis in a protein–protein system. This is largely the result of free energy convergence difficulties and computational limitations (??); using maximum force or simple AUC can solve these problems. We also used the traditional dual topology FEP technique to compute relative free energy differences for comparison to our simplified force biased approach.

~~We applied our method~~ We used SMD and FEP to interrogate the interaction between machupo virus (MACV) spike glycoprotein (GP1) and the human transferrin receptor (hTfR1) (??). Machupo virus is an ambisense RNA virus of the arenavirus family (?). Worldwide, arenaviruses represent a significant source of emerging zoonotic diseases for the human population (?). Members of the arenavirus family include the Lassa fever virus endemic to West Africa, the lymphochoriomeningitis virus (LCMV) endemic to rodents in several areas of the United States, and the Guanarito, Junin, and Machupo viruses endemic to rodents in South America (?). The South American arenaviruses typically infect humans after rodent contamination and can cause a devastating hemorrhagic fever with high mortality (?).

The hTfR1 is the primary receptor used by MACV for binding its host cell prior to infection. The primary role of hTfR1 *in vivo* is to bind transferrin for cellular iron uptake. The hTfR1 protein contains three extracellular domains: two basilar domains and an apical domain. The two basilar domains serve most of the transferrin-binding function (??). Viral entry is initiated by GP1 binding to the apical domain of hTfR1. Previous work has indicated that the GP1/hTfR1 binding interaction is the primary determinant of MACV host range variation (??). The co-crystal structure shows the high affinity interaction between GP1 and hTfR1 forces the normally flexible loop in the apical domain of hTfR1 into a rigid ~~beta-pleated~~  $\beta$ -pleated sheet domain. For GP1,

several extended loops mediate binding to hTfR1 (??), and many of the interface interactions are mediated by extensive hydrogen-bonding networks (?). Experimental alanine-scanning and whole-cell infectivity assays have identified several sites in both GP1 and hTfR1 that are probably critical for establishing infection (??).

We applied our computational method to wild type (WT) and mutant complexes, and found that we could resolve relative differences in unbinding and predict significant affinity changes. Importantly, the affinity changes predicted using only max force or AUC show a strong correlation with rigorous relative free energy differences computed by FEP. At sites known to be important for successful viral entry, we found that the biochemical cause of reduced infectivity may not be as simple as the static structure suggests. For example, the static structure shows a hydrogen-bonding network connected to site Asn348 in hTfR1. According to our simulations, this network may not affect binding affinity directly. In addition, our study offers an all-atom steered molecular dynamic approach to avoid some of the pitfalls of several existing methods used to evaluate mutations in protein-protein interfaces.

## 2 Materials and Methods

### 2.1 System Modeling

For our experiments, we used the experimentally determined GP1/hTfR1 structure (PDBID: 3KAS) (?). The apical domain of hTfR1 interacts directly with GP1 while the other two domains are closer to the cell membrane and have essentially no interaction with GP1. The biophysical independence of the apical domain allowed us to isolate it without significantly affecting the GP1/hTfR1 interaction.

We used the protein visualization software PyMOL (?) to remove residues 121-190, 301-329, and 383-756 in the hTfR1. No residues were removed from the viral protein. ~~We also~~ Figure 1 shows a model of the initial structure and that of the pared structure. Although GP1 has several glycosylatable residues, we opted to use the de-glycosylated protein for this study. The complexity of correctly parameterizing diverse sugar moieties is outside of the scope of this paper. Furthermore, although it is known that GP1 is glycosylated, and some of those sugars contact

hTfR1, the sugars in the available PDB structure are not physiological for mammals (?). In total we removed 10 ~~sugar residues from the native structure~~ sugars from the crystal structure for this study.

After system reduction, the Visual Molecular Dynamics (VMD) (?) package along with its system of back-ends was used for all subsequent modeling. The Orient add-on package allowed us to rotate the system axis such that the direction of steering was oriented directly down the z-axis. De-glycosylation simplified the system such that Autopsf could easily find the chain terminations and patch them appropriately. The Solvate package was used to generate a TIP3P water model with a 5 Ångstrom buffer (relative to the maximum dimensions of the proteins) on all sides except down the positive z-axis where a 20 Ångstrom buffer was created. Finally, we used the Autoionize package to place 150 millimolar NaCl and neutralize the total system charge. In the end, each modeled system had approximately 28,000 atoms.

## 2.2 Equilibration

NAMD was used for all simulations in this study (?). In addition to the modeled system, for equilibration we generated a configuration file that fixed the  $\alpha$ -carbon backbone. This was accomplished by setting  ~~$\beta=1$~~  the B-factor column to 1 for the fixed atoms and to zero for all other atoms. Further, we generated a configuration file with fixed  $\alpha$ -carbon atoms at residues 41-92 (numbered linearly, in this case, starting at 1 for the first amino acid as was required for ~~VMD~~ NAMD) in the hTfR1. The second file was used to affix a harmonic restraint, thus preventing any unfolding due to system reduction. More importantly, the harmonic restraint allowed the protein complex to equilibrate while preventing any drift from its predefined ~~orientation~~ position; the restraint did not constrain the structure of each protein, or the relative position or orientation of the two proteins to each other. Finally, we calculated the system center and dimensions for use in molecular dynamics settings.

The exact NAMD configuration file is made available as a supplement and a github repository with all necessary set up files is provided. We used the Charmm27 (?) all-atom force field. The initial system temperature was set to 310K. Several typical MD settings were used including switching and cutoff distances (see provided configuration file). In addition, we used a 2 femtosecond time step with rigid bonds ~~(the exact configuration is provided in the supplement)~~. We used periodic boundary conditions with the particle mesh ewald (PME) method of computing full system elec-

116 trostatics outside of the explicit box. Furthermore, we used a group pressure cell, flexible box,  
117 ~~and langevin-piston~~ langevin barostat and langevin thermostat during equilibration. ~~Finally, a~~  
118 harmonic restraint (called harmonic constraint in VMD) was set as stated previously.

119 To start the simulation, the ~~langevin-piston~~ barostat was switched off and the system was mini-  
120 mized for 1000 steps. Next, the fixed backbone was released, and the system was minimized for an  
121 additional 1000 time steps. Subsequently, the system was released into all-atom molecular dynam-  
122 ics for 3000 steps. Finally, the langevin piston was turned on and the system was simulated for 2 ns  
123 (1,000,000 steps) of chemical time. For each mutant, twenty independent equilibration replicates  
124 were run with an identical protocol.

## 125 2.3 Steered Molecular Dynamics

126 We used the final state from each equilibrated system to restart another MD simulation. Our steering  
127 protocol is fundamentally similar to ? with slightly different parameter choices. Perhaps the one  
128 significant difference lies in our choosing to not use a thermostat or barostat. However, this should  
129 not be a problem since we are not trying to calculate the binding free energy by any physically  
130 rigorous approach (the Jarzynski inequality being one example). Following equilibration, the final  
131 state of each simulation was used to generate a configuration file fixing the  $\alpha$ -carbon on residues  
132 1, 58, 73-83, 96, 136, 137, 138, and 161 (again with linear numbering) in the hTfR1. ~~Also, the~~  
133 These residues were selected as they are far from the binding interface and sufficiently distributed  
134 to prevent any orientational motion of the receptor relative to the viral spike protein. The center of  
135 mass of the  ~~$\alpha$ -carbon-carbons~~ of all residues (163-318 in linear numbering) in ~~the~~ GP1 ~~were given~~  
136 ~~an occupancy of 1, indicating they had an~~ received an applied force during the simulation.

137 ~~In every way possible the subsequent steering simulation had the same parameters as the~~  
138 ~~equilibration simulation. Thus, the steered molecular dynamics (SMD) simulation was a typical~~  
139 ~~restart with slightly different settings~~ The NAMD convention does not actually apply a force to all  
140  $\alpha$ -carbon atoms, but rather uses the selection to compute an initial center of mass. Then, during the  
141 steering run, the single center of mass point is pulled with the parameters described below. We used  
142 the same force field parameters (exclude, cutoff, switching, etc.), the same integrator parameters  
143 (time step, rigidbonds on, all molecular being wrapped, etc.), and the same particle mesh ewald  
144 parameters as in equilibration. Periodic boundary conditions were incorporated as part of the sys-

tem (as is the convention in NAMD restart) and PME was again used to approximate full system electrostatics. ~~In addition, we first tuned the pulling velocity for practical time considerations. Subsequently, the spring constant was adjusted to ensure a sufficient~~

We ran test simulations at several force constants, and visually inspected the results. A force constant of  $5 \text{ kcal/mol/\AA}^2$  was chosen due to its relatively low signal-to-noise ratio ~~with the given velocity. As a result, the spring constant was set to 5.~~ This is slightly lower than the more common  $7 \text{ kcal/mol/\AA}^2$ , ~~velocity set to~~ found in several recent studies; however, it seems that value is so commonly selected primarily because it is the force constant found in the NAMD, SMD tutorial available through the developers. Moreover, the force constant could very likely be set to a range of nearby values with little loss in predictive power. In SMD experiments the pulling velocity should be as low as possible for the available computational time ~~??~~. Thus we choose a velocity of  $0.000001 \text{ \AA/fs} = 1 \text{ \AA/ns}$ , and direction down the positive  $z$ -axis. This number could be set to pull faster if the computing time must be reduced, but slower than necessary pulling speeds are not typically considered problematic.

~~Finally,~~ SMD was run for 15 ns (7,500,000 time steps) of chemical time. For each simulation, we randomly selected one of the equilibration runs for restart. We ran 50 replicate simulations per mutant for a total of 550 SMD simulations. All GP1/hTfR1 complexes separated by greater than 4  $\text{\AA}$  and many separated to 10 or more.

To leave the final trajectory of a tractable size, only 1000 evenly spaced frames were retained from each simulation, leaving a final trajectory size of 323 MB. See the supplemental movie for a representative unbinding trajectory. Initial development of the SMD protocol was carried out on the Lonestar cluster at the Texas Advanced Computing Center (TACC). All production SMD simulations were performed on the Hrothgar cluster at Texas Tech University, using NAMD 2.9. Each simulation was parallelized over 60 computational cores and utilized approximately 20 hours of computing time. The total chemical time simulated for this project was nearly  $10 \mu\text{s}$ , requiring slightly over 1 million cpu-hours.

## 2.4 Free Energy Perturbation

Briefly, we used the traditional dual topology approach to FEP. This involves a thermodynamic cycle where a set of atoms are progressively decoupled from the environment while another set of

atoms are progressively coupled. To compute the relative free energy difference requires knowing the free energy change when the transformation is carried out for the bound complex and the individual protein. Then, one can compute the relative free energy difference between a WT and mutant complex by taking the difference between the energy required to decouple/couple the atoms in solution from the energy required to decouple/couple the atoms in the bound complex.

Again, the NAMD configuration file is made available as a supplemental file. We used a similar configuration to that in equilibration. One significant difference was to make a cubic water box with a side length equal to the long axis of the complex plus a 10 Å buffer on either side, and simply restrict center of mass motion with the NAMD setting. This was done to avoid affecting the system energy while calculating free energy differences.

The transition protocol for bound and free protein systems were identical. They started with 1000 steps of minimization and 250,000 steps of equilibration in the starting state for the forward and reverse directions. Phase transitions were carried out in steps of  $\lambda=0.05$ . Each transition was carried out for 250,000 steps. The first 100,000 steps after phase transition were reserved for equilibration and the final 150,000 steps were used for data collection.

The FEP analysis tool in VMD was used for subsequent analysis. We used it to perform error analysis and compute the Bennett acceptance ratio as the maximum likelihood free energy difference of the two states under consideration. Though the larger transitions presented difficulty in a small number of windows, forward and reverse hysteresis was generally in good agreement for all complexes. The double mutants were performed by first doing the Y211A mutation followed by the other of the two mutants. Then, the numbers were simply added together to get the total energetic difference.

## 2.5 Post-processing

The python packages MDAnalysis (?) and ProDy (?) were both used at various points in post-processing. The molecular trajectory (comprising the atomic coordinates per time) was parsed to compute the center-of-mass for each of the two complexes. The starting center-of-mass distance was set to zero and the distance was re-computed at each time step relative to the starting distance.

~~The force was output directly to the standard log file.~~

The statistical package R was used for all further analysis and visualization. Each of the 50



independent trajectories per mutant produced a fairly noisy force curve. ~~To reduce the noise, we averaged each point with~~ The force curves for each mutant were smoothed either by averaging over many replicates as in Figure 4 or by selecting a 50 ~~point interpolation window frame interpolation window for averaging.~~ The latter approach was used for statistical analysis. The ~~window ensured that the averaged maximum applied force was never cut off near the beginning of the simulation due to lost time steps (caused by averaging).~~ The two primary descriptive statistics we used were maximum interpolated applied force and total area under the interpolated curve (AUC). We used the pairwise.t.test function with the FDR p-value correction to test for significance. The ggplot (?) package was used to generate most of the figures.

## 3 Results

### 3.1 The GP1/hTfR1 system

The GP1/hTfR1 interface marks a particularly important and useful test system. There are several sites on both the human and viral protein known to affect the infectivity phenotype of MACV. ~~Virtually all~~ Many of the important sites have been mapped by *in vitro* flow-cytometry based entry assays. The GP1/hTfR1 interface appears not to be dominated by one particular type of interaction (electrostatics, hydrogen-bonding, or van der Waals). In addition, much of the binding domain on hTfR1 is on a ~~flexible loop and undergoes a large structural rearrangement upon binding, making~~ loop that is flexible prior to viral binding, but organizes to become a strand of a  $\beta$ -sheet on binding. ~~As a result,~~ many other computational techniques (??) only marginally useful. The complex nature of this interface represents a particularly difficult challenge for traditional computational analysis.

~~For our experiments, we used the experimentally determined GP1/hTfR1 structure (PDBID: 3KAS). The apical domain of hTfR1 interacts directly with GP1 while the other two domains are closer to the cell membrane and have essentially no interaction with GP1. The biophysical independence of the apical domain allowed us to isolate it without significantly affecting the GP1/hTfR1 interaction. Figure 1 shows a model of the initial structure and that of the pared structure. Although GP1 has several glycosylatable residues, we opted to use the de-glycosylated protein for this study. The complexity of correctly parameterizing diverse sugar moieties is outside~~

~~of the scope of this paper. Furthermore, although it is known that GP1 is glycosylated, and some of those sugars contact hTfR1, the sugars in the available PDB structure are not physiological for mammals.~~

In total, we tested 7 point mutants and 3 double mutants in addition to the WT complex (Table 1 and Figure 7). All of the mutations are within 5 Å of the protein-protein interface. Mutations at site hTfR1 site 211 have proven capable of causing loss-of-entry according to *in vitro* flow-cytometry infection assays or known host-range limitations (???). Most likely, this effect is caused by the destruction of a critical hydrogen bond to Ser113 or Ser111 in GP1. The lost hydrogen bond would lead to the subsequent loss of a large hydrogen-bonding network seen in the crystal structure (Table 1) (?). In a manner similar to site 211, Asn348 appears to be important for binding by participating in a critical hydrogen bonding network (??) to GP1. In particular, Asn348Lys is reported in the literature to cause significantly reduced viral entry *in vivo* (Table 1) (??). Finally, an alanine mutation at site 111 in GP1 (mutation vArg111Ala) has also been shown to cause decreased entry (Table 1) (?). For notation purposes, the viral site is always referred to with a preceding ‘v’.

### 3.2 Steered molecular dynamics simulations

Despite, the fact that viral binding occurs at the site of a flexible loop in the free hTfR structure, our data shows after binding the strand is extremely rigid. None of the sites in the interfacial strand have root mean squared fluctuation (RMSF) values in the top half of all sites in the complex during equilibration (Figure 2). This is unsurprising considering the high degree of burial that occurs as a result of viral binding. In addition, computing the root mean squared deviation (RMSD) of the entire structure over the trajectory shows that none of the mutations are so deleterious as the cause rapid unbinding. In fact, the RMSD over trajectory looks incredibly invariant across mutants (Figure 3).

~~We proceeded to analyze this system using molecular dynamics. To ensure a natural solution structure, we performed an initial equilibration phase to reduce the intrinsic crystallization distortions. First, we generated a physically realistic simulation world around the protein complex. The environment included an explicitly modeled bath of water, and a physiologically relevant concentration of ions. Next, we carried out successive rounds of energy minimization with a fixed  $\alpha$ -carbon backbone, followed by heating, and backbone release. Last, we performed a short round of all-atom~~

259 molecular dynamics with minimal restraints. We used the final state from each equilibrated system  
260 to restart another MD simulation. For SMD, at the centroid of several atoms a force was applied to  
261 a single point in GP1. By

## 262 3.2 Molecular dynamics simulations

263 We proceeded to analyze this system using two molecular dynamics techniques. First, in SMD by  
264 using a known spring force constant, and pulling with a constant velocity, we could calculate the  
265 applied force during protein-protein dissociation (??). A typical averaged force curve comparison  
266 can be seen in Figure 4. The force curves for each mutant were smoothed either by averaging  
267 over many replicates as in Figure 4 or by selecting an interpolation window for averaging. The  
268 latter approach was used for statistical analysis. In addition, to minimize the computational time in  
269 unproductive molecular dynamics, we shortened the simulation time such that the interaction just  
270 barely dropped to zero near the end of each run. As seen in Figure 4, this distance was relatively  
271 consistent among mutants. The supplementary movie visually illustrates the separation distance  
272 between peptide domains. The quantities maximum applied force and AUC were derived from the  
273 force versus distances curves. Their summary statistics are reported in Table ???. As we are more  
274 interested in the phenotypic impact of interface mutations we avoided many of the more physically  
275 rigorous, but technically complicated calculations that are possible with SMD (??).

276 Before systematically applying SMD to the GP1/hTfR1 interaction, we needed to ensure the  
277 method was sufficiently sensitive to distinguish between relatively minor point mutations. While  
278 SMD has been applied previously to measure the binding energy of high-affinity T-cell receptor  
279 interactions (??), to our knowledge the method has never been used previously to parse small  
280 differences in a complex energy landscape. For this initial sensitivity analysis, we tested alanine  
281 substitutions congruent with the traditional experimental and computational approach.

282 We proceeded to compare our SMD results to that of the standard dual topology FEP approach  
283 to calculate relative free energy differences. The correlation between the energetically rigorous  
284 FEP and our statistical approach is high. For all 11 complexes tested, the correlation between max  
285 force and FEP was  $r=-0.767$  at  $p=0.005857$  (Figure 6), and the correlation between AUC and FEP  
286 was  $r=-0.729$  at  $p=0.011$ . As a result of the strong correlation, we will generally refer to the SMD  
287 differences only.

We found that relative to WT, one alanine mutation (Tyr211Ala) produced a very large and statistically significant difference in the maximum applied force and AUC (Figure 4, Table 2), while the other two did not (Table 2). When considering additional mutants (also discussed below), we found that maximum applied force was generally sufficient to distinguish mutants (Tables 2 and 3), and AUC was able to add a few more statistically significant differences (Table 4). In general, however, maximum applied force seemed to be the more sensitive statistic than AUC.

### 3.3 Comparative analysis of the GP1/hTfR1 interface

Considering the involvement of extended hydrogen-bonding networks in the GP1/hTfR1 interface, it was not clear that individual alanine mutations, even those that should destroy such networks, would significantly change the strength of interaction. One major advantage of first principles simulations is the ability to test mutations other than alanine without additional underlying assumptions in the energy function. As shown in Table 1, we made additional mutations based on biochemical intuition or available experimental data to chemically diverse amino acids including tryptophan, lysine, aspartate, and threonine. Several mutations caused significant relative affinity changes. In addition, to detect synergistic effects, we tested several double mutants where both mutations appeared to cause similar changes in binding. Then, we compared the size of those differences to single mutants (Figure 5 and 7).

Although Tyr211Ala appears to have a large impact on binding affinity, no single mutant can provide enough evidence to understand the biochemical difference in binding mechanism. Since alanine is both smaller than tyrosine and also incapable of participating in hydrogen-bond interactions, we tested further mutations to identify the critical biochemical difference responsible for change in binding affinity. In particular, we substituted smaller side chains that, like tyrosine, were capable of hydrogen bonding. We chose Tyr211Asp and Tyr211Thr, two mutations that have been discussed in the context of selection pressure on hosts in rodent populations (???). Both mutations proved capable of causing a significant change in binding affinity in our simulations, but the change appeared to be increased affinity (Figures 5 and 7, and Table 3).

We also simulated several point mutations at Asn348 in the hTfR1. As discussed above, the alanine mutation at this site showed no significant difference in maximum applied force or AUC from WT (Tables 3 and 4). In addition, neither the Asn348Lys nor the Asn348Trp mutation showed a sig-

nificant difference from WT. For both of these mutations, however, mean maximum applied force and mean AUC was lower than for WT (See Table ??). On the other hand, there was a detectable difference between Asn348Ala and Asn348Lys (Tables 3 and 4), with Asn348Lys being a weaker binder. Moreover, Asn348Trp showed nearly identical results to Asn348Lys. The mutations to large amino acids (Asn348Trp and Asn348Lys) produced nearly identical affinity changes, whereas the mutations to amino acids not capable of hydrogen bonding (Asn348Ala and Asn348Trp) produced significantly different affinity changes (Table 2). To check the consistency of our results, we hypothesized that the combination of Tyr211Ala and Asn348Trp, being chemically disconnected in two different hydrogen-bonding networks, would lead to a synergistic loss-of-binding. As expected, the double mutant was the weakest binding mutant tested ( $p < 10^{-6}$ , Tables 3 and 4) in this study. Further, according to maximum applied force (but not AUC), the combination of Tyr211Ala and Asn348Trp also showed significantly weaker binding than Tyr211Ala by itself (Tables 3 and 4). We suspect that the effect of Asn348Trp alone is near the limit of detection using our method. A larger number of replicates would possibly have resolved affinity differences between Asn348Trp and WT or other mutants more consistently.

Last, we further analyzed a single mutation in GP1, vArg111Ala. As mentioned in the previous subsection, in our simulations this mutant showed no significant change in either maximum applied force or AUC (Tables 3 and 4), even though both quantities were, on average, lower than in WT (Table ??). This result was somewhat surprising, since Tyr211Ala, presumably disrupting the same hydrogen-bonding network as vArg111Ala, displayed a significant reduction in affinity. To probe the interaction between position 111 in the GP1 and position 211 in the hTfR1 further, we also tested the double mutant vArg111Ala/Tyr211Ala. This double mutant showed affinity indistinguishable from WT and significantly higher than Tyr211Ala alone (Table 2). This result shows that the two sites do indeed interact, and that replacing the hydrogen-bonding network at these sites with a hydrophobic interaction could lead to comparable binding affinity.

## 4 Discussion

We have applied a method utilizing steering forces in all-atom molecular dynamics simulations to evaluate the effects of mutations at the GP1/hTfR1 interface. We modeled mutations at sev-

eral sites in the GP1/hTfR1 interface, and verified that our computational protocol was sensitive enough to distinguish point mutants in hTfR1. Further, we identified two test statistics, maximum applied force and AUC, that can be used as proxies for binding affinity. Both of these statistics correlate well with FEP, but offer the simplicity of not requiring a large commitment to planning for the theoretical issues inherent to free energy methods. We systematically tested several point mutations to understand their contribution to the binding interaction. In the case of Asn348Lys, we have shown that the static structure provides little insight into why this mutation causes loss-of-infectivity *in vivo*. While Asn348 appears to be involved in a hydrogen-bonding network in the static structure, change in binding at that site may actually be caused by size and charge restriction. We also found that a negatively polar residue at site 211 in hTfR1 seem critical for a tight binding interaction. Any non-polar mutation at Tyr211 in hTfR1 is likely to completely halt viral entry and dramatically decrease the chances of MACV infection.

~~By extending the SMD technique to investigate a protein-protein interaction we have, in many ways, changed the scope of its application. Traditionally this technique~~ Traditionally SMD has been either applied to compute equilibrium ~~quantities~~ free energies via a non-equilibrium approximation (??), used to estimate protein stability through unfolding (?), or used to calculate the absolute free energy of small molecule ligand binding (?). Likewise, others have used SMD to understand the process of binding and unbinding at a resolution unmatched by experiment (??). Here, we have shown that SMD can provide insight into the *relative* strength of protein-protein interactions. ~~In contrast to more traditional absolute affinity calculations, relative binding is much more useful for experimental application.~~ Via SMD, one can separate mutations whose likely effect is altered binding affinity ~~from mutations that may affect other factors such as protein expression, protein folding, or interactions with other proteins not considered in the simulation~~ with simple statistics like maximum force of separation. Thus, SMD may open avenues for subsequent experimental work in some situations where FEP may be prohibitively difficult.

Our findings ~~may~~ rationalize several effects observed in both infectivity data and rodent populations (??). First, we found that some substitutions at positions 211 and 348 did affect the strength of receptor binding. However, the computational data suggest that the reason and nature of the effects at these two sites are very different. At position 211, mutations to non-polar residues cause a large change in binding. This is congruent with what is known from viral entry data (??). By contrast,

375 mutations at position 348 need only be small to maintain WT binding. The ability to hydrogen bond  
376 appears to be insignificant. This can be inferred from the fact that Tyr211Ala paired with large (Trp)  
377 and positively charged (Lys) substitutions at position 348 results in a larger than expected synergis-  
378 tic difference. That is, the double mutant Tyr211Ala/Asn348Trp caused a much larger decrease in  
379 binding than we expected from either mutation individually. Third, the GP1 mutation vArg111Ala  
380 causes a loss-of-infection during *in vitro* infectivity assays (?), yet it was indistinguishable from  
381 the WT complex in our simulations. Although Tyr211Ala was the most disruptive single mutant  
382 we tested, vArg111Ala in the GP1 was able to restore mean maximum applied force to WT levels  
383 (Table ??), and to levels significantly higher than observed for Tyr211Ala alone.

384 We would like to emphasize here that we cannot expect perfect agreement between our simula-  
385 tions and the available experimental data, but the correspondence to a well established free energy  
386 method is bolsters our conclusions. While we have shown that our method can distinguish individ-  
387 ual point mutations, we do not know the limit of detection with our method. First, it is possible  
388 that some mutants display measurable phenotypic effects in experiments yet appear identical ~~under~~  
389 ~~our method~~ in simulation. More extensive sampling or refinement of the simulation protocol could  
390 help to differentiate such mutants (see also next paragraph). Second, ~~our~~ the SMD method is fun-  
391 damentally limited by the accuracy of our starting structure. Third, the available experimental data  
392 for the GP1/hTfR1 system were generally obtained from entry assays or whole-cell binding assays  
393 rather than molecular binding assays. A mutant may cause a phenotypic difference in infectivity  
394 without generating a signal by our method. For example, entry could be lost in the experimental  
395 system because the protein is grossly or partially misfolded. An additional analytical step with  
396 circular dichroism or an analogous technique could clarify such large-scale folding differences.  
397 Further, since our simulations start with a bound structure, any changes that may dramatically af-  
398 fect the rate of association (different folds, trafficking issues, etc.) or relative orientation of the two  
399 proteins would be underestimated by our method.

400 There are a few additional challenges for investigating host-virus interaction via molecular  
401 dynamics simulation. As with any atomistic simulation, there is going to be a fairly large noise  
402 to signal ratio. To reduce noise, one could further customize each simulation, e.g. by determining  
403 the optimal pulling speed. Furthermore, larger amounts of computational resources will have a  
404 direct and powerful impact on the strength of any atomistic study (?). Such resources could come

in the form of increased compute time, improved code, or customized hardware for floating point operations (?). With improved resources, we could investigate thousands of individual permutations in the GP1/hTfR1 binding interface. In addition, with additional compute time it would be possible to incorporate equilibrium sampling approaches (?) or use brute force equilibrium approaches (?) to improve resolution.

For future studies, although our approach offers the simplicity of not requiring prior knowledge about a system of interest (other than a bound model), ~~all-atom, unrestrained SMD is likely at this point SMD may~~ not the best approach for ~~most-many~~ relative affinity calculations. To ensure one's results are independent of the dissociation path one selects would require computing the work of separation for all likely paths. Such an approach eventually requires using the Jarzynski inequality (?) to establish a lower limit for binding energy and would quickly become computationally inefficient for evaluating a large number of mutations in most systems. ~~By contrast, However,~~ considering the strong correlation between FEP and SMD in this system, it may not be important to ensure one's results are path independent for relative affinity calculations.

More importantly, with no *a priori* knowledge of the appropriate number of equilibration samples, the during of equilibration, the appropriate number of pulling runs, or the best pulling speed means the computational expense in our SMD protocol may not be commensurate with the information provided. For example, another all atom approach that makes calculations via short simulations of spatially restrained complexes has proven capable of generating relatively accurate binding affinities with less compute time than is required from our steering strategy ~~—(??)~~. That being said, there is no reason to believe this SMD approach to mutagenic studies could not be optimized to reduce computational expense. Further analysis would be needed to understand the lower limits of resources required for accurate predictions.

## 5 Acknowledgements

This work was carried out using high-performance computing resources provided by the High Performance Computing Center (HPCC) at Texas Tech University at Lubbock (<http://www.hpcc.ttu.edu>) and the Texas Advanced Computing Center (TACC) at The University of Texas at Austin (<http://www.tacc.utexas.edu>). The authors acknowledge Bryan Sutton for opening



433 access to the Hrothgar cluster. The authors also acknowledge the reviewers Ilan Samish and Matteo  
434 Masetti for their helpful comments in preparing this manuscript.




Figure 1: The GP1/hTfR1 complex. GP1 is shown in blue and hTfR1 is shown in green. (A) The full, de-glycosylated GP1/hTfR1 co-crystal structure. (B) The reduced structure used in SMD simulations.




Figure 2: RMSF values. The RMSF values for every site in the bound complex computed during the equilibration phase of the protocol. Each color represents the average over 20 trajectories of a single mutant. The plot shows the flexibility of each site is essentially independent of mutation, and none of the sites above 0.72 Å are found in the interface.




Figure 3: RMSD values. The RMSD values over the time of the trajectory computed during the equilibration phase of the protocol. Each color represents the average over 20 trajectories of a single mutant. The plot shows none of the mutants causes immediate unbinding of the protein–protein complex. In addition, the universal upward trend near the end of the equilibration trajectories may indicate the crystal is more tightly packed than would normally occur in solution.




Figure 4: Force versus distance curve of WT and the Y211A mutant. The average force curve for 50 replicates of the WT complex is shown in black, and the average of 50 replicates of the Y211A mutant is shown in red. There is a large difference in both maximum applied force and AUC between the two complexes.




Figure 5: Distribution of interpolated maximum force for three different GP1/hTfR1 complexes. The WT GP1-hTfR1 complex in the middle is flanked by the tighter binding mutant Y211D on the right and the weaker binding double mutant N348W/Y211A on the left. The large non-overlapping areas indicate a large and statistically significant difference in these three complexes.




Figure 6: [Max force versus free energy perturbation](#) Scatter plot of maximum force in SMD versus the relative free energy difference calculated by FEP for all 10 mutants tested plus the WT complex. The WT complex for FEP was simply set to 0.0. The correlation between the two is  $r=-0.767$  with  $p=0.005857$ .




Figure 7: Distribution of interpolated maximum force for all bound complexes tested. Stars above the boxplots indicate a statistically significant difference in mean maximum force relative to the WT complex.



Table 1: Summary of prior information available for each mutation tested. Observed *in vivo* refers to mutations that have been observed in rodent populations. Phenotype *in vitro* refers to the observed phenotype in *in vitro* viral entry assays.

Mutation	Observed <i>in vivo</i>	Phenotype <i>in vitro</i>
WT	Yes	Normal Entry
N348A	No	-
N348K	Yes	Diminished Entry
N348W	No	-
Y211A	No	No Expression
Y211D	Yes	No Expression
Y211T	No	Diminished Entry
vR111A	No	Diminished Entry
N348A/Y211A	No	-
N348W/Y211A	No	-
vR111A/Y211A	No	-

Table 2: Pairwise differences (row variable minus column variable) in mean maximum applied force. Bolded values are statistically significant at  $p < 0.05$ . Values marked with \* and \*\* are significant at  $p < 0.01$  and  $p < 0.001$ , respectively.

	WT	N348A	N348W	N348K	Y211A	Y211D	Y211T	vR111A	N348A/Y211A	N348W/Y211A
N348A	27.33									
N348W	-39.87	<b>-67.20</b>								
N348K	-40.64	<b>-67.97</b>	-0.77							
Y211A	<b>-87.97*</b>	<b>-115.30**</b>	-48.10	-47.33						
Y211D	<b>103.44**</b>	<b>76.12*</b>	<b>143.31**</b>	<b>144.08**</b>	<b>191.42**</b>					
Y211T	<b>89.17*</b>	<b>61.85</b>	<b>129.04**</b>	<b>129.81**</b>	<b>177.15**</b>	-14.27				
vR111A	-17.76	-44.94	22.26	23.03	<b>70.36</b>	<b>-121.05**</b>	<b>-106.78**</b>			
N348A/Y211A	-9.04	-36.37	30.83	31.60	<b>78.93</b>	<b>-112.48**</b>	<b>-98.21*</b>	8.57		
N348W/Y211A	<b>-154.79**</b>	<b>-182.12**</b>	<b>-114.92**</b>	<b>-114.15**</b>	<b>-66.2</b>	<b>-258.23**</b>	<b>-243.96**</b>	<b>-137.18**</b>	<b>-145.75**</b>	
vR111A/Y211A	8.71	-18.12	48.58	49.35	<b>96.68*</b>	<b>-94.74*</b>	<b>-80.46*</b>	26.31	0.59	<b>163.50**</b>

Table 3: Pairwise difference  $p$ -values for maximum applied force. Bolded values are statistically significant at  $p < 0.05$ .

	WT	N348A	N348W	N348K	Y211A	Y211D	Y211T	vR111A	N348A/Y211A	N348W/Y211A
N348A	0.35									
N348W	0.22	<b>0.012</b>								
N348K	0.22	<b>0.012</b>	0.98							
Y211A	<b>0.0045</b>	<b>1.6x10<sup>-05</sup></b>	0.14	0.15						
Y211D	<b>0.00083</b>	<b>0.0045</b>	<b>3.9x10<sup>-06</sup></b>	<b>3.9x10<sup>-06</sup></b>	<b>5.7x10<sup>-10</sup></b>					
Y211T	<b>0.0042</b>	<b>0.022</b>	<b>3.0x10<sup>-05</sup></b>	<b>3.0x10<sup>-05</sup></b>	<b>1.0x10<sup>-08</sup></b>	0.67				
vR111A	0.59	0.11	0.51	0.51	<b>0.024</b>	<b>8.9x10<sup>-05</sup></b>	<b>0.00056</b>			
N348A/Y211A	0.78	0.20	0.35	0.35	<b>0.011</b>	<b>0.00026</b>	<b>0.0016</b>	0.78		
N348W/Y211A	<b>6.2x10<sup>-07</sup></b>	<b>9.3x10<sup>-12</sup></b>	<b>0.00021</b>	<b>0.00026</b>	<b>0.032</b>	<b>&lt; 2x10<sup>-16</sup></b>	<b>2.4x10<sup>-15</sup></b>	<b>9.1x10<sup>-06</sup></b>	<b>2.8x10<sup>-06</sup></b>	
vR111A/Y211A	0.78	0.52	0.14	0.14	<b>0.0019</b>	<b>0.0024</b>	<b>0.010</b>	0.44	0.59	<b>1.6x10<sup>-07</sup></b>

Table 4: Pairwise difference  $p$ -values for interpolated AUC. Bolded values are statistically significant at  $p < 0.05$ .

	WT	N348A	N348W	N348K	Y211A	Y211D	Y211T	vR111A	N348A/Y211A	N348W/Y211A
N348A	0.77									
N348W	0.67	0.42								
N348K	0.17	0.056	0.38							
Y211A	<b>0.00094</b>	<b>3.5x10<sup>-05</sup></b>	<b>0.0045</b>	0.063						
Y211D	<b>0.0072</b>	<b>0.0051</b>	<b>0.0016</b>	<b>4.6x10<sup>-05</sup></b>	<b>1.2x10<sup>-09</sup></b>					
Y211T	<b>0.016</b>	<b>0.014</b>	<b>0.0041</b>	<b>0.00016</b>	<b>6.1x10<sup>-09</sup></b>	0.77				
vR111A	0.38	0.18	0.68	0.67	<b>0.016</b>	<b>0.00032</b>	<b>0.00094</b>			
N348A/Y211A	0.061	<b>0.013</b>	0.18	0.68	0.17	<b>5.2x10<sup>-06</sup></b>	<b>2.1<sup>-05</sup></b>	0.38		
N348W/Y211A	<b>0.00017</b>	<b>3.9x10<sup>-06</sup></b>	<b>0.0011</b>	<b>0.021</b>	0.68	<b>1.6x10<sup>-10</sup></b>	<b>6.3x10<sup>-10</sup></b>	<b>0.0047</b>	0.063	
vR111A/Y211A	0.56	0.34	0.86	0.47	<b>0.0072</b>	<b>0.0010</b>	<b>0.0025</b>	0.77	0.24	<b>0.0019</b>