

# Daily arXiv Report:

## Extreme Eigenvalues and Eigenvectors for Finite Rank Additive Deformations of Non-Hermitian Sparse Random Matrices

Report on arXiv:2602.20956 by W. Hachem, M. Louvaris, J. Najim

February 27, 2026

### 1 Core Question

Given a sparse non-Hermitian random matrix  $X_n$  perturbed by a deterministic finite-rank matrix  $E_n$ , do the outlier eigenvalues of  $Y_n = X_n + E_n$  still match those of  $E_n$ , and—in the rank-one case—does the associated eigenvector align with the “true” direction as in the classical Hermitian BBP transition?

### 2 Setup and Key Definitions

**Definition 1** (Sparse non-Hermitian random matrix). Let  $\chi$  be a complex random variable with  $\mathbb{E}\chi = 0$ ,  $\mathbb{E}|\chi|^2 = 1$ . Let  $A_n = (A_{ij})$  be  $n \times n$  with i.i.d. entries distributed as  $\chi$ . Let  $B_n = (B_{ij})$  be an independent  $n \times n$  matrix with i.i.d. Bernoulli( $K_n/n$ ) entries, where  $K_n \leq n$  and  $K_n \rightarrow \infty$ . Define the *sparse random matrix*

$$X_n = (X_{ij}^{(n)}), \quad X_{ij}^{(n)} = \frac{1}{\sqrt{K_n}} B_{ij} A_{ij}.$$

Note  $\mathbb{E}X_{11} = 0$  and  $\mathbb{E}|X_{11}|^2 = 1/n$ . The parameter  $K_n$  controls sparsity: when  $K_n = n$  we recover the dense i.i.d. model.

**Definition 2** (Finite-rank deformation). Fix  $r \geq 1$  and deterministic vectors  $u_{t,n}, v_{t,n} \in \mathbb{C}^n$  ( $t = 1, \dots, r$ ) with

$$\sum_{t=1}^r (\|u_{t,n}\| + \|v_{t,n}\|) \leq C$$

for some absolute constant  $C > 0$  (bounded operator-norm condition). Set

$$E_n = \sum_{t=1}^r u_{t,n} v_{t,n}^*, \quad Y_n = X_n + E_n.$$

**Definition 3** (Reverse characteristic polynomial).

$$q_n(z) = \det(I_n - z Y_n), \quad z \in \mathbb{D} = \{z \in \mathbb{C} : |z| < 1\}.$$

The zeros of  $q_n$  in  $\mathbb{D}$  are exactly  $1/\lambda$  for eigenvalues  $\lambda$  of  $Y_n$  with  $|\lambda| > 1$ , i.e. the *outliers*.

### 3 Concrete Example

Take the simplest case:  $r = 1$ ,  $u_n = v_n$  a unit vector, and  $E_n = \alpha u_n u_n^*$  with  $\alpha > 1$ . Then  $E_n$  has a single nonzero eigenvalue  $\alpha$  (an outlier outside the unit disk).

**Theorem (informal):** The matrix  $Y_n = X_n + \alpha u_n u_n^*$  has, with high probability, exactly one eigenvalue  $\lambda_{\max}(Y_n)$  near  $\alpha$ , and the squared projection of the corresponding eigenvector  $\tilde{u}_n$  onto  $u_n$  satisfies

$$|\langle \tilde{u}_n, u_n \rangle|^2 \xrightarrow{P} 1 - \frac{1}{\alpha^2}.$$

This is the *same formula* as in the Hermitian BBP transition for spiked Wigner matrices.

### 4 Main Results (Bulleted)

- **Asymptotic equivalence of the characteristic polynomial (Theorem 1.1).** Define

$$b_n(z) = \det(I - zE_n), \quad \kappa(z) = \sqrt{1 - z^2 \mathbb{E}A_{11}^2}, \quad F(z) = \sum_{k=1}^{\infty} \frac{z^k Z_k}{\sqrt{k}},$$

where  $(Z_k)$  are independent complex Gaussians with  $\mathbb{E}Z_k = 0$ ,  $\mathbb{E}|Z_k|^2 = 1$ ,  $\mathbb{E}Z_k^2 = (\mathbb{E}A_{11}^2)^k$ . Then as  $\mathcal{H}$ -valued (holomorphic functions on  $\mathbb{D}$ ) random variables:

$$q_n \sim b_n \kappa \exp(-F), \quad n \rightarrow \infty.$$

- **Outlier eigenvalue matching (Theorem 1.2).** Under the *gap condition* (no eigenvalue of  $E_n$  in the annulus  $1 < |z| < 1 + \varepsilon$ ), the number of eigenvalues of  $Y_n$  outside the disk of radius  $1 + \varepsilon$  equals the number of eigenvalues of  $E_n$  outside the unit disk, w.h.p., and the Hausdorff distance between the two sets converges to zero.
- **Spectral radius convergence (Theorem 1.4).**  $\rho(X_n) \xrightarrow{P} 1$ , generalising the dense result of Bordenave et al. (2022) to all sparsity regimes  $K_n \rightarrow \infty$ .
- **Eigenvector projection for rank-one deformation (Theorem 1.6).** If additionally the entries are sub-Gaussian and  $K_n \gg \log^9 n$ , then for the unit eigenvector  $\tilde{u}_n$  of  $Y_n$  at  $\lambda_{\max}$ :

$$\left| \langle \tilde{u}_n, u_n / \|u_n\| \rangle \right|^2 - \left( 1 - \frac{1}{|\langle u_n, v_n \rangle|^2} \right) \xrightarrow{P} 0.$$

### 5 Intuition

The reverse characteristic polynomial  $q_n(z)$  encodes the outlier eigenvalues of  $Y_n$  as its zeros in  $\mathbb{D}$ . The key insight (following Bordenave–Chafaï–García-Zelada, 2022) is:

1.  $q_n$  factorises asymptotically as  $b_n(z)$  (the deterministic part from  $E_n$ ) times a *universal random analytic function*  $\kappa(z) \exp(-F(z))$  that depends only on the entry distribution, not on  $E_n$ .
2. Since  $\kappa \exp(-F)$  has *no zeros* in  $\mathbb{D}$ , the zeros of  $q_n$  are governed entirely by  $b_n$ , hence by  $E_n$ .
3. The eigenvector result requires a *universality comparison*: one replaces  $X_n$  by a Gaussian analogue (via the Brailovskaya–van Handel 2024 singular-value comparison), reducing the problem to known Gaussian computations.

## 6 Proof Sketch

### Step 1: Tightness of $(q_n)$ in $\mathcal{H}$

Using the multilinearity of the determinant and the singular-value decomposition  $I - zE_n = U\Sigma V^*$ , one shows  $\mathbb{E}|q_n(z)|^2 \leq C/(1-|z|^2)$ , which gives tightness in  $\mathcal{H}$  (the space of holomorphic functions on  $\mathbb{D}$  with the topology of uniform convergence on compacta).

### Step 2: Truncation

One may assume the entries  $A_{ij}$  are bounded a.s. (truncation at level  $D$ ) with negligible error as  $D \rightarrow \infty$ , uniformly in  $n$ .

### Step 3: Asymptotic equivalence $q_n \sim G_n = b_n \det(I - zX_n)$

Using the matrix determinant lemma (since  $E_n$  has rank  $r$ ):

$$q_n(z) = \det(I - zX_n) \det(I - z(I - zX_n)^{-1}E_n).$$

The second factor is an  $r \times r$  determinant involving the resolvent  $(I - zX_n)^{-1}$ . Via a careful analysis of the resolvent's bilinear forms  $v_t^*(I - zX_n)^{-1}u_s$ , one shows the second factor converges to  $b_n(z)/\det(I - zE_n)$ ... more precisely,  $q_n \sim b_n \det(I - zX_n)$ .

### Step 4: Asymptotic equivalence $\det(I - zX_n) \sim \kappa \exp(-F)$

This is the core random-analytic-function step. One shows the power-series coefficients of  $\log \det(I - zX_n)$  converge jointly to those of  $\log \kappa - F$  by computing moments of traces  $\text{tr}(X_n^k)$  and showing they match the Gaussian limit.

### Step 5: Outlier location via Rouché's theorem

With the asymptotic equivalence  $q_n \sim b_n \kappa \exp(-F)$  in hand, and since  $\kappa \exp(-F)$  is a.s. zero-free in  $\mathbb{D}$ , Rouché's theorem (via Skorokhod coupling) shows that for large  $n$ , the zeros of  $q_n$  in  $\{|z| \leq 1/(1+\varepsilon)\}$  are close to those of  $b_n$ , yielding Theorem 1.2.

### Step 6: Eigenvector projection (rank-one case)

- (a) **Gaussian comparison.** Under sub-Gaussianity and  $K_n \gg \log^9 n$ , the Brailovskaya–van Handel (2024) result implies that the singular-value distribution of  $zX_n + M$  is close to that of the Gaussian analogue  $zG_n + M$  for any deterministic  $M$ . This transfers eigenvector statistics.
- (b) **Reduction to resolvent.** The projection  $|\langle \tilde{u}_n, u_n / \|u_n\| \rangle|^2$  is expressed via the adjugate of  $(I - \lambda_{\max}^{-1} X_n)$  and the rank-one structure of  $E_n$ .
- (c) **Gaussian computation.** For the Gaussian analogue, one exploits rotational invariance and known formulas for the resolvent, yielding the limit  $1 - 1/|\langle u_n, v_n \rangle|^2$ .

## References

- Hachem, Louvaris, Najim. *Extreme eigenvalues and eigenvectors for finite rank additive deformations of non-Hermitian sparse random matrices*. arXiv:2602.20956, 2026.
- Bordenave, Chafaï, García-Zelada. *Convergence of the spectral radius of a random matrix through its characteristic polynomial*. Probab. Theory Related Fields, 2022.

Brailovskaya, van Handel. *Universality and sharp matrix concentration inequalities*. Geom. Funct. Anal., 2024.

Tao. *Outliers in the spectrum of iid matrices with bounded rank perturbations*. Probab. Theory Related Fields, 2013.

Baik, Ben Arous, Pécché. *Phase transition of the largest eigenvalue for nonnull complex sample covariance matrices*. Ann. Probab., 2005.