

# Paper Report: The Eigenvalues of I.I.D. Matrices Are Hyperuniform

G. Cipolloni, L. Erdős, O. Kolupaiev (arXiv:2602.17628)

Auto-generated report

February 24, 2026

## 1 Core Question

Given an  $N \times N$  random matrix  $X$  with independent, identically distributed (i.i.d.) entries, do its eigenvalues exhibit *hyperuniformity*—i.e., does the variance of the number of eigenvalues in a nice subdomain  $\Omega$  of the spectrum grow *strictly slower* than the expected count  $\mathbb{E}[N_\Omega] \sim N$ ?

## 2 Main Statement

**Definition 1** (I.I.D. matrix ensemble). An  $N \times N$  matrix  $X$  belongs to the *i.i.d. ensemble* if its entries  $x_{ab} = N^{-1/2}\chi$  are i.i.d. copies of a centered random variable  $\chi$  with  $\mathbb{E}|\chi|^2 = 1$  (and  $\mathbb{E}\chi^2 = 0$  in the complex case), satisfying moment bounds  $\mathbb{E}|\chi|^p \leq C_p$  for all  $p$ .

**Definition 2** (Hyperuniformity (Torquato)). A point process in  $\mathbb{R}^d$  is *hyperuniform* if the variance of the number of points in a ball of radius  $R$  grows strictly slower than its volume  $R^d$  as  $R \rightarrow \infty$ .

**Theorem 3** (Complex case, Theorem 2.4 in the paper). *Let  $X$  be a complex i.i.d. matrix satisfying the assumptions above. Fix  $\alpha \in [0, 1/2)$  and let  $\Omega_N \subset \mathbb{D}$  (the open unit disk) be a “nice” domain of diameter  $\sim N^{-\alpha}$  with  $C^2$  boundary, contained well inside  $\mathbb{D}$ . Define*

$$N_\Omega := \#\{\sigma_i \in \Omega_N\},$$

where  $\sigma_1, \dots, \sigma_N$  are the eigenvalues of  $X$ . Then for any  $\xi > 0$ ,

$$\text{Var}(N_\Omega) \leq N^{1-2\alpha-q_0(1/2-\alpha)+\xi}, \quad q_0 = \frac{1}{20}.$$

In particular, for macroscopic domains ( $\alpha = 0$ ) one obtains  $\text{Var}(N_\Omega) \leq N^{1-1/40+\xi}$ , which is  $\ll N \sim \mathbb{E}[N_\Omega]$ .

**Theorem 4** (Real case, Theorem 2.6). *Under the same setup but with real entries (and a transversality condition on  $\partial\Omega_N$  at the real axis),*

$$\text{Var}(N_\Omega) \leq N^{1-2\alpha-q_0(1/2-\alpha)+\xi} + N^{1-2\alpha-q_1+\xi}, \quad q_0 = \frac{1}{20}, \quad q_1 = \frac{1}{106}.$$

## 3 Concrete Example

Take  $X$  to be the *complex Ginibre ensemble*:  $X_{ij} = N^{-1/2}g_{ij}$  with  $g_{ij} \sim \mathcal{N}_{\mathbb{C}}(0, 1)$  i.i.d. The eigenvalues are known to form a determinantal point process on  $\mathbb{D}$ , and for that ensemble the exact number variance is  $\text{Var}(N_\Omega) \sim \sqrt{N}$  (“class I hyperuniform”).

For a *general* i.i.d. matrix—say with Bernoulli  $\pm 1$  entries—no determinantal structure is available. The paper proves, for the first time, that the eigenvalue process is still hyperuniform:  $\text{Var}(N_\Omega) \leq N^{1-1/40+\xi}$ , a power-law improvement over the trivial Poisson bound  $\sim N$ .

## 4 Intuition and Setup

**Circular law.** By the circular law, the empirical spectral measure of  $X$  converges to the uniform measure on  $\mathbb{D}$ , so  $N_\Omega \approx N|\Omega|/\pi$ . Fluctuations around this are the object of study.

**Linear statistics and Girko's formula.** Write  $N_\Omega - \mathbb{E}N_\Omega = L_N(\mathbf{1}_\Omega)$  as a *linear statistic*. For smooth test functions  $f$ , the variance of  $L_N(f)$  is  $O(1)$  (proven in prior work). But  $\mathbf{1}_\Omega$  is not smooth, and approximating it by smooth functions  $f^\pm$  on a sub-microscopic scale  $N^{-a}$  ( $a > 1/2$ ) introduces large Laplacian norms  $\|\Delta f\|_\infty \sim N^{2a}$ .

**Girko's formula** converts eigenvalue statistics to resolvents of the *Hermitization*:

$$H_z := \begin{pmatrix} 0 & X - z \\ (X - z)^* & 0 \end{pmatrix}, \quad G_z(w) := (H_z - w)^{-1}.$$

Then  $\sum_i f(\sigma_i) = \frac{i}{4\pi} \int_{\mathbb{C}} \Delta f(z) \int_0^\infty \text{Tr} G_z(i\eta) d\eta d^2z$ . The variance of  $L_N(f)$  thus reduces to bounding  $\text{Cov}(\langle G_{z_1}(i\eta_1) \rangle, \langle G_{z_2}(i\eta_2) \rangle)$ .

**Why existing bounds are insufficient.** Prior work (Cipolloni–Erdős–Schröder, 2022) gave

$$\text{Cov}(\langle G_1 \rangle, \langle G_2 \rangle) = \frac{\text{Main Term}}{N^2} + O\left(\frac{1}{\sqrt{N}\eta_*(N\eta_1\eta_2)^2}\right),$$

but the error lacks decay in  $|z_1 - z_2|$ , which is essential for the double integral over  $(z_1, z_2)$ .

## 5 Main Results (Bulleted)

- **Hyperuniformity for complex i.i.d. matrices:**  $\text{Var}(N_\Omega) \leq N^{1-1/40+\xi}$  (macroscopic  $\Omega$ ).
- **Hyperuniformity for real i.i.d. matrices:**  $\text{Var}(N_\Omega) \leq N^{1-1/106+\xi}$ , even when  $\Omega$  intersects the real axis (new even for real Ginibre).
- **Mesoscopic domains:** Result extends to  $N$ -dependent domains  $\Omega_N$  of diameter  $N^{-\alpha}$ ,  $0 \leq \alpha < 1/2$ .
- **New two-resolvent local law:** For two distinct spectral parameters  $z_1, z_2$  and deterministic matrices  $B_1, B_2$ ,

$$\langle G_{z_1} B_1 G_{z_2} B_2 \rangle = \text{deterministic approximation} + O_{\prec}\left(\frac{1}{N(|z_1 - z_2|^2 + \eta_1 + \eta_2)}\right).$$

- **Improved resolvent covariance (Ginibre-type bound):**

$$\text{Cov}(\langle G_1 \rangle, \langle G_2 \rangle) = \frac{\text{Main}}{N^2} + O\left(\frac{1}{N(|z_1 - z_2|^2 + \eta_1 + \eta_2)(N\eta_1\eta_2)^2}\right).$$

- **Singular vector overlaps:** For  $|z_1|, |z_2| \leq 1 - \delta$ ,

$$|\langle u_i^{z_1}, u_j^{z_2} \rangle|^2 + |\langle v_i^{z_1}, v_j^{z_2} \rangle|^2 \prec \frac{1}{N(|z_1 - z_2|^2 + N^{-1}|i - j|)},$$

identifying decorrelation in the singular-vector index  $|i - j|$  for the first time.

## 6 Proof Sketch

**Step 1. Portmanteau smoothing.** Approximate  $\mathbf{1}_\Omega$  from above/below by smooth functions  $f_{a,N}^\pm := \varphi_N^\pm * \omega_{a,N}$ , where  $\omega_{a,N}(z) = N^{2a}\omega(N^a z)$  is a bump function at scale  $N^{-a}$ ,  $a > 1/2$ , and  $\varphi_N^\pm$  are indicators of  $\Omega_N$  enlarged/shrunk by  $N^{-a}$ . By the Portmanteau lemma:

$$\text{Var}(N_\Omega) \leq 2[\text{Var}(L_N(f^+)) + \text{Var}(L_N(f^-))] + (\mathbb{E}L_N(f^+ - f^-))^2.$$

**Step 2. Expectation computation (Proposition 3.1).** Using Girko's formula and existing local laws, show

$$\mathbb{E}L_N(f_{a,N}) = \frac{N|\Omega_N|}{\pi}(1 + O(N^{-c})),$$

for  $a \in [1/2, 1/2 + 1/14]$ . This controls the “discrepancy” term  $(\mathbb{E}L_N(f^+ - f^-))^2 \lesssim N^{2(1-a-\alpha)}$ .

**Step 3. Variance bound via Girko (Proposition 3.2).** By Girko's formula,  $\text{Var}(L_N(f))$  is expressed as a double integral of  $\text{Cov}(\langle G_{z_1} \rangle, \langle G_{z_2} \rangle)$  weighted by  $\Delta f(z_1)\Delta f(z_2)$ . Split the  $\eta$ -integrals into three regimes:

- *Sub-microscopic* ( $\eta \ll 1/N$ ): controlled by lower-tail bounds on the smallest singular value of  $X - z$ .
- *Microscopic* ( $\eta \sim 1/N$ ): handled via Dyson Brownian motion (DBM) comparison with the Ginibre ensemble.
- *Macro-mesoscopic* ( $\eta \gg 1/N$ ): requires the new resolvent covariance estimate—the main novelty.

**Step 4. Chaos expansion for the covariance (Section 7).** To prove the improved covariance bound with  $|z_1 - z_2|^2$  decay:

- Start from the trivial bound  $\text{Cov}(\langle G_1 \rangle, \langle G_2 \rangle) \prec (N\eta_1\eta_2)^{-1}$  (from single-resolvent local law).
- Perform iterative cumulant expansions: each step gains a factor  $(N\eta_*)^{-1}$  (small in the mesoscopic regime  $\eta_* \geq N^{-1+\epsilon}$ ), but also generates higher-order covariances  $\text{Cov}(\langle (G_1 - M_1) \cdots \rangle, \langle G_2 \rangle)$ .
- These form a *hierarchy*; after  $\sim 1/\epsilon$  iterations, truncate using multi-resolvent local laws.
- The accumulated  $(N\eta_*)^{-1/\epsilon}$  factors compensate the truncation error, yielding the  $|z_1 - z_2|^2 + \eta_1 + \eta_2$  decay.

**Step 5. Two-resolvent local law (Section 4 & Supplementary S2).** Prove a new *averaged* local law for products  $\langle G_{z_1} B_1 G_{z_2} B_2 \rangle$  with  $z_1 \neq z_2$ . The key difficulty: unlike Wigner resolvents, there is no resolvent identity linking  $G_{z_1}$  and  $G_{z_2}$  when  $z_1 \neq z_2$ , so the decorrelation factor does not appear automatically. Instead, it emerges from a *two-body stability analysis* of the matrix Dyson equation (Appendix A).

**Step 6. DBM interpolation for the microscopic regime.** Interpolate between the general i.i.d. matrix  $X$  and a Ginibre matrix  $X_{\text{Gin}}$  via Ornstein–Uhlenbeck flow  $X_t = e^{-t}X + \sqrt{1 - e^{-2t}}X_{\text{Gin}}$ . Use the new two-resolvent local law to show that the resolvent covariance of  $X_t$  at time  $t \sim N^{-1+\epsilon}$  is close to that of  $X$ , while at the Ginibre end exact formulas give sharp bounds.

**Step 7. Optimization.** The parameter  $a$  (smoothing scale) is optimized: the variance terms scale as  $N^{2(a-\alpha)-2q_0(1/2-\alpha)}$  while the expectation discrepancy scales as  $N^{2(1-a-\alpha)}$ . Balancing gives the final exponent  $q = q_0(1/2 - \alpha)$  in the complex case.

**Step 8. Real case (Supplementary S3).** Additional complications from eigenvalues on/near the real axis. Use a refined analysis near  $\Im z = 0$ , where the density of real eigenvalues is  $\sim \sqrt{N}$ . An extra error term with exponent  $q_1 = 1/106$  appears.

## References

G. Cipolloni, L. Erdős, O. Kolupaiev, *The eigenvalues of i.i.d. matrices are hyperuniform*, arXiv:2602.17628, February 2026.  
<https://arxiv.org/abs/2602.17628>