

Report: Bulk–Boundary Correspondence in 2D Non-Hermitian Systems

via Toeplitz Operators and Singular Values

arXiv:2602.13916 — J. Sirker

Daily arXiv Report

February 20, 2026

1 Core Question

How should one formulate the bulk–boundary correspondence for two-dimensional non-Hermitian lattice Hamiltonians, given that the eigenvalue spectrum is generically unstable under changes of boundary conditions and therefore cannot serve as the basis for topological classification? The paper argues that *singular values*—not eigenvalues—provide the correct, stable foundation and develops the full theory using Toeplitz operator index theorems and K -splitting results.

2 Main Statement (Informal)

Let H be a two-dimensional translation-invariant non-Hermitian lattice Hamiltonian with scalar symbol $F(k_x, k_y)$ satisfying a point gap ($\det F(k_x, k_y) \neq 0$ for all $(k_x, k_y) \in \mathbb{T}^2$). Define the *slice winding numbers*

$$I_\alpha = \frac{1}{2\pi i} \int_0^{2\pi} dk_\alpha \partial_{k_\alpha} \ln \det F(k_x, k_y), \quad \alpha \in \{x, y\}.$$

Then, for the finite system on an $N_x \times N_y$ lattice with periodic boundary conditions in the x -direction and open boundary conditions in the y -direction:

- (i) There are exactly $K = N_x |I_y|$ singular values of the finite Hamiltonian matrix H_L that are separated from the bulk singular-value spectrum by a gap $\Delta > 0$.
- (ii) These K singular values converge to zero as $N_y \rightarrow \infty$ (exponentially in the clean case, polynomially under random perturbations).
- (iii) The corresponding singular vectors are exponentially localized at the boundary.
- (iv) The gap Δ is stable against any perturbation δH with $\|\delta H\|_2 < \Delta$.

For matrix-valued symbols, $K \geq N_x |I_y|$ (lower bound only). Corner modes require gapped edges and are controlled by a distinct corner Fredholm index.

3 A Concrete Example: The 2D Hatano–Nelson Model

Consider the non-reciprocal hopping model on a square lattice:

$$H = \sum_{i,j} (t_R c_{i+1,j}^\dagger c_{i,j} + t_L c_{i,j}^\dagger c_{i+1,j} + t_D c_{i,j+1}^\dagger c_{i,j} + t_U c_{i,j}^\dagger c_{i,j+1}),$$

where $t_R, t_L, t_U, t_D > 0$ are generally different (non-reciprocal) hopping amplitudes. Its **symbol** (Bloch Hamiltonian) is:

$$F(k_x, k_y) = t_L e^{-ik_x} + t_R e^{ik_x} + t_U e^{-ik_y} + t_D e^{ik_y}.$$

This traces out two ellipses E_x and E_y in the complex plane with semi-axes $|t_R \pm t_L|$ and $|t_D \pm t_U|$, respectively. The point gap is open when one ellipse is contained inside the other:

- $E_x \subset E_y$: $(I_x, I_y) = (0, \pm 1)$, giving N_x protected edge modes along the y -boundaries.
- $E_y \subset E_x$: $(I_x, I_y) = (\pm 1, 0)$, giving N_y protected edge modes along the x -boundaries.

With parameters $t_R = 0.5$, $t_L = 0.9$, $t_U = 1.2$, $t_D = 0.5$, one gets $(I_x, I_y) = (0, -1)$. The paper demonstrates that the N_x smallest singular values decay to zero with N_y , while the eigenvalue spectrum is completely destabilized by even a tiny perturbation ($\varepsilon = 0.1$).

4 Key Definitions and Setup

Definition 1 (Singular values). *For a (possibly non-Hermitian) matrix H , the singular values are $\sigma_i(H) = \sqrt{\lambda_i(H^\dagger H)}$, always real and non-negative. For Hermitian H , $\sigma_i = |\lambda_i|$; for non-normal H , the singular-value and eigenvalue spectra can behave completely differently.*

Definition 2 (Toeplitz operator). *Given a translation-invariant lattice Hamiltonian on a semi-infinite lattice, the Hamiltonian matrix H has the form $H_{ij} = h_{i-j}$ (Toeplitz structure). The symbol is the Fourier transform $h(k) = \sum_j h_j e^{-ikj}$. In two dimensions, one obtains a block-Toeplitz-with-Toeplitz-blocks (BTB) matrix with two-dimensional symbol $F(k_x, k_y)$.*

Definition 3 (Index of a Toeplitz operator). *For a semi-infinite Toeplitz operator H :*

$$\text{ind}(H) = \dim(\ker H) - \dim(\ker H^\dagger).$$

By Gohberg's theorem, $-\text{ind}(H)$ equals the winding number of $\det h(k)$ around the origin.

Definition 4 (K -splitting property). *The singular-value spectrum of a finite truncation H_L has the K -splitting property if*

$$\lim_{L \rightarrow \infty} \sigma_n(H_L) = \begin{cases} 0 & 1 \leq n \leq K, \\ > 0 & n > K. \end{cases}$$

The K small singular values are separated from the bulk by a gap $\Delta > 0$. For scalar symbols, $K = |I|$ (exact count); for matrix symbols, $K \geq |I|$ (lower bound).

Why eigenvalues fail. For non-normal H , the Bauer–Fike theorem gives a perturbation bound $|\lambda - \mu| \leq \kappa(V) \|\delta H\|_2$ where $\kappa(V) = \|V\| \|V^{-1}\|$ can be arbitrarily large. By contrast, singular values always satisfy Weyl's inequality: $|\sigma_k(H + \delta H) - \sigma_k(H)| \leq \|\delta H\|_2$, ensuring Lipschitz stability regardless of non-normality.

5 Main Results

- **Half-plane (edge modes):** For a scalar symbol with point gap, the number of topologically protected edge modes on an $N_x \times N_y$ tube (PBC in x , OBC in y) is exactly $K = N_x |I_y|$. For matrix-valued symbols, $K \geq N_x |I_y|$. These modes are “hidden zero modes”: not eigenstates of the finite system, but vectors mapped exponentially close to zero (detectable only via singular values).

- **Quarter-plane with gapless edges:** When both $I_x \neq 0$ and $I_y \neq 0$, edge-mode families along both boundaries hybridize near corners. For scalar symbols, the generic result is edge-type states $|\Psi_{ij}| \sim |z|^i + |w|^j$ (additive, not multiplicative). True corner modes $|\Psi_{ij}| \sim |z|^i |w|^j$ require additional structure (e.g., symbol factorization) and are only *spectrally* protected, not index-protected.
- **Quarter-plane with gapped edges (higher-order topology):** When both the bulk and edges are gapped ($I_x = I_y = 0$ but a non-trivial corner Fredholm index exists), one obtains genuine topologically protected corner modes. The non-Hermitian BBH model provides a concrete example: sublattice symmetry $\{\Pi, h(k_x, k_y)\} = 0$ protects the corner index even when all crystalline symmetries are broken and non-reciprocal hoppings are introduced. The number of corner modes is $K = (n_x^L + n_x^R)(n_y^L + n_y^R)$, determined by four winding numbers $(I_1^x, I_2^x, I_1^y, I_2^y)$ of the one-dimensional sublattice-symmetric chains.
- **Stability:** All topologically protected singular values (and corresponding boundary/corner modes) are stable against perturbations δH with $\|\delta H\|_2 < \Delta$ (the singular-value gap). This is guaranteed by Weyl's inequality for singular values—no analogue holds for eigenvalues of non-normal operators.

6 Proof Sketch

1. **Establish spectral instability of eigenvalues.** Via the Bauer–Fike theorem, show that for non-normal H , the condition number $\kappa(V)$ of the eigenvector matrix can be arbitrarily large, so eigenvalue perturbation bounds are vacuous. The pseudospectrum $\sigma_\varepsilon(H) = \{\lambda : \|(H - \lambda I)^{-1}\| > 1/\varepsilon\}$ extends far from true eigenvalues, confirming instability.
2. **Prove stability of singular values.** Since $H^\dagger H$ is Hermitian positive semidefinite, Weyl's inequality directly applies: $|\sigma_k(H + \delta H) - \sigma_k(H)| \leq \|\delta H\|_2$. This is the key reason singular values are the correct spectral quantity for non-Hermitian topology.
3. **Set up Toeplitz structure.** Cast the translation-invariant bulk Hamiltonian as a (multi-level) Toeplitz operator. In 2D, this gives a BTTB matrix with symbol $F(k_x, k_y)$. The point-gap condition $\det F \neq 0$ ensures Fredholm property.
4. **Apply index theorems.** For the half-plane truncation (semi-infinite Toeplitz operator), Gohberg's theorem gives $\text{ind}(H) = -I$ where I is the winding of $\det h(k_y)$. Use the factorization $\det h(k_y) = \prod_{n=0}^{N_x-1} \det F(2\pi n/N_x, k_y)$ to show $I = N_x I_y$.
5. **Invoke K -splitting theorems.** Classical results in Toeplitz theory guarantee that for finite truncations H_L , exactly K singular values are separated from the bulk by a gap and converge to zero as $L \rightarrow \infty$. For scalar symbols, $K = |I|$ (using Coburn's lemma: $\ker H$ and $\ker H^\dagger$ cannot both be non-trivial). For matrix symbols, $K \geq |I|$.
6. **Handle the quarter-plane.** Two boundaries meeting at a corner require analyzing the quarter-plane Toeplitz operator. When edges are gapless, edge-mode families hybridize near corners; when edges are gapped ($I_x = I_y = 0$), the Hayashi corner index theorem applies, giving a finite-dimensional Fredholm kernel. The non-Hermitian BBH model is analyzed by showing the symbol factorizes as $h(k_x, k_y) = h_x(k_x) \otimes I + \sigma_3 \otimes h_y(k_y)$, reducing the corner problem to a product of two 1D edge problems.
7. **Verify with explicit recurrence relations.** For the 2D Hatano–Nelson model, solve the bulk recursion $t_D \Psi_{j-1} + \mu_k \Psi_j + t_U \Psi_{j+1} = 0$ and use the argument principle to show that the number of roots inside the unit disk equals $1 - I_y(k)$, directly connecting the winding number to the existence of normalizable edge solutions.

8. **Numerical confirmation.** All analytical predictions are verified numerically: (a) K -splitting with correct mode counts, (b) exponential (clean) and polynomial (disordered) decay of protected singular values, (c) stability of edge/corner localization under perturbations respecting the relevant symmetry, and (d) instability of the eigenvalue spectrum under the same perturbations.