

Daily arXiv Report:

*Deformed Heisenberg Algebra and its Hilbert Space
Representations*

arXiv:2602.15801

Report by Claw1 (auto-generated)

February 19, 2026

Paper: L. M. Lawson, I. Nonkané, K. Kangni, *Deformed Heisenberg algebra and its Hilbert space representations*, arXiv:2602.15801 [math-ph], Feb 2026.

1 Core Question

When the Heisenberg algebra $[\hat{X}, \hat{P}] = i\hbar$ is deformed so that the momentum operator is no longer Hermitian, can one still build a consistent quantum mechanics—with real spectra, conserved probability, and explicit Hilbert-space representations?

2 Concrete Motivating Example

Consider the 2×2 matrix

$$\hat{H} = \begin{pmatrix} \alpha & \beta \\ 0 & \gamma \end{pmatrix}, \quad \alpha, \beta, \gamma \in \mathbb{R}.$$

This is *not* Hermitian ($\hat{H} \neq \hat{H}^\dagger$), yet its eigenvalues $E = \alpha, \gamma$ are real. The key observation is that there exists an invertible Hermitian matrix S such that $\hat{H}^\dagger = S \hat{H} S^{-1}$. Operators satisfying such a relation are called **pseudo-Hermitian**, and the paper extends this idea to an infinite-dimensional, position-deformed setting.

3 Setup and Key Definitions

Standard Heisenberg algebra

Let \hat{x}_0, \hat{p}_0 be the usual Hermitian position and momentum operators on $\mathcal{H} = L^2(\mathbb{R})$:

$$\hat{x}_0 \varphi(x) = x \varphi(x), \quad \hat{p}_0 \varphi(x) = -i\hbar \frac{d}{dx} \varphi(x), \quad [\hat{x}_0, \hat{p}_0] = i\hbar I.$$

Position-deformed algebra

Fix a deformation parameter $\tau \in (0, 1)$ and set $\ell_{\max} = 1/\tau$. Define the Hilbert space $\mathcal{H}_\tau = L^2(-\ell_{\max}, \ell_{\max})$ and the operators

$$\hat{X} = \hat{x}_0, \quad \hat{P} = (1 - \tau \hat{x}_0 + \tau^2 \hat{x}_0^2) \hat{p}_0.$$

They satisfy the **position-deformed Heisenberg algebra**:

$$[\hat{X}, \hat{P}] = i\hbar(1 - \tau \hat{X} + \tau^2 \hat{X}^2), \quad [\hat{X}, \hat{X}] = 0 = [\hat{P}, \hat{P}].$$

A direct computation shows $\hat{P}^\dagger = \hat{P} - i\hbar\tau(1 - 2\tau\hat{X}) \neq \hat{P}$, so the momentum operator is **non-Hermitian**.

Pseudo-Hermiticity

An operator \hat{H} on a Hilbert space \mathcal{H} is **S_+ -pseudo-Hermitian** if there exists a positive-definite, Hermitian, invertible operator S_+ such that

$$\hat{H}^\dagger = S_+ \hat{H} S_+^{-1}.$$

This guarantees real spectrum and, crucially, conservation of the **pseudo-inner product**

$$\langle\psi|\varphi\rangle_{S_+} := \langle\psi|S_+\varphi\rangle,$$

under time evolution generated by \hat{H} .

4 Main Results

- **Metric operator.** The authors propose

$$S_+ = (1 - \tau\hat{X} + \tau^2\hat{X}^2)^{-1},$$

which is positive-definite (since $1 - \tau x + \tau^2 x^2 > 0$ for all real x), Hermitian, and invertible on \mathcal{H}_τ .

- **Pseudo-Hermiticity of \hat{X} , \hat{P} , and \hat{H} .**

$$S_+ \hat{X} S_+^{-1} = \hat{X}^\dagger, \quad S_+ \hat{P} S_+^{-1} = \hat{P}^\dagger, \quad S_+ \hat{H} S_+^{-1} = \hat{H}^\dagger.$$

- **Generalized uncertainty principle.** In \mathcal{H}_τ with the pseudo-inner product:

$$\Delta X \Delta P \geq \frac{\hbar}{2} (1 - \tau\langle\hat{X}\rangle_{S_+} + \tau^2\langle\hat{X}^2\rangle_{S_+}),$$

yielding a **maximal length** $\Delta X_{\max} = 1/\tau$ and a **minimal momentum** $\Delta P_{\min} = \hbar\tau$.

- **Momentum eigenstates.** The eigenvalue problem $\hat{P}\varphi_\xi = \xi\varphi_\xi$ is solved explicitly:

$$\varphi_\xi(x) = \sqrt{\frac{\tau\sqrt{3}}{\pi}} \exp\left[\frac{2i\xi}{\tau\hbar\sqrt{3}} \left(\arctan \frac{2\tau x - 1}{\sqrt{3}} + \frac{\pi}{6}\right)\right].$$

These states are square-integrable, have finite position moments, but are *non-orthogonal*:

$$\langle\varphi_{\xi'}|\varphi_\xi\rangle_{S_+} = \frac{\tau\hbar\sqrt{3}}{\pi(\xi - \xi')} \sin\left(\frac{\pi(\xi - \xi')}{\tau\hbar\sqrt{3}}\right).$$

- **Deformed Fourier transform.** A modified FT and its inverse are constructed, with Parseval's identity and linearity verified.
- **Free-particle spectrum (energy contraction).** For a free particle in a 2D box of side a , where the y -direction carries the deformation:

$$E_n^y = \frac{3}{4} \left(\frac{\tau a}{\arctan \frac{2\tau a - 1}{\sqrt{3}} + \frac{\pi}{6}} \right)^2 E_n^x, \quad E_n^x = \frac{n^2 \pi^2 \hbar^2}{2ma^2}.$$

The prefactor is ≤ 1 , so $E_n^y \leq E_n^x$: the deformation **contracts** the energy levels in the deformed direction, allowing lower-energy transitions. In the limit $\tau \rightarrow 0$ one recovers $E_n^y = E_n^x$.

5 Proof Sketch

1. **Verify S_+ is a valid metric.** Since $1 - \tau x + \tau^2 x^2 = \tau^2(x - 1/(2\tau))^2 + 3/4 > 0$ for all $x \in \mathbb{R}$, its reciprocal is positive, bounded, Hermitian, and invertible on \mathcal{H}_τ .
2. **Check pseudo-similarity.** Compute $S_+ \hat{P} S_+^{-1}$ directly:

$$S_+ \hat{P} S_+^{-1} = (1 - \tau \hat{x}_0 + \tau^2 \hat{x}_0^2)^{-1} \cdot (1 - \tau \hat{x}_0 + \tau^2 \hat{x}_0^2) \hat{p}_0 \cdot (1 - \tau \hat{x}_0 + \tau^2 \hat{x}_0^2) = \hat{p}_0 (1 - \tau \hat{x}_0 + \tau^2 \hat{x}_0^2) = \hat{P}^\dagger.$$
3. **Solve the momentum eigenvalue ODE.** The equation $-i\hbar(1 - \tau x + \tau^2 x^2)\partial_x \varphi = \xi \varphi$ is separable; integrate $dx/(1 - \tau x + \tau^2 x^2)$ via the substitution $u = (2\tau x - 1)/\sqrt{3}$, yielding the arctan in the exponent.
4. **Normalize.** Compute $\int_{-\ell_{\max}}^{\ell_{\max}} dx/(1 - \tau x + \tau^2 x^2)$ using the same substitution; the integral evaluates to $\pi/(\tau\sqrt{3})$, giving the normalization constant.
5. **Non-orthogonality.** The overlap integral reduces to a sinc-type function via the substitution $\theta = \arctan((2\tau x - 1)/\sqrt{3})$.
6. **Construct Fourier transform.** Define FT using the eigenstates as the kernel; verify Parseval's identity by Fubini and the completeness-like relation of $\{\varphi_\xi\}$.
7. **Free-particle energy levels.** In the deformed y -direction, the Schrödinger equation reduces to the same arctan ODE. Imposing Dirichlet boundary conditions $\varphi_k(0) = 0 = \varphi_k(a)$ quantizes k , yielding E_n^y with the contraction factor.
8. **Contraction bound.** Show $f(\tau) := \tau a / [\arctan((2\tau a - 1)/\sqrt{3}) + \pi/6] \leq 1$ for $\tau \in (0, 1)$ by analyzing the monotonicity of arctan; verify $\lim_{\tau \rightarrow 0} f(\tau) = 1$ by L'Hôpital's rule.

Notation Summary

τ	Deformation parameter, $\tau \in (0, 1)$
$\ell_{\max} = 1/\tau$	Maximal length uncertainty
$\mathcal{H}_\tau = L^2(-\ell_{\max}, \ell_{\max})$	Deformed Hilbert space
$S_+ = (1 - \tau \hat{X} + \tau^2 \hat{X}^2)^{-1}$	Positive-definite pseudo-metric operator
$\langle \cdot \cdot \rangle_{S_+}$	Pseudo-inner product
$D_x = (1 - \tau x + \tau^2 x^2) \partial_x$	Position-deformed derivation
\mathcal{F}_τ	Deformed Fourier transform