# Daily arXiv Report: Convergence Rate of the Diagonal-Valued Cauchy Transform for Permutation Invariant Random Matrices

Auto-generated report – arXiv:2603.02068
by Alexis Imbert

March 5, 2026

## 1 Core Question

Given two large $N \times N$ random Hermitian matrices $A$ and $B$, one of which is *permutation invariant* (i.e. its law is unchanged under conjugation by permutation matrices), how fast does the *diagonal-valued Cauchy transform* of $A + B$ converge to that of the operator-valued free sum $a \boxplus_\Delta b$, as $N \to \infty$?

In the classical (unitarily invariant) setting the convergence rate of the *scalar* Cauchy transform is $N^{-1}$. This paper establishes the first *quantitative* bounds in the more general permutation-invariant regime, where the relevant algebraic structure is freeness with amalgamation over the diagonal algebra.

## 2 Main Results

### Setup and notation

- **Permutation invariance.** A random matrix $M \in M_N(\mathbb{C})$ is *permutation invariant* if for every $\sigma \in S_N$, the matrix $V_\sigma M V_\sigma^{-1}$ has the same law as $M$, where $V_\sigma$ is the permutation matrix of $\sigma$.
- **Diagonal algebra.** $\mathcal{D}_N(\mathbb{C})$ denotes the algebra of $N \times N$ diagonal matrices. The *diagonal projection* is $\Delta(M) = (M_{i,j}\,\delta_{i,j})_{1 \le i,j \le N}$.
- **Diagonal-valued Cauchy transform.** For a self-adjoint $X$ and a diagonal matrix $D$ with $\mathrm{Im}(D_{ii}) > 0$,
$$G_X(D) := \Delta\big((D - X)^{-1}\big).$$
When $D = zI$ for $z \in \mathbb{C} \setminus \mathbb{R}$ we write $G_X(z)$.
- **Operator-valued free sum $a \boxplus_\Delta b$.** The authors construct $a + b$ as the adjacency operator of an infinite locally-finite weighted graph (see Section 5 below), such that $a$ and $b$ are *free over* $\mathcal{D}_N$ in the $C^*$-operator-valued probability space $(A, \mathcal{D}, \Delta)$.
- **Frobenius norm.** $\|M\|_F = \sqrt{\mathrm{Tr}(MM^*)}$.

### Theorem 1 (General permutation invariant case)

**Theorem 1** (Theorem 2.6 in the paper)**.** *Let $A, B$ be random Hermitian matrices satisfying:*

1. ***Assumption 4.5*** *(traffic moment control) at speed $M$ with constants $h_1, h_2$;*

2. ***Assumption 4.6*** *(Frobenius/Schatten-$M$ norm control) at speed $M$ with constant $C$;*

3. *One of them is permutation invariant.*

*Let $0 \le c \le \frac{1}{h_1+h_2+6}$ and $|z|^2 > r_0^2 := \max\{4C, 2c^2\}$. Then*

$$\mathbb{E}\left[\frac{1}{N}\left\|G_{A+B}(z) - G_{a+b}(z)\right\|_F^2\right] \le \frac{4}{|z|^2}\left(\frac{r_0}{|z|}\right)^{2cM}.$$

*Since $M \le \frac{\log N}{\log\log N}$, the best achievable rate is $N^{-\beta/\log\log N}$ for an explicit $\beta > 0$.*

**Theorem 2 (Sparse & bounded case)**

**Theorem 2** (Theorem 2.7 in the paper)**.** *Additionally assume that $A, B$ are:*

1. *Uniformly bounded in operator norm: $\|A\|_{\mathrm{op}}, \|B\|_{\mathrm{op}} \le C$;*

2. *Uniformly sparse: $\sum_{j=1}^{N}(\mathbf{1}_{A_{ij}\neq 0} + \mathbf{1}_{B_{ij}\neq 0}) \le C$ for all $i$.*

*Let $\eta_0 > \|A\|_{\mathrm{op}} + \|B\|_{\mathrm{op}}$ and $D \in \mathcal{D}_N(\mathbb{C})$ with $\min_i \mathrm{Im}(D_{ii}) > \eta_0$, $d := \min_i |D_{ii}|$. Then*

$$\mathbb{E}\left[\frac{1}{N}\left\|G_{A+B}(D) - G_{a+b}(D)\right\|_F^2\right] \le \frac{6}{d^2}\, N^{-\frac{2\log 2C/d}{1+2\log C}}.$$

*This gives a* polynomial *rate $N^{-\gamma}$ and controls the Cauchy transform at any diagonal argument $D$, not just scalar multiples of the identity.*

# 3 Concrete Example

Consider two independent adjacency matrices of uniformly random $d$-regular graphs on $N$ vertices (with $d$ fixed). Each is permutation invariant, bounded in operator norm by $d$, and $d$-sparse. Theorem 2 applies and gives:

$$\mathbb{E}\left[\frac{1}{N}\|G_{A+B}(D) - G_{a+b}(D)\|_F^2\right] = O\big(N^{-\gamma}\big),$$

for an explicit $\gamma = \gamma(d, \eta_0) > 0$. This extends prior single-matrix results of Bordenave–Collins [4] to the *sum* of two independent random regular graphs.

# 4 Intuition and Setup

The classical Voiculescu theory says: if $A$ is *unitarily* invariant, then $A$ and any independent $B$ are asymptotically *free*, and the limiting spectral measure of $A + B$ is the free convolution $\mu_A \boxplus \mu_B$. The Cauchy transform converges at rate $N^{-1}$.

When $A$ is only *permutation* invariant (e.g. adjacency matrices of random graphs), ordinary freeness fails. Instead, $A$ and $B$ are asymptotically *free over the diagonal* $\mathcal{D}_N(\mathbb{C})$. The limiting object is the *operator-valued* free convolution $\boxplus_{\mathcal{D}_N}$, which retains more information (the full diagonal of the resolvent, not just its trace). The key question becomes: at what rate does this convergence happen?

The answer depends on two assumptions that quantify how "nice" the matrices are:

- **Assumption 4.5 (traffic moments):** For graph monomials of size $\le M$, the number of injective embeddings into the adjacency structure of $A$ (or $B$) grows in a controlled way. Satisfied by Wigner matrices, Erdős–Rényi graphs, sparse bounded matrices, and products thereof.

- **Assumption 4.6 (Frobenius norm):** $\mathbb{E}[\|A^M\|_F^2/N]$ stays bounded. This is more restrictive: it fails for heavy-tailed Wigner matrices but holds for Erdős–Rényi with $d \to \infty$ and for bounded sparse models.

## 5 Explicit Construction of $a \boxplus_\Delta b$

The free sum $a + b$ is built as the adjacency operator of an infinite locally-finite graph:

1. Start with vertex $x \in [N]$.
2. Attach a copy of the graph of $A$ at $x$ (edges colored blue/"$a$").
3. To each new vertex $j \neq x$ created, attach a copy of the graph of $B$ at $j$ (edges colored red/"$b$").
4. Continue alternating: at each vertex reached for the first time, attach the other graph.

This yields a tree-like structure when viewed at the level of colored components. The resulting Hilbert space is $\mathcal{H}_N^x = \ell^2(V_N^x)$ where $V_N^x$ consists of tuples $(j_1, \ldots, j_n, \varepsilon)$ with $\varepsilon \in \{a, b\}$ and $j_k \neq j_{k+1}$.

The operators $a, b \in B(\mathcal{H}_N^x)$ act within their respective colored components:

$$\langle a\, \delta_{v_1}, \delta_{v_2} \rangle = A_{j(v_2), j(v_1)}, \qquad v_1, v_2 \in R_a(v),$$

where $R_a(v)$ is the "$a$-component" containing $v$ and $j(\cdot)$ extracts the last index. By the tree structure, $a$ and $b$ are provably $\Delta$-free (Proposition 3.1).

## 6 Proof Sketch

Both theorems are proved via a **moment method**: expand the resolvent as a power series and compare term by term.

### Step 1: Series expansion of the resolvent

For $|z|$ large enough, expand

$$(zI - X)^{-1} = \sum_{k=0}^{\infty} \frac{X^k}{z^{k+1}}.$$

The diagonal-valued Cauchy transform becomes $G_X(z) = \sum_{k \geq 0} z^{-(k+1)}\, \Delta(X^k)$.

### Step 2: Moment matching up to order $M$

The key technical work shows that for $k \leq M$,

$$\mathbb{E}\big[\Delta\big((A + B)^k\big)\big] \approx \Delta\big((a + b)^k\big),$$

using the traffic distribution framework. For the sum $(A + B)^k$, the diagonal entry $[(A + B)^k]_{ii}$ counts weighted walks of length $k$ from $i$ to $i$ on $[N]$, alternating between edges of $A$ and $B$. The analogous count on the infinite tree graph of $a + b$ agrees up to corrections controlled by Assumption 4.5 (non-injective embeddings contribute lower-order terms by permutation invariance).

### Step 3: Remainder control (Theorem 1)

For $k > M$, Assumption 4.6 (Frobenius norm of $A^M$) gives

$$\frac{1}{N}\mathbb{E}\big[\|\Delta(X^k)\|_F^2\big] \leq C^k,$$

which makes the tail of the series geometrically small for $|z| > r_0$. The factor $(r_0/|z|)^{2cM}$ arises from summing the tail starting at order $M$.

### Step 4: Remainder control (Theorem 2)

When $\|A\|_{\mathrm{op}}, \|B\|_{\mathrm{op}} \leq C$ and both are $C$-sparse, the operator norm directly controls the series remainder without needing Assumption 4.6. The sparsity allows a refined counting of non-backtracking walks on the tree, improving the rate to $N^{-\gamma}$. The argument proceeds by:

1. Truncating the Neumann series at order $M_0 = c_0 \log N$;
2. Using $\|X\|_{\mathrm{op}} \leq C$ to bound $\|X^k\|_{\mathrm{op}} \leq C^k$;
3. Using sparsity to show that the number of *injective* closed walks of length $k$ from vertex $i$ is at most $C^k$, so the diagonal entry $[(A+B)^k]_{ii}$ is well controlled;
4. Combining to get the polynomial-in-$N$ convergence rate.

### Step 5: From moments to Cauchy transform

The moment-level bounds are assembled into the $L^2$-Frobenius bound on $G_{A+B}(z) - G_{a+b}(z)$ via the Parseval-type identity

$$\frac{1}{N}\|G_{A+B}(z) - G_{a+b}(z)\|_F^2 = \frac{1}{N}\sum_{i=1}^{N}\left|\sum_{k\geq 0}\frac{[\Delta((A+B)^k) - \Delta((a+b)^k)]_{ii}}{z^{k+1}}\right|^2.$$

## 7 Summary

This paper provides the first quantitative convergence rates for the *diagonal-valued* Cauchy transform of sums of permutation invariant random matrices toward their operator-valued free convolution. The two main results trade off generality against rate:

- General traffic/Frobenius assumptions $\Rightarrow$ sub-polynomial rate $N^{-\beta/\log\log N}$.
- Bounded operator norm + sparsity $\Rightarrow$ polynomial rate $N^{-\gamma}$.

The construction of the free sum as an adjacency operator of an explicit infinite graph is of independent interest.

**Reference:** A. Imbert, *Convergence rate of the diagonal-valued Cauchy-transform for permutation invariant random matrices*, arXiv:2603.02068, March 2026.