arXiv Report: Existence, Properties, and Parametric Inference for Possibly Hyperuniform Gibbs Perturbed Lattices

Auto-generated report for arXiv:2603.01858
J.-F. Coeurjolly and C. Renaud-Chan

March 4, 2026

1 Core Question

Can one build a flexible class of point processes that (i) retains a lattice backbone, (ii) allows Gibbs-type interactions (including hard-core repulsion), and (iii) may be *hyperuniform* — a property that neither classical Gibbs point processes nor simple perturbed lattices can simultaneously achieve?

2 Exact Main Result

Theorem 1 (Existence, Theorem 1 of the paper)**.** *Let $\mathcal{L} \subset \mathbb{R}^d$ be a full-rank lattice with fundamental domain D. Assume the Hamiltonian H satisfies* (H1)–(H4) *(stability, non-degeneracy, heredity, translation-invariance) and either*

- (H5.1) *finite range* + (Q1)[1] *(exponential moment of order* 1 *for the move distribution Q), or*
- (H5.2) *pairwise summability* + (Q2) *(bounded support of Q).*

Then the set $\mathcal{G}_{\mathcal{L}}(H, Q)$ of Gibbs measures is non-empty, and the associated Gibbs perturbed lattice

$$\Gamma = \{i + X_i + U : i \in \mathcal{L}\}, \qquad X \sim P \in \mathcal{G}_{\mathcal{L}}(H, Q), \quad U \sim \mathrm{Unif}(D),$$

is a stationary point process on $\mathbb{R}^d$. Moreover, for $d \leq 2$, under strengthened moment/boundedness conditions on Q, Γ is ***hyperuniform****:*

$$\frac{\mathrm{Var}[\#\Gamma \cap B(0,R)]}{|B(0,R)|} \longrightarrow 0 \quad \textit{as } R \to \infty.$$

3 Concrete Example

Strauss hard-core perturbed lattice. Take $\mathcal{L} = \mathbb{Z}^2$ and define the pairwise Hamiltonian

$$H(\gamma) = \sum_{\{x,y\} \subset \gamma} \Big[A\, \mathbf{1}_{[r,R]}(|x-y|) + \infty \cdot \mathbf{1}_{[0,r)}(|x-y|) \Big],$$

with interaction range $R > 0$, hard-core distance $r \geq 0$, and energy parameter $A \geq 0$. Choose the move distribution Q as the uniform distribution on a ball $B(0, \varepsilon)$ with $\varepsilon < (\delta - 2r)/2$ (where δ is the lattice spacing).

The resulting Gibbs perturbed lattice:

- has a *guaranteed* hard-core: no two points can be closer than r;

- exhibits Strauss-type soft repulsion at distances in $[r, R]$;
- is **hyperuniform** in $d = 2$ (by Proposition 3.6 of the paper).

None of these three properties can be simultaneously achieved by a standard Gibbs point process on $\mathbb{R}^2$ or a simple i.i.d. perturbed lattice.

4 Intuition and Setup

Key objects.

- **Lattice** $\mathcal{L} \subset \mathbb{R}^d$: a full-rank discrete subgroup generated by d linearly independent vectors $a_1, \ldots, a_d$. Fundamental domain $D = \left\{\sum t_i a_i : t_i \in [0, 1)\right\}$. Lattice spacing $\delta = \min\{|i| : i \in \mathcal{L} \setminus \{0\}\}$.
- **Move distribution** Q: a probability measure on $\mathbb{R}^d$ controlling how far each particle can be displaced from its lattice site.
- **Perturbed lattice**: $\Gamma = \{i + X_i + U : i \in \mathcal{L}\}$ where each $X_i \sim Q$ (not necessarily i.i.d.—the joint law is governed by P) and $U \sim \mathrm{Unif}(D)$ is an independent global shift that stationarizes Γ.
- **Hamiltonian** $H : \mathcal{C}_f \to \mathbb{R} \cup \{\infty\}$: measures interaction energy of a finite point configuration. The local energy of a point x given the rest of the configuration γ is $h(x, \gamma) = \lim_{n\to\infty}[H(\{x\} \cup \gamma_{[-n,n]^d}) - H(\gamma_{[-n,n]^d})]$.
- **Gibbs measure on** $\mathcal{L}$: the joint law P of $(X_i)_{i\in\mathcal{L}}$ satisfying DLR equations (1) below.

Why a new model class? Known hyperuniform processes (Ginibre, determinantal, i.i.d. perturbed lattices) lack flexibility: one cannot impose hard-core constraints. Conversely, Coeurjolly–Flint–Laverny (2024) showed that most Gibbs point processes on $\mathbb{R}^d$ are *not* hyperuniform. The authors bridge this gap by making the Hamiltonian act on the *perturbed locations* $\{i+X_i\}$ rather than on the marks $\{X_i\}$ alone, creating a model that is simultaneously:

1. lattice-based (hence potentially hyperuniform),
2. Gibbs-type (hence flexible, with tunable interactions).

5 Main Results (Summary)

- **Existence (Theorem 1).** Under stability, non-degeneracy, heredity, translation-invariance of H, plus a range/moment condition, infinite-volume Gibbs measures on $\mathcal{L}$ exist and yield stationary point processes.

- **Hyperuniformity (Proposition 3.6).**
  (i) Under (H5.1)+(Q1)[d] or (H5.2)+(Q2): *some* Gibbs perturbed lattice in $d \leq 2$ is hyperuniform.
  (ii) Under (H5.3)+(Q3) (bounded local energy + d-th moment): *every* Gibbs perturbed lattice in $d \leq 2$ is hyperuniform.

- **DLR-type equations (Propositions 3.7–3.9).** Three families of equilibrium equations (first-order, second-order, variational) analogous to GNZ equations for classical Gibbs point processes.

- **Parametric inference (Theorem 2).** For exponential-family models observed in an expanding window W_n:
  - A Takacs–Fiksel estimator $\hat{\theta}_n$ is defined, encompassing pseudo-likelihood and variational estimators as special cases.
  - $\hat{\theta}_n \to \theta^\star$ a.s. (consistency).

- $\sqrt{|W_n|}(\hat{\theta}_n - \theta^\star) \Rightarrow \mathcal{N}(0, \Sigma)$ (asymptotic normality), where Σ is an explicit sandwich-type covariance matrix.

- **Two observation frameworks:** (1) both lattice sites i and displacements X_i are observed; (2) only the perturbed locations $i + X_i$ are observed (harder; handled via the variational equation for specific move distributions).

6 Proof Sketch

Step 1: Finite-volume Gibbs measures

On a finite sub-lattice $\Lambda \subset \mathcal{L}$, define the "lifted" Hamiltonian $\tilde{H}(x_\Lambda) = H(\{i + x_i : i \in \Lambda\})$. The Gibbs measure on Λ is

$$P_\Lambda(dx_\Lambda) = \frac{1}{Z_\Lambda} e^{-\tilde{H}(x_\Lambda)} Q^{\otimes\Lambda}(dx_\Lambda).$$

Stability (H1) and non-degeneracy (H2) guarantee $Z_\Lambda \in (0, \infty)$. DLR equations on the finite lattice follow by elementary conditioning (Proposition 3.3).

Step 2: Infinite-volume limit (specific entropy)

The authors adapt the *specific entropy* approach of Georgii (2011). For an increasing sequence $\Lambda_n \uparrow \mathcal{L}$, they show:

1. The specific entropy $h(P \,|\, Q^{\otimes\mathcal{L}})$ is well-defined and lower-semicontinuous.
2. The specific energy $e(P) = \lim_n \frac{1}{|\Lambda_n|} E_P[\tilde{H}(X_{\Lambda_n})]$ exists under the stability/range assumptions.
3. The free-energy functional $f(P) = h(P \,|\, Q^{\otimes\mathcal{L}}) + e(P)$ is minimized by some $P^\star$, which is then shown to satisfy the DLR equations (1).

The DLR equations for the infinite lattice are:

$$P(dx'_\Delta \mid x_{\Delta^c}) = \frac{1}{Z_\Delta(x_{\Delta^c})} e^{-\tilde{H}_\Delta(x'_\Delta \cup x_{\Delta^c})} Q^{\otimes\Delta}(dx'_\Delta), \quad \forall \text{ finite } \Delta \subset \mathcal{L}. \tag{1}$$

Step 3: Hyperuniformity

The key input is the result of Klatt–Last–Yogeshwaran (2020): a perturbed lattice $\{i + X_i + U\}$ is hyperuniform in $d \leq 2$ whenever $E[|X_0|^d] < \infty$.

- Under (H5.1)+(Q1)[d]: the specific-entropy minimizer $P^\star$ inherits finite d-th moment from Q, giving at least one hyperuniform Gibbs perturbed lattice.
- Under (H5.3)+(Q3): the bounded local energy allows a uniform integrability argument showing $E_P[|X_0|^d] < \infty$ for *every* $P \in \mathcal{G}_\mathcal{L}(H, Q)$.

Step 4: DLR-type (GNZ-type) equations

Starting from the single-site DLR equation and applying it with test functions f, the authors derive:

- *First-order* equation (Proposition 3.7): an expectation identity involving the normalized Papangelou intensity $\Lambda(i, x, \gamma) = \lambda(i, x, \gamma) / \int_Q \lambda(i, y, \gamma)\, dy$.
- *Second-order* equation (Proposition 3.8): obtained by iterating the single-site DLR twice.
- *Variational* equation (Proposition 3.9): integration by parts on the mark space using the DLR kernel, yielding an identity involving ∇h and ∇q.

Step 5: Parametric inference

For exponential-family models $-\log \lambda(i, x, \gamma; \theta) = \theta^\top S(i, x, \gamma)$:

1. Define the Takacs–Fiksel contrast $\mathrm{TF}_n(f, \Gamma; \theta) = \sum_{l=1}^{L} [\mathrm{DLR}_n(f_l, \Gamma; \theta)]^2$ where DLR_n is the empirical DLR residual with border correction.
2. **Consistency:** Use a novel ergodic theorem for the empirical DLR functionals (adapting standard lattice ergodic theory to the continuous-mark, perturbed-location setting) to show $\mathrm{TF}_n/|W_n| \to \mathrm{TF}(\theta)$ uniformly, which has a unique minimum at $\theta^\star$.
3. **Asymptotic normality:** Taylor-expand $\mathrm{TF}_n^{(1)}$ around $\theta^\star$; use a CLT for the empirical DLR functionals (proved via cumulant bounds exploiting finite range) to obtain $\sqrt{|W_n|}(\hat{\theta}_n - \theta^\star) \Rightarrow \mathcal{N}(0, \Sigma)$ with sandwich covariance $\Sigma = I(\theta^\star)^{-1} V(\theta^\star) I(\theta^\star)^{-1}$.

Reference

J.-F. Coeurjolly and C. Renaud-Chan, *Existence, properties, and parametric inference for possibly hyperuniform Gibbs perturbed lattices*, arXiv:2603.01858, March 2026.