Daily arXiv Report:

Limit Theorems for Anisotropic Functionals of Stationary Gaussian Fields with Gneiting Covariance Function

Report on arXiv:2603.07188 by
N. Leonenko, L. Maini, I. Nourdin, F. Pistolato

March 10, 2026

1 Core Question

Given a stationary Gaussian field $B = (B_x)_{x\in\mathbb{R}^d}$ with a *non-separable* covariance of Gneiting type, what is the limiting distribution of the normalised nonlinear additive functional

$$\widetilde{Y}(t) = \frac{Y(t) - \mathbb{E}[Y(t)]}{\sqrt{\mathrm{Var}(Y(t))}}, \qquad Y(t) = \int_{t_1(t)D_1\times t_2(t)D_2} \varphi(B_x)\,dx,$$

as the two windows $t_1(t)D_1$ and $t_2(t)D_2$ grow at *different* (anisotropic) rates?

2 Setup and Key Definitions

Gaussian field and Hermite expansion. Let $B = (B_x)_{x\in\mathbb{R}^d}$ ($d = d_1 + d_2$, $d_i \geq 1$) be a continuous, centred, unit-variance, stationary Gaussian field with covariance $C(x) = \mathrm{Cov}(B_x, B_0)$. A square-integrable function $\varphi : \mathbb{R} \to \mathbb{R}$ admits the **Hermite expansion**

$$\varphi \overset{L^2}{=} \sum_{q\geq 0} a_q\, H_q, \qquad a_q = \frac{1}{q!}\,\mathbb{E}[H_q(N)\,\varphi(N)], \quad N \sim \mathcal{N}(0,1),$$

where H_q is the q-th Hermite polynomial (e.g. $H_0 = 1$, $H_1(x) = x$, $H_2(x) = x^2 - 1$). The **Hermite rank** R of φ is the smallest $q \geq 1$ with $a_q \neq 0$.

Gneiting covariance (Assumption 1.1). C belongs to the **Gneiting class** if

$$C(x_1, x_2) = C_2(x_2)\, C_1\Big(\frac{x_1}{C_2(x_2)^{1/d_1}}\Big), \qquad x = (x_1, x_2) \in \mathbb{R}^{d_1} \times \mathbb{R}^{d_2},$$

where C_1, C_2 are radial, c_1 (the radial part of C_1) is completely monotone, $C_2 > 0$, and $1/c_2$ has a completely monotone derivative. This is one of the few constructive recipes for valid *non-separable* space–time covariances on $\mathbb{R}^{d_1} \times \mathbb{R}^{d_2}$.

Anisotropic growth (Assumption 1.2). Let $D_i \subset \mathbb{R}^{d_i}$ be convex bodies and set $Y(t) = \int_{t_1 D_1\times t_2 D_2} \varphi(B_x)\,dx$ with $t_1, t_2 \to \infty$ at possibly different rates. Assumption 1.2 requires $t_1\, c_2(t_2)^{1/d_1} \to \infty$, ensuring the effective spatial rescaling diverges.

Regular variation. A function $G : \mathbb{R}^d \to \mathbb{R}$ is **regularly varying** at ∞ with index $-\rho < 0$ if $G(|x|) \sim |x|^{-\rho} L(|x|)$ as $|x| \to \infty$, where L is slowly varying ($L(\alpha x)/L(x) \to 1$).

3 Concrete Example

Consider a centred Gaussian field on $\mathbb{R}^{d+1}$ (space $\mathbb{R}^d$, time $\mathbb{R}$) with Gneiting covariance, C_1 and C_2 asymptotically power-law with exponents $-\rho_1$ and $-\rho_2$. The **excursion volume** above level $u > 0$,

$$M_T = \int_0^T \int_{T^\gamma K} \mathbf{1}(|B_x| \geq u)\, dx,$$

where $K \subset \mathbb{R}^d$ is a convex body and $\gamma \in (0, 1/\rho_2)$, has Hermite rank $R = 2$ (for $u > 0$). Depending on (ρ_1, ρ_2):

- If $C \in L^2$: $\mathrm{Var}(M_T) \asymp T^{\gamma d+1}$, CLT holds.
- If $C_1 \in L^2$, $C_2 \notin L^1$, $\rho_2 < 1$: $\mathrm{Var}(M_T) \asymp T^{\gamma d+2-\rho_2}$, CLT holds.
- If $C_1 \notin L^2$, $\rho_1 < d/2$, $C_2 \in L^{2(1-\rho_1/d)}$: $\mathrm{Var}(M_T) \asymp T^{2\gamma(d-\rho_1)+1}$, CLT holds.
- If both exhibit sufficient long-range dependence ($2\rho_1 < d$, $\rho_2 < \frac{d}{2(d-\rho_1)}$): **non-Gaussian** limit—a 2-domain Rosenblatt random variable.

4 Main Results

Throughout: C is Gneiting, Assumptions 1.1–1.2 hold, $R \geq 2$ is the Hermite rank of φ.

- **Theorem 1.1 (CLT, short-range in C_1).** If $C \in L^R(\mathbb{R}^d)$, or if $C_1 \in L^R(\mathbb{R}^{d_1})$ with C_2 regularly varying (even if $C_2 \notin L^{R-1}$), then

  $$\widetilde{Y}(t) \xrightarrow{d} \mathcal{N}(0, 1).$$

  Variance: $\mathrm{Var}(Y(t)) \sim \ell\, t_1^{d_1} t_2^{d_2}$ (first case) or grows super-linearly in the volume (second case).

- **Theorem 1.2 (CLT, long-range in C_1, short-range in C_2).** If $C_1 \notin L^R(\mathbb{R}^{d_1})$ with power-law exponent $-\rho_1$, $R\rho_1 < d_1$, and $C_2 \in L^{R(1-\rho_1/d_1)}(\mathbb{R}^{d_2})$, then

  $$\widetilde{Y}(t) \xrightarrow{d} \mathcal{N}(0, 1).$$

- **Theorem 1.3 (Non-CLT, $R = 2$, both long-range).** If $R = 2$, $C_1 \notin L^2(\mathbb{R}^{d_1})$ and $C_2 \notin L^1(\mathbb{R}^{d_2})$, both power-law with exponents $-\rho_1$, $-\rho_2$ satisfying $2\rho_1 < d_1$ and $\rho_2 < \frac{d_1 d_2}{2(d_1-\rho_1)}$, then

  $$\widetilde{Y}(t) \xrightarrow{d} \mathrm{sgn}(a_2)\, \mathcal{H}^{2, D_1 \times D_2}_{\rho_1, \rho_2(1-\rho_1/d_1)},$$

  a **2-domain Rosenblatt** random variable (second Wiener chaos, non-Gaussian). Variance: $\mathrm{Var}(Y(t)) \sim \ell_3\, t_1^{2d_1-2\rho_1}\, t_2^{2d_2-2\rho_2(1-\rho_1/d_1)}$.

- **Theorem 1.5 (Asymptotic separability transfer).** If C is asymptotically separable (Definition 1.13), then the second-chaos functional $\widetilde{Y}(t)$ has the *same* distributional limit as the analogous functional built from a truly separable covariance $C^{\mathrm{sep}} = C_1 \otimes C_2^*$.

5 The 2-Domain Rosenblatt Distribution

For $\alpha_i \in (0, 1)$ and convex bodies $D_i \subset \mathbb{R}^{d_i}$, the **2-domain Rosenblatt random variable** $\mathcal{H}^{\alpha, D_1 \times D_2}_{\rho_1, \rho_2}$ is defined as a double Wiener–Itô integral:

$$\mathcal{H} = I_2(f), \qquad f(y_1, y_2) = c \int_{D_1 \times D_2} \prod_{i=1}^{2} \|x_i - y_i\|^{-\alpha_i d_i}\, dx,$$

living in the second Wiener chaos of a white-noise field. Its cumulants are given by traces of powers of the associated Hilbert–Schmidt operator, and its characteristic function involves Fredholm determinants. It is the natural non-Gaussian limit for rank-2 functionals under anisotropic long-range dependence.

6 Proof Sketch

Step 1. Chaotic projection. Since φ has Hermite rank R, write $Y(t) = \sum_{q \geq R} a_q \int_{t_1 D_1 \times t_2 D_2} H_q(B_x)\, dx$. Each integral lives in the q-th Wiener chaos.

Step 2. Cumulant characterisation. For the dominant $q = R$ component (a second-chaos r.v. when $R = 2$), cumulants are

$$\kappa_k(\widetilde{Y}) = 2^{k-1}(k-1)!\, c_k(t),$$

where $c_k(t) = \frac{\int_{(t_1 D_1 \times t_2 D_2)^k} C(x^{(1)} - x^{(2)}) \cdots C(x^{(k)} - x^{(1)})\, dx}{\left(\int_{(t_1 D_1 \times t_2 D_2)^2} C^2(x^{(1)} - x^{(2)})\, dx\right)^{k/2}}$.

Step 3. Fourth Moment Theorem (CLT cases). By the Nualart–Peccati theorem, for a sequence in a fixed Wiener chaos, $\widetilde{Y}(t) \xrightarrow{d} \mathcal{N}(0,1)$ iff $\kappa_4(\widetilde{Y}(t)) \to 0$, i.e. $c_4(t) \to 0$. In Theorems 1.1 and 1.2, the authors show $c_4(t) \to 0$ by splitting the cyclic integral over $\mathbb{R}^{d_1}$ and $\mathbb{R}^{d_2}$ blocks and using integrability of C_1 (or C_2) to kill one factor.

Step 4. Asymptotic separability of Gneiting covariances. Key structural lemma: the Gneiting coupling $x_1/C_2(x_2)^{1/d_1}$ becomes negligible at large scales. Formally, for each $k \geq 2$,

$$c_k(t_1 D_1 \times t_2 D_2;\, C) \;\longrightarrow\; c_k(D_1;\, C_1) \cdot c_k(D_2;\, C_2^*),$$

so the non-separable problem reduces to a separable one. The proof uses Potter bounds for regularly varying functions: for any $\varepsilon > 0$ and large $|x|$, $(1-\varepsilon)|x|^{-\rho-\varepsilon} \leq G(|x|)/L(|x|)|x|^{-\rho} \leq (1+\varepsilon)|x|^{-\rho+\varepsilon}$.

Step 5. Cumulant convergence (non-CLT case). When both $C_1 \notin L^2$ and $C_2 \notin L^1$, the coefficients $c_k(t)$ converge to the cumulants of a 2-domain Rosenblatt distribution. This is computed explicitly by substituting the power-law tails and evaluating the resulting "covariogram" integrals on growing domains using asymptotic formulae for regularly varying functions (Appendix A).

Step 6. Moment determinacy. Random variables in the second Wiener chaos are *moment-determined* (their distributions are characterised by their cumulant sequences). Therefore, cumulant convergence $\Rightarrow$ convergence in distribution, completing the proof.

Step 7. Variance asymptotics. The variance $\mathrm{Var}(Y(t)) = \kappa_2(Y(t))$ is computed by evaluating $\int C^2$ over the growing product domain, splitting via Gneiting structure, and applying regularly varying integral asymptotics to each block.

Summary

This paper establishes the first complete CLT/non-CLT phase diagram for nonlinear functionals of Gaussian fields with *genuinely non-separable* (Gneiting-type) covariance, under anisotropic domain growth. The key insight is that Gneiting covariances are **asymptotically separable**: at large scales the non-separable coupling washes out, and the limiting distribution is governed by the same phase diagram as a separable model. Depending on the long-range dependence parameters (ρ_1, ρ_2), the limit is either Gaussian (CLT) or a 2-domain Rosenblatt distribution (non-Gaussian, second Wiener chaos).