

# Theory-mode report: A transition in the hole probability at finite temperature for free fermions in $d$ dimensions

Generated from Scholar Inbox digest

Digest date: 2026-05-08

## Paper information

- **Title:** A transition in the hole probability at finite temperature for free fermions in  $d$  dimensions
- **Authors:** Giuseppe Del Vecchio Del Vecchio, Pierre Le Doussal, Grégory Schehr
- **arXiv:** 2605.05431
- **Scholar digest score:** 94.0

## Executive summary

This paper analyzes the probability  $P(R, T)$  that a spherical region of radius  $R$  in a homogeneous free Fermi gas is completely empty at finite temperature. The central question is how this *hole probability* crosses over from classical Poisson-like behavior at high temperature to strongly correlated Pauli-driven behavior at low temperature. The main result is a sharp low-temperature large-radius scaling law

$$P(R, T) \sim \exp \left( - (k_F R)^{d+1} \Phi_d(u) \right), \quad u = \frac{2RT}{k_F} = \frac{2}{\pi} \frac{R}{\xi_T},$$

where  $k_F$  is the Fermi momentum and  $\xi_T$  the thermal correlation length. The striking point is that the scaling function  $\Phi_d(u)$  has a genuine nonanalytic transition at the universal critical value

$$u_c = \frac{2}{\pi},$$

of order  $\frac{3}{2}(d+1)$ . In the Coulomb-gas reformulation, this transition corresponds to the opening of a macroscopic gap in the optimal momentum-space density that realizes the rare event.

So this is not just a crossover story. In the joint limit  $k_F R \rightarrow \infty$ ,  $T/T_F \rightarrow 0$  with  $u$  fixed, the rare-fluctuation mechanism itself changes phase.

## Physical setting and observable

The model is the grand-canonical free Fermi gas in  $d$  dimensions at temperature  $T = 1/\beta$  and chemical potential  $\tilde{\mu}$ , with fixed average density  $\rho$ . The one-body occupation factor is

$$\sigma(k) = \frac{1}{e^{\beta(k^2/2 - \tilde{\mu})} + 1},$$

and the free fermions form a determinantal point process. The observable is the hole probability

$$P(R, T) = \mathbb{P}(\mathcal{N}_R = 0) = \lim_{s \rightarrow \infty} \langle e^{-s\mathcal{N}_R} \rangle,$$

where  $\mathcal{N}_R$  counts particles in the ball of radius  $R$ .

At very high temperature one expects classical independent-particle behavior, so for a Poisson process one has

$$P^{\text{Poi}}(R) = e^{-\rho V_d R^d}.$$

At zero temperature, the Pauli principle makes emptiness much rarer, and the known asymptotic is

$$P(R, 0) \sim e^{-\kappa_d (k_F R)^{d+1}}, \quad \kappa_d = \frac{2}{(d+1)^2 \Gamma(d+1)}.$$

The paper asks: what happens between these two regimes, and can the crossover be described sharply in any dimension?

## Main asymptotic regimes

The paper identifies two different large- $R$  regimes.

### 1. High-temperature / classical regime

For fixed  $T \sim T_F$  and  $R/\lambda_T \gg 1$ , with  $\lambda_T$  the thermal de Broglie wavelength,

$$\log P(R, T) \simeq -(R\sqrt{T})^d A_d(T/T_F) + (R\sqrt{T})^{d-1} B_d(T/T_F).$$

The functions  $A_d$  and  $B_d$  are given explicitly in terms of polylogarithms:

$$A_d(t) = -\frac{1}{2^{d/2} \Gamma(1 + d/2)} \text{Li}_{d/2+1}(-e^{\beta \tilde{\mu}}),$$

$$B_d(t) = \frac{2^{(d-3)/2}}{\pi \Gamma(d)} \int_0^\infty dv v^{(d-1)/2} [\text{Li}_{1/2}(-e^{\beta \tilde{\mu}} - v)]^2.$$

In the extreme high-temperature limit  $T/T_F \gg 1$ ,  $A_d$  reduces so that the Poisson law is recovered. So the leading term is volume-like and classical, exactly as expected.

### 2. Low-temperature / quantum large-deviation regime

For  $T \ll T_F$  and  $k_F R \gg 1$  with

$$u = \frac{2RT}{k_F} = \frac{2}{\pi} \frac{R}{\xi_T}$$

held fixed, the hole probability obeys

$$\log P(R, T) \simeq -(k_F R)^{d+1} \Phi_d(u).$$

This is the main scaling regime. The authors compute  $\Phi_d(u)$  exactly in any dimension and show that it has two branches,

$$\Phi_d(u) = \begin{cases} \Phi_{d,+}(u), & u > 2/\pi, \\ \Phi_{d,-}(u), & u < 2/\pi. \end{cases}$$

The two branches meet nonanalytically at  $u_c = 2/\pi$ , with

$$\Phi_{d,+}(u) - \Phi_{d,-}(u) \propto (u_c - u)^{\frac{3}{2}(d+1)} \quad (u \uparrow u_c).$$

Hence the transition is of order  $\frac{3}{2}(d+1)$ .

## Explicit scaling functions

For  $u > 2/\pi$ , the function  $\Phi_{d,+}(u)$  is a polynomial in  $1/u$  of degree  $d + 1$ . The paper gives explicit formulas, e.g.

$$\begin{aligned}\Phi_{1,+}(u) &= \frac{4}{3\pi u} - \frac{1}{\pi^2 u^2}, \\ \Phi_{2,+}(u) &= \frac{1}{8u} - \frac{8}{15\pi^2 u^2} + \frac{1}{12\pi^2 u^3}, \\ \Phi_{3,+}(u) &= \frac{4}{45\pi u} - \frac{1}{6\pi^2 u^2} + \frac{16}{105\pi^3 u^3} - \frac{1}{18\pi^4 u^4}.\end{aligned}$$

These large- $u$  forms match the asymptotics from the high-temperature/Widom regime, so the two asymptotic theories glue consistently.

For  $u < 2/\pi$ , the zero-temperature-like branch depends on parity of  $d$ . Examples given in the paper are

$$\Phi_{1,-}(u) = \frac{1}{2} - \frac{\pi^2 u^2}{48}, \quad \Phi_{3,-}(u) = \frac{1}{48} - \frac{\pi^2 u^2}{1440} + \frac{\pi^4 u^4}{53760},$$

and in  $d = 2$ ,

$$\Phi_{2,-}(u) = \sqrt{4 - \pi^2 u^2} \left( -\frac{\pi^2 u^2}{1440} + \frac{9}{40\pi^2 u^2} + \frac{7}{360} \right) - \frac{8}{15\pi^2 u^2} + \left( \frac{1}{6\pi^3 u^3} + \frac{1}{4\pi u} \right) \sin^{-1} \left( \frac{\pi u}{2} \right).$$

At  $u = 0$ , one recovers exactly the known zero-temperature coefficient  $\kappa_d$ , as one should.

## What the transition means

The really nice part is the mechanism. The hole probability is mapped to an effective Coulomb gas in momentum space. In that picture, the rare event “a large real-space hole is empty” is realized by an optimal momentum density  $\rho^*(p)$  minimizing an energy functional. The support of that optimizer changes qualitatively at  $u_c = 2/\pi$ .

- For  $u > u_c$ , the optimal density reaches down to zero momentum: no macroscopic momentum gap.
- For  $u < u_c$ , the support is shifted away from the origin: a genuine momentum gap opens.

So the nonanalyticity of  $\Phi_d(u)$  is not an accident of formulas; it reflects a change in the constrained optimal configuration of the Coulomb gas. The paper interprets this as a sharp change in the mechanism of rare fluctuations.

A good intuition is: when thermal effects are still strong enough compared with the region size ( $u > u_c$ ), the optimal rare event uses a “soft” reorganization of occupations. Below the threshold ( $u < u_c$ ), emptiness is so demanding that the system creates a hard depletion window in momentum space.

## Determinantal and angular decomposition

The exact starting point is the Fredholm determinant formula

$$P(R, T) = \det(I - \Pi_R K),$$

where  $K$  is the finite-temperature fermionic kernel. In dimensions  $d > 1$ , the authors use angular decomposition to factor the problem into independent radial sectors labeled by angular momentum  $\ell$ :

$$P(R, T) = \prod_{\ell \geq 0} (P^{(\ell)}(R, T))^{g_d(\ell)},$$

where  $g_d(\ell)$  is the degeneracy and each sector is expressed as a Fredholm determinant involving a finite-temperature Bessel kernel,

$$P^{(\ell)}(R, T) = \det(I - \Pi_{[0, R]} \hat{K}_\nu), \quad \nu = \ell + \frac{d}{2} - 1.$$

This reduction is important because it turns the  $d$ -dimensional problem into a family of effectively one-dimensional problems, which can then be attacked by random-matrix/Coulomb-gas tools.

## Coulomb-gas reformulation

A duality relation rewrites each sector's Fredholm determinant as a linear statistic for the zero-temperature Bessel determinantal process. From there, the event is represented by minimizing a Coulomb-gas energy

$$E_\nu[\rho] = -\frac{1}{2} \iint \log |p - p'| \hat{\rho}(p) \hat{\rho}(p') dp dp' + \frac{1}{2} \int \rho(p) \log(1 + e^{-\beta(p/(2R^2) - \tilde{\mu})}) dp,$$

with positivity constraints and a background equilibrium density. In the large- $R$  limit,

$$P^{(\ell)}(R, T) \sim e^{-2E_{0,\nu}}, \quad E_{0,\nu} = \min_{\rho \geq 0} E_\nu[\rho].$$

The minimizer develops the support transition mentioned above.

For fixed angular sector  $\ell = O(1)$ , the paper already finds a transition at  $u_c = 2/\pi$ , with an asymptotic of the form

$$\log P^{(\ell)}(R, T) \simeq -z^2 \begin{cases} \frac{2}{3\pi u} - \frac{1}{2\pi^2 u^2}, & u > 2/\pi, \\ \frac{1}{4} - \frac{\pi^2 u^2}{96}, & u < 2/\pi, \end{cases} \quad z = k_F R.$$

But for the full  $d$ -dimensional hole probability, the dominant contribution comes from large angular momentum sectors  $\ell \sim R$ , not finite  $\ell$ . The authors therefore solve the Coulomb-gas problem in the scaling  $\ell \sim \nu \sim \lambda z$  and integrate over  $\lambda$ .

That yields the exact representation

$$-\log P(R, T) \simeq \frac{2}{\Gamma(d-1)} z^{d+1} \int_0^1 d\lambda \lambda^{d-2} \phi(\lambda, \tilde{a}(u, \lambda)),$$

which becomes the explicit scaling function  $\Phi_d(u)$  after a change of variables. The lower limit of the integral is controlled by the critical support edge  $\tilde{a}_c(u)$ , and this is exactly where the transition enters.

## Critical region

The paper also identifies a critical scaling window around the transition. If

$$y = (2k_F R)^{2/3} \frac{u - u_c}{u_c},$$

then in the regime  $y = O(1)$ ,

$$P(R, T) \simeq G_d(y) e^{-(k_F R)^{d+1} \Phi_{d,+}(u)}.$$

The scaling function  $G_d(y)$  tends to 1 for  $y \rightarrow +\infty$ , while for  $y \rightarrow -\infty$  it reproduces the singular matching needed for the two branches. The temperature scale of the critical window is

$$\delta T \sim T_c (k_F R)^{-2/3} \sim T_F (k_F R)^{-5/3}, \quad T_c = \frac{k_F}{\pi R}.$$

So the transition sharpens with increasing  $R$ , and the critical region becomes parametrically narrow.

## Why the result matters

There are several reasons this is a strong paper.

- It gives an exact large-deviation scaling function for a nontrivial finite-temperature fermionic observable in any dimension.
- It identifies a universal critical value  $u_c = 2/\pi$  and a clear order of nonanalyticity.
- It ties together free-fermion determinantal structure, Widom asymptotics, Bessel/sine kernels, and Coulomb-gas variational methods.
- It is experimentally relevant: the hole probability is directly measurable in quantum gas microscope setups.

Conceptually, the nicest point is that even for *free* fermions, a sufficiently sharp rare-event question reveals a bona fide phase transition in the optimal fluctuation problem.

## Numerics and checks

The authors back the formulas with precise numerical evaluations of the relevant Fredholm determinants using Bornemann-type methods and finite- $z$  extrapolation. They also recover known rigorous results in  $d = 1$  and known  $T = 0$  formulas in any  $d$ , which makes the whole asymptotic picture feel very solid.

## For our group

- The most reusable structural idea is the **rare-event-as-Coulomb-gas-optimum** viewpoint: the nonanalyticity comes from a change in support of the optimizer, not from any interaction in the underlying fermion Hamiltonian.

- The scaling variable

$$u = \frac{2RT}{k_F} = \frac{2}{\pi} \frac{R}{\xi_T}$$

is the right knob: it compares the hole size directly with the thermal correlation length.

- For asymptotic matching problems, this paper is a nice example of combining **Widom/high- $T$  asymptotics** with a **low- $T$  Coulomb-gas large-deviation theory** and showing they meet consistently.

## Bottom line

A very elegant paper. It turns a natural full-counting-statistics observable for free fermions into an exactly analyzable variational problem and finds a universal finite-temperature transition in the hole-probability scaling function. If you're interested in determinantal processes, Coulomb gases, quantum large deviations, or the geometry of rare events, this one is absolutely worth reading.