Theory-mode report
Many-body dynamical localization in Fock space

Authors. Nathan Dupont, Bruno Peaudecerf, David Guéry-Odelin, Gabriel Lemarié, Bertrand Georgeot, Christian Miniatura, Nathan Goldman
arXiv. arXiv:2604.09224 **Date.** 2026-04-14

1. Paper information and one-paragraph orientation

This paper studies a periodically driven interacting bosonic two-mode system and argues that it realizes an *Anderson-like dynamical localization in Fock space*. The model is a kicked two-mode Bose system, equivalently a quantum kicked top with spin length $N/2$, where N is the conserved boson number. The main message is that although the classical mean-field dynamics becomes globally chaotic and diffusive on the Bloch sphere, the finite-N quantum dynamics can remain sharply localized in the basis of population-imbalance states. The localization is controlled by the kick angle a and the effective Planck constant $\hbar_{\text{eff}} = 2/N$, and it disappears in the classical limit.

The paper is short, but the conceptual structure is clean: (i) derive the classical kicked-top map and show bounded diffusion in the population imbalance; (ii) rewrite the Floquet problem as a finite Anderson-like tight-binding problem in Fock space; (iii) estimate the localization length from early-time diffusion; (iv) verify the ergodic-to-localized crossover through the long-time profile and quasienergy statistics; and (v) reinterpret discrete time-crystal behavior near $a \approx \pi$ as a consequence of localization of Floquet eigenstates near the poles.

2. Executive summary

The system consists of N interacting bosons in two modes with periodic delta-kicks in the inter-mode tunneling. Using Schwinger bosons, the Hamiltonian becomes a kicked collective spin model,

$$\hat{H}(t) = U\hat{J}_z^2 - k\hat{J}_x \sum_{m\in\mathbb{Z}} \delta(t - mT) + \text{const.}$$

with Floquet operator

$$\hat{U}_F = e^{-i\hbar_{\text{eff}} b\hat{J}_z^2} e^{-ia\hat{J}_x}, \qquad a = -k/\hbar, \quad b = \frac{NUT}{2\hbar}, \quad \hbar_{\text{eff}} = \frac{2}{N}.$$

The basis $|n_1, n_2\rangle$ of fixed occupation numbers acts as a one-dimensional Fock-space lattice of size $L = N+1$, indexed by the population imbalance $n_1 - N/2$. In the classical limit, trajectories spread diffusively over the Bloch sphere and eventually fill it ergodically. In the quantum problem, however, interference suppresses this transport, producing exponential localization in the Fock basis,

$$|\psi(n_1)|^2 \sim e^{-|n_1 - N/2|/\xi}.$$

The paper's central estimate is

$$\xi \approx \frac{D}{\hbar_{\text{eff}}^2} = \frac{\sin^2 a}{4\hbar_{\text{eff}}^2},$$

where $D = \sin^2(a)/4$ is the short-time classical diffusion constant. Thus localization is strongest for small kick angle $|\sin a| \ll 1$, and the finite system looks localized whenever $\xi \ll L$. This gives the crossover condition

$$\xi \ll L \quad \Longleftrightarrow \quad |\sin a| \ll \frac{4}{\sqrt{N}}.$$

Equivalent evidence comes from spectral statistics: the quasienergy-spacing law crosses over from GOE-like level repulsion in the ergodic regime to Poisson statistics in the localized regime. Near $a \approx \pi$, the same localization mechanism stabilizes a discrete time crystal by keeping the relevant Floquet eigenstates exponentially pinned near the fully polarized Fock states.

3. Problem setup and intuition

Degrees of freedom. The microscopic system is a driven two-mode bosonic model. If $\hat{a}_1, \hat{a}_2$ annihilate bosons in the two modes, then the interaction term is onsite and the kick periodically couples the modes. Because $\hat{N} = \hat{n}_1 + \hat{n}_2$ is conserved, the Hilbert space at fixed N has dimension $N + 1$.

Spin representation. Introduce collective spin operators

$$\hat{J}_x = \frac{\hat{a}_1^\dagger \hat{a}_2 + \hat{a}_2^\dagger \hat{a}_1}{2}, \qquad \hat{J}_y = \frac{\hat{a}_1^\dagger \hat{a}_2 - \hat{a}_2^\dagger \hat{a}_1}{2i}, \qquad \hat{J}_z = \frac{\hat{n}_1 - \hat{n}_2}{2}.$$

Then fixed total particle number means a fixed spin length $J = N/2$. The occupation basis $|n_1, n_2\rangle$ is exactly the $\hat{J}_z$ eigenbasis. So the Fock-space question is literally a transport problem along the discrete J_z axis.

Classical intuition. In the mean-field limit $N \to \infty$, the rescaled spin becomes a point on the Bloch sphere. One kick period applies a rotation around x and a nonlinear torsion around z. For sufficiently large torsion parameter b, the stroboscopic dynamics is globally chaotic. Starting near the equator ($z_0 = 0$), the classical variance of z first grows diffusively and then saturates to the uniform-sphere value $1/3$.

Quantum intuition. Finite-N quantization prevents indefinite diffusion. The Floquet eigenvalue problem can be rewritten as an effective one-dimensional tight-binding model with quasi-random onsite term

$$W_{j,n} = \tan\left(\frac{\varepsilon_j}{2} - \frac{b(n - N/2)^2}{N}\right),$$

and rapidly decaying hoppings $t_{n',n}$. That is the Anderson picture: classically the system wanders through phase space, but quantum mechanically the quasi-random phases generated by the nonlinear torsion act like disorder in Fock space and localize the wave function.

4. Main result, stated informally but precisely

The main result is not a single formal theorem, but the paper makes a precise scaling claim:

> For the kicked two-mode bosonic system (equivalently the quantum kicked top), in the regime of globally chaotic classical motion and small kick angle $|\sin a| \ll 1$, the quantum evolution from a balanced Fock state localizes exponentially in Fock space with localization length
>
> $$\xi \approx \frac{\sin^2 a}{4\hbar_{\text{eff}}^2} = \frac{N^2 \sin^2 a}{16}.$$
>
> Hence the dynamics is localized when this length is much smaller than the Fock-space size $L = N + 1$, i.e.
>
> $$|\sin a| \ll \frac{4}{\sqrt{N}},$$
>
> while the system becomes ergodic-like when ξ exceeds the system size.

A complementary formulation is in terms of observables. Let

$$\sigma_z^2(m) = \hbar_{\text{eff}}^2(\langle \hat{J}_z^2 \rangle - \langle \hat{J}_z \rangle^2)$$

for the quantum dynamics after m kicks. Classically,

$$\sigma_z^2(m) \approx \frac{1}{3}(1 - e^{-m/\tau}), \qquad \tau = \frac{2}{3\sin^2 a},$$

so early growth is diffusive with $D = \sin^2(a)/4$. Quantum mechanically, in the localized regime, $\sigma_z^2(m)$ saturates far below 1/3, and the long-time Fock profile is exponentially localized around the initial imbalance.

5. Proof architecture / derivation logic

Step 1: map the bosons to a kicked top. The Schwinger-boson map turns the two-mode interacting Bose system into a collective spin with nonlinear twist $\hat{J}_z^2$ and periodic rotation $\hat{J}_x$. This is the standard kicked-top structure, but here its Fock-space meaning is essential.

Step 2: analyze the classical limit. Set $\hbar_{\text{eff}} = 2/N$ and take $N \to \infty$ with b fixed. The spin becomes a classical unit vector (x, y, z) on the Bloch sphere. The resulting stroboscopic map is rotation plus torsion. In the chaotic regime the variance along z obeys

$$\sigma_z^2(m) \approx \frac{1}{3}\Big(1 - (1 - 1/\tau)^m\Big),$$

which for large τ yields the short-time diffusive law $\sigma_z^2(m) \approx 2Dm$.

Step 3: turn the Floquet eigenproblem into an Anderson-like lattice problem. In the J_z basis, the eigenvalue equation $\hat{U}_F|\phi_j\rangle = e^{-i\varepsilon_j T/\hbar}|\phi_j\rangle$ can be recast as an effective finite-range one-dimensional Hamiltonian on sites $n = 0, \ldots, N$. The diagonal terms depend quasi-randomly on n through the quadratic phase generated by $\hat{J}_z^2$. Rapidly decaying off-diagonal couplings make this look like a finite Anderson model rather than a fully connected random matrix.

Step 4: estimate the localization length from the break time. This is the key physics estimate. If the classical diffusion constant is D, then a wave packet initially diffuses as if classical until interference builds up. The localization time is taken to scale like the number of sites effectively occupied, $t_{\text{loc}}/T \sim \xi$. Matching the diffusive spread reached by that time,

$$\sigma_z^2 \sim 2D\xi,$$

with the variance of an exponentially localized state,

$$\sigma_z^2 \sim 2\hbar_{\text{eff}}^2\xi^2,$$

gives

$$\xi \sim \frac{D}{\hbar_{\text{eff}}^2}.$$

Substituting $D = \sin^2(a)/4$ yields the announced scaling.

Step 5: compare ξ to the system size. Because the Fock lattice has only $L = N+1$ sites, localization is physically visible only when $\xi \ll L$. This gives the crossover line $|\sin a| \ll 4/\sqrt{N}$. If instead $\xi \gtrsim L$, the state fills the available Fock space and the dynamics looks ergodic.

Step 6: verify using long-time profiles and spectral statistics. Numerically, the long-time probability distribution in the Fock basis is exponential in the localized regime and approximately uniform in the ergodic regime. Independently, quasienergy spacings move from GOE-like level repulsion to Poisson-like statistics. This is exactly the expected Anderson/ergodic dichotomy.

Step 7: connect to discrete time crystals. Near $a = \pi$, ideal π pulses pair Fock states across the two poles. When a is perturbed away from π, localization keeps the relevant Floquet states pinned near the poles, so the quasienergy splitting away from $h/(2T)$ becomes exponentially small in N/ξ. That produces robust subharmonic oscillations for exponentially long times.

6. Why it matters, limitations, and takeaway

Why it matters. The paper is a nice bridge between three literatures that often stay separate: kicked-top quantum chaos, Anderson-style localization, and many-body / Floquet time-crystal physics. The conceptual move is to treat the Fock basis itself as the physical lattice. That makes localization visible in a minimal interacting model with only two modes, which is experimentally much simpler than a disordered chain. It also gives a controlled finite-size setting where one can study the ergodic-to-localized crossover and potentially higher-dimensional Fock-space analogs for more modes.

What is genuinely strong.

- The scaling argument for ξ is transparent and physically motivated.
- The classical/quantum contrast is very sharp: classical chaos coexists with quantum localization.
- The GOE-to-Poisson crossover provides an independent spectral signature.
- The discrete-time-crystal reinterpretation is conceptually elegant rather than ornamental.

Limitations and caveats.

- This is a finite-dimensional Fock-space localization problem, not a proof of a thermodynamic many-body-localized phase.
- The key estimate is scaling-level and heuristic; it is not a rigorous theorem with error bounds.
- Localization is strongest for small kick amplitude, where the interaction term dominates. So the "disorder" is effective and dynamical, not externally tunable static disorder.
- The cleanest spectral analysis uses a random-phase kicked-top variant to avoid resonances; that improves diagnosis but slightly shifts the model away from the literal experimental protocol.
- Because the system is only two-mode, its relation to conventional spatial MBL is conceptual rather than direct.

Takeaway. The paper's core claim is that a driven, interacting, yet very small bosonic system can show a bona fide Anderson-like suppression of transport when viewed in the right basis: the many-body Fock basis. The controlling parameter is the competition between classical diffusion and quantum interference, summarized by the localization length $\xi \sim D/\hbar_{\text{eff}}^2$. In plain terms: classical chaos tries to spread the state across occupations, but quantum phases stop it after a finite Fock-space distance. That same mechanism then stabilizes long-lived subharmonic oscillations near the π-pulse regime.

Generated automatically on 2026-04-14 from the latest Scholar Inbox digest; theory-mode summary based on arXiv abstract/HTML content for 2604.09224.