Reading Report:
Uniform Concentration for α-Subexponential Random Operators

Auto-generated report for arXiv:2603.09487
T. Diao, X. Hu, V.V. Ulyanov, H. Wang

March 12, 2026

1 Core Question

When a random matrix $A \in \mathbb{R}^{m\times n}$ acts on a bounded set $T \subset \mathbb{R}^n$, classical results show that A is a near-isometry on T provided its entries (or rows/columns) are *subgaussian*. This paper asks: **to what extent do near-isometric properties persist when the subgaussian assumption is relaxed to α-subexponential tails ($\alpha \in (0, 2]$)?**

2 Key Definitions and Setup

Definition 1 (α-subexponential random variable)**.** *A random variable ξ is called* α-subexponential *(with $\alpha > 0$) if*

$$\|\xi\|_{\psi_\alpha} \;:=\; \inf\Big\{t > 0 : \mathbb{E}\exp\Big(\frac{|\xi|}{t}\Big)^{\alpha} \le 2\Big\} \;<\; \infty.$$

When $\alpha = 2$ this recovers the subgaussian class; $\alpha = 1$ gives the classical subexponential class. Smaller α allows heavier (but still exponential-type) tails.

Definition 2 (Isotropic random vector)**.** *A random vector $X \in \mathbb{R}^n$ is* isotropic *if $\mathbb{E}[XX^\top] = I_n$.*

Definition 3 (Talagrand's γ_α-functional)**.** *Let (T, d) be a metric space. An* admissible sequence *of partitions of T is an increasing sequence $(\mathcal{A}_k)_{k\ge 0}$ with $|\mathcal{A}_0| = 1$ and $|\mathcal{A}_k| \le 2^{2^k}$ for $k \ge 1$. For $\alpha > 0$,*

$$\gamma_\alpha(T, d) \;:=\; \inf_{(\mathcal{A}_k)} \sup_{t\in T} \sum_{k\ge 0} 2^{k/\alpha}\, \Delta_d\big(\mathcal{A}_k(t)\big),$$

where $\mathcal{A}_k(t)$ is the element of $\mathcal{A}_k$ containing t and $\Delta_d(A) = \sup_{s,t\in A} d(s, t)$ is the diameter of A. When d is the Euclidean metric we write $\gamma_\alpha(T)$. The quantity $\mathrm{rad}(T) := \sup_{t\in T} \|t\|_2$.

3 A Concrete Example

Johnson–Lindenstrauss with heavy tails. Consider N points $x_1, \dots, x_N \in \mathbb{R}^n$ and let A be an $m \times n$ random matrix whose rows are i.i.d., isotropic, α-subexponential with ψ_α-norm $\le K$. Apply the row-wise theorem (Corollary 5) with $T = \{x_i - x_j\}$: for

$$m \;\gtrsim\; K^{8/\alpha}\, \frac{(\log N)^{2/\alpha}}{\varepsilon^2},$$

the rescaled map $\frac{1}{\sqrt{m}}A$ satisfies $(1-\varepsilon)\|x_i - x_j\|_2 \le \big\|\frac{1}{\sqrt{m}}A(x_i - x_j)\big\|_2 \le (1+\varepsilon)\|x_i - x_j\|_2$ for all pairs simultaneously, with high probability. When $\alpha = 2$ (subgaussian) this recovers the classical JL bound $m \gtrsim \varepsilon^{-2} \log N$.

4 Main Results

Theorem 4 (Row-wise model — Theorem 1.1 of the paper)**.** *Let $B \in \mathbb{R}^{\ell \times m}$ be fixed and $A \in \mathbb{R}^{m \times n}$ have independent, isotropic rows with ψ_α-norm $\leq K$, $\alpha \in (0, 2]$. For any bounded $T \subset \mathbb{R}^n$,*

$$\mathbb{E} \sup_{x \in T} \big| \|BAx\|_2 - \|B\|_{\mathrm{HS}} \|x\|_2 \big| \;\leq\; C(\alpha)\, K^{4/\alpha}\, \|B\|_{\mathrm{op}} \big(\gamma_\alpha(T) + \mathrm{rad}(T)\big).$$

Moreover, for every $u \geq 0$, with probability $\geq 1 - C\, e^{-u^\alpha}$,

$$\sup_{x \in T} \big| \|BAx\|_2 - \|B\|_{\mathrm{HS}} \|x\|_2 \big| \;\leq\; C(\alpha)\, K^{4/\alpha}\, \|B\|_{\mathrm{op}} \big(\gamma_\alpha(T) + u \cdot \mathrm{rad}(T)\big).$$

Corollary 5 (Identity pre-factor)**.** *Setting $B = I_m$ in Theorem 4:*

$$\sup_{x \in T} \big| \|Ax\|_2 - \sqrt{m}\, \|x\|_2 \big| \;\leq\; C(\alpha)\, K^{4/\alpha} \big(\gamma_\alpha(T) + u \cdot \mathrm{rad}(T)\big)$$

with probability $\geq 1 - Ce^{-u^\alpha}$.

Theorem 6 (Column-wise model — Theorem 1.2 of the paper)**.** *Let $A \in \mathbb{R}^{m \times n}$ have independent, mean-zero columns $A_i \in \mathbb{R}^m$ with $\|A_i\|_2 = 1$ a.s. and ψ_α-norm $\leq K$, $\alpha \in (0, 2]$. Then for bounded $T \subset \mathbb{R}^n$,*

$$\mathbb{E} \sup_{x \in T} \big| \|Ax\|_2 - \|x\|_2 \big| \;\leq\; C(\alpha)\, K \, \big(\gamma_\alpha(T) + \mathrm{rad}(T)\big),$$

and with probability $\geq 1 - Ce^{-u^\alpha}$,

$$\sup_{x \in T} \big| \|Ax\|_2 - \|x\|_2 \big| \;\leq\; C(\alpha)\, K \, \big(\gamma_\alpha(T) + u \cdot \mathrm{rad}(T)\big).$$

Remark 7 (Column normalization is necessary)**.** *The column-wise result* requires $\|A_i\|_2 = \lambda$ *a.s. A Bernoulli-scaled counterexample shows that isotropy alone is insufficient: if $b \sim$* Bernoulli$(1/2)$ *and $X \sim$* Unif$(\sqrt{m}\, S^{m-1})$*, then bX is isotropic and subgaussian, yet $\|Ae_1\|_2$ deviates from any target by $\geq \sqrt{m}/2$ with probability $1/2$.*

Summary of main results:

- **Row-wise model:** Near-isometry controlled by $\gamma_\alpha(T)$ with factor $K^{4/\alpha}$; recovers optimal subgaussian bounds at $\alpha = 2$.
- **Column-wise model:** Near-isometry controlled by $\gamma_\alpha(T)$ with linear factor K; requires column normalization.
- **Applications:** JL embeddings for α-subexponential matrices (target dimension $\sim K^{8/\alpha}(\log N)^{2/\alpha}/\varepsilon^2$); restricted isometry property for compressed sensing; column-normalized structured matrices.
- **Methodological novelty:** A direct decomposition approach that avoids subgaussian-specific tools, yielding a more transparent proof even at $\alpha = 2$.

5 Proof Sketch

The proofs of Theorems 4 and 6 share a common strategy but differ in the technical ingredients at the single-vector level.

Step 1: Reduce to a chaining problem

Define the random process $Z_x := \|BAx\|_2 - \|B\|_{\mathrm{HS}}\|x\|_2$ (row-wise) or $Z_x := \|Ax\|_2 - \|x\|_2$ (column-wise). The goal is to bound

$$\sup_{x\in T} |Z_x| \quad \text{or equivalently} \quad \sup_{x,y\in T} |Z_x - Z_y|.$$

By Talagrand's generic chaining theorem (Theorem 2.1 in the paper), it suffices to show that the process has *α-subexponential increments*:

$$\|Z_x - Z_y\|_{\psi_\alpha} \;\le\; M\,\|x-y\|_2,$$

since then $\mathbb{E}\sup|Z_x - Z_y| \;\le\; C(\alpha)\,M\,\gamma_\alpha(T)$ and the exponential tail follows with an extra $u\cdot \mathrm{rad}(T)$ term.

Step 2 (Row-wise): Establish α-subexponential increments via Hanson–Wright

For the row-wise model (Lemma 3.1 in the paper), the key input is an *α-subexponential Hanson–Wright inequality* due to Sambale: for centered α-subexponential $X = (X_1,\dots,X_n)$ with $\|X_i\|_{\psi_\alpha} \le K$ and a symmetric matrix M,

$$\mathbb{P}\big(|X^\top MX - \mathbb{E}[X^\top MX]| \ge t\big) \;\le\; 2\exp\Big(-C(\alpha)\min\Big\{\frac{t^2}{K^4\|M\|_{\mathrm{HS}}^2},\; \Big(\frac{t}{K^2\|M\|_{\mathrm{op}}}\Big)^{\alpha/2}\Big\}\Big).$$

To bound $|Z_x - Z_y|/\|x-y\|_2$, split into two regimes:

1. **Large deviation ($s \ge 2\|B\|_{\mathrm{HS}}$):** Use the triangle inequality to reduce to a single-vector concentration $\mathbb{P}(\|BAu\|_2 \ge s)$, then apply the ψ_α norm bound from Sambale's Theorem 3.2.

2. **Moderate deviation ($1 < s < 2\|B\|_{\mathrm{HS}}$):** Polarize $\|BAx\|_2^2 - \|BAy\|_2^2 = \langle BAu, BAv\rangle$ with $u = (x-y)/\|x-y\|_2$ and $v = x+y$. Expand via $2\langle w_1, w_2\rangle = \|w_1+w_2\|^2 - \|w_1\|^2 - \|w_2\|^2$ to reduce to three centered quadratic forms $Y_w = \|BX_w\|_2^2 - \mathbb{E}\|BX_w\|_2^2$, each controlled by the Hanson–Wright inequality.

Combining both regimes gives $\|Z_x - Z_y\|_{\psi_\alpha} \le C(\alpha)\,K^{4/\alpha}\|B\|_{\mathrm{op}}\|x-y\|_2$.

Step 3 (Column-wise): A direct decomposition avoiding subgaussian tools

For the column-wise model, the proof uses a fundamentally different approach from the original Plan–Vershynin argument (which relies on subgaussian-specific moment bounds).

Write $Ax = \sum_{i=1}^n x_iA_i$ (sum of rescaled columns). Because columns have unit norm, $\|Ax\|_2^2 = \sum_{i,j} x_ix_j\langle A_i, A_j\rangle$, and the diagonal contributes $\|x\|_2^2$ deterministically. The off-diagonal terms $x_ix_j\langle A_i, A_j\rangle$ ($i \ne j$) are mean-zero (by independence and zero-mean columns) and have α-subexponential tails.

The increment $Z_x - Z_y$ is then controlled by bounding the ψ_α-norm of the off-diagonal bilinear form. Independence of the columns, combined with basic ψ_α algebra (Hölder-type inequalities for Orlicz norms, Proposition 2.1), yields

$$\|Z_x - Z_y\|_{\psi_\alpha} \;\le\; C(\alpha)\,K\,\|x-y\|_2.$$

The absence of $K^{4/\alpha}$ (compared to the row-wise bound) reflects the column-normalization constraint $\|A_i\|_2 = 1$, which eliminates the main source of variance inflation.

Step 4: Apply generic chaining (Theorem 2.2)

With the increment bound M established in Steps 2 or 3, invoke Theorem 2.2:

- **Expectation bound:** Use L^{2n} moment growth $\|Z_x - Z_y\|_{2n} \leq CM \cdot 2^{n/\alpha} \|x-y\|_2$ to translate into the γ_α-functional via an admissible partition.
- **Tail bound:** Talagrand's inequality (2.2) adds the $u \cdot \mathrm{rad}(T)$ term; reparametrize u to convert the Gaussian-type e^{-u^2} tail into e^{-u^α}.

This completes the proofs of Theorems 4 and 6. □

References

[1] R. Adamczak et al., "Tail and moment estimates for sums of independent random vectors with logarithmically concave tails," *Studia Math.*, 2012.

[2] T. Diao, X. Hu, V.V. Ulyanov, H. Wang, "Uniform Concentration for α-Subexponential Random Operators," arXiv:2603.09487, 2026.

[3] J. Jeong et al., "Sub-Gaussian matrices on sets: optimal tail dependence and applications," 2020.

[4] Y. Plan, R. Vershynin, "Dimension reduction by random hyperplane tessellations," *Discrete Comput. Geom.*, 2014.

[5] H. Sambale, "Some notes on concentration for α-subexponential random variables," 2023.

[6] M. Talagrand, *Upper and Lower Bounds for Stochastic Processes*, Springer, 2014.