Report: Quantitative Equidistribution of Eigenvalues of Random Normal Matrices in the Wasserstein Distance

arXiv:2603.18313 by P. García Arias

Report generated: March 20, 2026

1 Core Question

How fast does the empirical spectral measure of a Random Normal Matrix (RNM) converge to its equilibrium measure, when "fast" is measured by the expected 2-Wasserstein distance $\mathbb{E}\, W_2$? More generally, what quantitative convergence rates can be obtained for determinantal and homogeneous point processes?

2 Main Statement

Theorem 1 (Theorem 2 of the paper)**.** *Let $Q : \mathbb{C} \to \mathbb{R}$ be a potential satisfying the regularity assumptions (A1)–(A4) below, and assume the droplet S is convex. If $z_1, \dots, z_N$ are the eigenvalues of the $N \times N$ Random Normal Matrix model with potential Q, and*

$$d\sigma(z) = \tfrac{1}{4}\Delta Q(z)\, \chi_S(z)\, dA(z)$$

is the equilibrium measure (where $dA(z) = \frac{1}{\pi} dx\, dy$), then

$$\mathbb{E}\, W_2\Big(\frac{1}{N}\sum_{n=1}^{N} \delta_{z_n},\ \sigma\Big) \ \lesssim\ \frac{\sqrt{\log N}}{\sqrt{N}}.$$

The regularity assumptions on Q are:
(A1) Q is real-analytic in a neighborhood of S.
(A2) $\Delta Q \neq 0$ on S.
(A3) ∂S is a C^ω smooth Jordan curve.
(A4) The pre-droplet equals the droplet: $S^* = S$.

For homogeneous point processes the paper also proves:

Theorem 2 (Theorem 3)**.** *Let Ξ be a homogeneous point process in $\mathbb{R}^d$ with the variance bound $\mathrm{Var}\big(\sum_{x\in\Xi^L} f(x)\big) \lesssim L\|f\|_{L^2}^2$, and let $\Omega \subset \mathbb{R}^d$ be compact and convex. Then*

$$\mathbb{E}\, W_2\Big(\frac{1}{N}\sum_{x\in\Xi^L\cap\Omega} \delta_x,\ \frac{1}{|\Omega|}dx\big|_\Omega\Big) \ \lesssim\ \begin{cases} \dfrac{\sqrt{\log L}}{\sqrt{L}} & d = 2, \\ \dfrac{1}{L^{1/d}} & d > 2. \end{cases}$$

This applies to the infinite Ginibre ensemble, the Bessel ensemble, and zeros of the planar Gaussian Analytic Function (GAF).

3 Early Concrete Example

Example 1 (Ginibre ensemble)**.** Take $Q(z) = |z|^2$. The droplet is the closed unit disk $S = \overline{\mathbb{D}}$, and the equilibrium measure is $\sigma = \frac{1}{\pi}\chi_{\mathbb{D}}\, dx\, dy$ (uniform measure on the disk). The eigenvalues of a random $N \times N$ matrix with i.i.d. complex Gaussian entries, rescaled by $1/\sqrt{N}$, form this model. The theorem says

$$\mathbb{E}\, W_2\Big(\frac{1}{N}\sum_{n=1}^{N}\delta_{z_n},\ \sigma\Big) = O\Big(\frac{\sqrt{\log N}}{\sqrt{N}}\Big).$$

Previously, Prod'homme obtained the rate $O(1/\sqrt{N})$ for Ginibre using rotational symmetry. This paper recovers an almost-optimal rate *without* symmetry, for *all* RNM models with convex droplet.

4 Intuition and Setup

Key objects and notation

Definition 1 (Wasserstein distance)**.** For two probability measures μ, ν on a metric space (M, d), the p-Wasserstein distance is

$$W_p(\mu,\nu) = \Big(\inf_{\pi\in\Pi(\mu,\nu)}\int_{M\times M} d(x,y)^p\, d\pi(x,y)\Big)^{1/p},$$

where $\Pi(\mu,\nu)$ is the set of couplings (transport plans) with marginals μ and ν.

Definition 2 (Random Normal Matrix model)**.** Given a potential $Q : \mathbb{C} \to \mathbb{R}$ with sufficient growth at infinity, consider $N \times N$ complex normal matrices ($M^*M = MM^*$) with probability density proportional to $e^{-N \operatorname{tr} Q(M)}$. The eigenvalues $z_1, \dots, z_N \in \mathbb{C}$ form a *determinantal point process* (DPP) with joint density

$$\frac{1}{Z_N}\prod_{i<j}|z_i - z_j|^2 \prod_{j=1}^{N} e^{-NQ(z_j)}\, dA(z_j).$$

Definition 3 (Droplet and equilibrium measure)**.** The *equilibrium potential* is $\hat{Q}(z) = \sup\{f(z) : f \text{ subharmonic}, f \le Q, f(w) \le \log^+|w|^2 + O(1)\}$. The *droplet* $S = \operatorname{supp}(\frac{1}{4}\Delta\hat{Q})$ is the compact set where eigenvalues accumulate. The equilibrium measure is $d\sigma = \frac{1}{4}\Delta Q\, \chi_S\, dA$.

Definition 4 (Homogeneous point process)**.** A point process Ξ in $\mathbb{R}^d$ whose first intensity ρ_1 is constant. One rescales Ξ by $L^{-1/d}$ to get Ξ^L with first intensity $\rho_1^L = cL$, concentrating more points in any fixed region Ω.

Central idea

The standard approach to bounding W_2 is Peyre's inequality: $W_2(\mu,\nu) \le 2\|\mu - \nu\|_{\dot{H}^{-1}(d\mu)}$, which connects optimal transport to negative Sobolev norms. But empirical measures (sums of Dirac masses) are "rough" and hard to control in Sobolev norms directly.

The key innovation is a **heat smoothing procedure** on convex domains with Neumann boundary conditions. One replaces μ and ν by their heat-flow regularizations $\mu_t = P_t[\mu]$ and $\nu_t = P_t[\nu]$, then estimates:

$$W_2(\mu,\nu) \ \le\ W_2(\mu,\mu_t) + W_2(\mu_t,\nu_t) + W_2(\nu_t,\nu).$$

- The "smoothing error" $W_2(\mu,\mu_t)$ is controlled by $O(\sqrt{t})$ via the heat kernel.

- The "smooth comparison" $W_2(\mu_t, \nu_t)$ is controlled by Fourier coefficients (Neumann eigenfunctions) weighted by $e^{-\lambda_k t}/\lambda_k$, which are in turn controlled by variance estimates for linear statistics of the point process.
- Optimizing t yields the final rate.

Previous work (Borda, Marzo–Ortega-Cerdà) applied this on compact manifolds *without boundary*. This paper extends it to **compact convex subsets of** $\mathbb{R}^d$ using the Neumann Laplacian, which is the main technical novelty.

5 Main Results (Summary)

- **Heat Smoothing Inequality (Theorem 6):** For measures μ, ν on a convex domain $\Omega \subset \mathbb{R}^d$ with $\nu \geq c\,\mathrm{Vol}|_\Omega$,

$$W_2(\mu, \nu) \leq C_1(dt)^{1/2} + \frac{2}{C_2(\nu, t)} \Big(\sum_{k \geq 1} \frac{e^{-\lambda_k t}}{\lambda_k} |\hat{\mu}(k) - \hat{\nu}(k)|^2 \Big)^{1/2},$$

  where λ_k are Neumann eigenvalues of Ω, and $\hat{\mu}(k) = \int \varphi_k \, d\mu$ are Neumann–Fourier coefficients.

- **RNM (Theorem 2):** $\mathbb{E}\, W_2(\mu_N, \sigma) \lesssim \sqrt{\log N}/\sqrt{N}$ for convex droplet with regular potential.

- **Homogeneous processes (Theorem 3):** Rate $\sqrt{\log L}/\sqrt{L}$ in $d = 2$; rate $L^{-1/d}$ in $d > 2$, under a variance bound on linear statistics.

- **Corollaries:** Applies immediately to the infinite Ginibre ensemble, Bessel ensemble (via the reproducing property of their kernels), and the planar GAF zeros (via known variance bounds).

- **Optimality:** For Ginibre, the known lower bound is $\Omega(1/\sqrt{N})$, so the result is optimal up to a $\sqrt{\log N}$ factor.

6 Proof Sketch

Step 1: Heat smoothing on convex domains (Theorem 6)

1. Given measures μ, ν on Ω, define $\mu_t = P_t[\mu]$, $\nu_t = P_t[\nu]$ where P_t is the Neumann heat semigroup.
2. **Smoothing cost** (Lemma 7–8): Using the explicit transport plan $d\pi(x, y) = P(t, x, y)\, d\mu(y)\, dx$ and the convexity of Ω (to ensure $\partial_n |x - y|^2 \geq 0$), one shows

$$W_2(\mu, \mu_t) \leq \sqrt{2dt\, \mu(\Omega)}.$$

3. **Smooth comparison** (Lemma 9): Apply Peyre's inequality to the smoothed measures. Expanding in the Neumann eigenbasis $\{\varphi_k\}$:

$$W_2(\mu_t, \nu_t) \leq \frac{2}{C_2(\nu, 2t)} \Big(\sum_{k \geq 1} \frac{e^{-2\lambda_k t}}{\lambda_k} |\hat{\mu}(k) - \hat{\nu}(k)|^2 \Big)^{1/2}.$$

4. Combine via triangle inequality and optimize t.

Step 2: Application to homogeneous processes (Theorem 3)

1. Set $\mu_L = \frac{1}{N}\sum_{x\in\Xi^L\cap\Omega}\delta_x$ and $\nu = \frac{1}{|\Omega|}dx|_\Omega$.
2. The variance hypothesis gives $\mathbb{E}|\hat{\mu}_L(k) - \hat{\nu}(k)|^2 \lesssim 1/L$.
3. Input into Corollary 10 (a bookkeeping lemma using Weyl's law $|\{m : \lambda_m \le x\}| \lesssim x^{d/2}$ to sum the Fourier series). Optimize $t = L^{-\alpha}$ to get $\gamma = a/(2b+d)$ with $a = 1$, $b = 0$ giving $\gamma = 1/d$ (or $\sqrt{\log}$ correction at the critical exponent $b + d/2 = 1$, i.e. $d = 2$).
4. For DPPs with reproducing kernels, the variance bound follows from the identity $\mathrm{Var}(\sum f(x)) = \frac{1}{2}\iint |f(x) - f(y)|^2|K(x,y)|^2$ and $\int |K(x,y)|^2\,dy = K(x,x) \approx L$.

Step 3: Application to Random Normal Matrices (Theorem 2)

The RNM proof is more delicate because the equilibrium measure σ is *not* uniform—it has varying density $\frac{1}{4}\Delta Q$—and the kernel behavior changes near the boundary ∂S.

1. **Decompose** $\mathbb{C}$ into three regions relative to the droplet S:
   - *Exterior* ($|z| \gg$ or $d(z, \partial S) > a_0\delta_N$ outside S, where $\delta_N = \sqrt{\log N}/\sqrt{N}$): the kernel decays as $K(z,z) \lesssim (2 + |z|^2)^{-3}$, so very few points lie here. Bound: $\mathbb{E}\,W_2 \lesssim 1/\sqrt{N}$.
   - *Edge* (E_N: a strip of width $\sim \delta_N$ around ∂S): handle by a dyadic partition of E_N into 2^k sectors, constructing intermediate measures that redistribute mass sector-by-sector. Telscoping over $K \sim \log N$ levels gives $\mathbb{E}\,W_2 \lesssim \sqrt{\log N}/\sqrt{N}$.
   - *Bulk* ($S_N = r_N\mathring{S}$, slightly contracted droplet): here $K(z,z) \approx \frac{N}{4}\Delta Q(z)$. Apply the heat smoothing (Theorem 6). The key estimate is

     $$\mathbb{E}\Big|\frac{1}{N}\sum_{z_n\in S_N}\varphi_k(z_n) - \sigma(\varphi_k)\Big|^2 \lesssim \frac{1}{N},$$

     obtained from the DPP variance formula and the kernel approximation in the bulk.
2. **Triangle inequality:** Chain exterior $\to$ edge $\to$ bulk $\to$ smoothing, each contributing $O(\sqrt{\log N}/\sqrt{N})$.
3. The convexity of S is used crucially in the heat smoothing step (Neumann boundary, Lemma 7) and in the geometric decomposition.

References

García Arias, P. "Quantitative equidistribution of eigenvalues of Random Normal Matrices in the Wasserstein distance." arXiv:2603.18313, March 2026.