# Reading Report: Outlier Eigenvectors for Sparse Non-Hermitian Perturbations

Based on arXiv:2603.03972 by M. Galanis and M. Louvaris

Auto-generated report

March 9, 2026

## 1 Core Question

Given a sparse non-Hermitian random matrix $X_n$ (with sparsity parameter $K_n$) and a deterministic finite-rank perturbation $E_n$, how does the outlier eigenvector of $Y_n = X_n + E_n$ align with the spike eigenspace of $E_n$? Specifically, the paper extends the rank-one eigenvector overlap result of Hachem–Louvaris–Najim (2026) to arbitrary finite rank, proving that for a spike $\mu$ with $|\mu| > 1$, the squared projection of the outlier eigenvector onto the spike eigenspace converges in probability to $1 - |\mu|^{-2}$.

## 2 Exact Main Statement

**Theorem 1** (Theorem 2.6, simplified)**.** *Let $X_n \in \mathbb{C}^{n\times n}$ be a sparse i.i.d. non-Hermitian random matrix with sparsity parameter $K_n$ satisfying $\log^9 n/K_n \to 0$, and entries drawn from a sub-Gaussian law $\chi$ with $\mathbb{E}\chi = 0$, $\mathbb{E}|\chi|^2 = 1$. Let $E_n$ be a deterministic rank-$r$ perturbation admitting a biorthogonal decomposition (Assumption 3 below). Fix an outlier spike $\mu$ with $|\mu| > 1$ and let $\tilde{u}_{\ell,n}$ be a unit right eigenvector of $Y_n = X_n + E_n$ associated with the eigenvalue $\lambda_{\ell,n} \xrightarrow{P} \mu$. Let $F_n = \ker(\mu_n I - E_n)$ be the spike eigenspace. Then:*

*(a)* ***Overlap with the correct spike:***

$$\langle \tilde{u}_{\ell,n},\, F_n\rangle^2 \;:=\; \|Q^*_{\ell,n}\tilde{u}_{\ell,n}\|^2 \;\xrightarrow{P}\; 1 - \frac{1}{|\mu|^2}.$$

*(b)* ***Orthogonality to other spikes:*** *If $\mu' \neq \mu$ is another spike with associated eigenspace $F_{\ell',n} \perp F_n$, then*

$$\langle \tilde{u}_{\ell,n},\, F_{\ell',n}\rangle^2 \;\xrightarrow{P}\; 0.$$

Here $Q_{\ell,n} \in \mathbb{C}^{n\times k_\ell}$ has orthonormal columns spanning $F_n$, and $\langle x, F\rangle := \|\mathrm{Proj}_F\, x\|$ is the norm of the orthogonal projection of $x$ onto $F$.

## 3 A Concrete Example

**Rank-one case ($r = 1$).** Take $E_n = \mu\, u_n v_n^*$ where $u_n, v_n \in \mathbb{C}^n$ are unit vectors with $\langle v_n, u_n\rangle \to \mu$ (so $E_n$ has a single eigenvalue $\mu$ with $|\mu| > 1$). Theorem 2.6 recovers the earlier rank-one result (Theorem 1.6 of [1]):

$$\left|\left\langle \tilde{u}_n,\, \frac{u_n}{\|u_n\|}\right\rangle\right|^2 \;\xrightarrow{P}\; 1 - \frac{1}{|\mu|^2}.$$

In words: once $|\mu| > 1$, most of the outlier eigenvector "lives" in the spike direction; only a fraction $|\mu|^{-2}$ leaks into the bulk.

**Rank-two illustration.** Let $E_n$ have eigenvalues $\mu_1, \mu_2$ with $|\mu_1|, |\mu_2| > 1$ and orthogonal eigenspaces $F_1, F_2$. Then the outlier eigenvector near $\mu_j$ satisfies $\langle \tilde{u}_j, F_j \rangle^2 \to 1 - |\mu_j|^{-2}$ and $\langle \tilde{u}_j, F_k \rangle^2 \to 0$ for $k \neq j$.

# 4 Intuition and Setup

## 4.1 The random matrix model

Define the sparse i.i.d. matrix $X_n = (X_{ij}) \in \mathbb{C}^{n \times n}$ via

$$X_{ij} = \frac{1}{\sqrt{K_n}} B_{ij} A_{ij},$$

where $A_{ij} \overset{\text{i.i.d.}}{\sim} \chi$ (zero-mean, unit-variance, sub-Gaussian) and $B_{ij} \overset{\text{i.i.d.}}{\sim} \text{Bernoulli}(K_n/n)$, independent of $A$. The parameter $K_n$ controls sparsity: when $K_n = n$ one recovers the standard i.i.d. model; when $K_n \ll n$ the matrix is sparse.

The perturbation has the biorthogonal form

$$E_n = P_n \Lambda_n W_n^*, \qquad W_n^* P_n = I_r,$$

with $\Lambda_n = \text{diag}(\mu_{(1,n)}, \ldots, \mu_{(r,n)})$.

## 4.2 Why the extension is non-trivial

In the non-Hermitian setting, multiplicities and interactions between distinct spike blocks create structural difficulties absent in rank one. One must:

- Control the kernel of a finite-dimensional matrix-valued function derived from the resolvent.
- Localise the kernel vector onto the correct spike block.
- Handle the fact that left and right eigenvectors need not coincide (non-normality).

## 4.3 Resolvent philosophy

The entire argument is resolvent-based: express the outlier eigenvector via $(X_n - \lambda I)^{-1}$, reduce the problem to a finite-dimensional kernel, then use probabilistic estimates on bilinear forms of the resolvent.

# 5 Main Results (Bulleted)

- **Finite-rank kernel–eigenspace bijection (Lemma 3.1):** For $\lambda \notin \sigma(X_n)$, the map $\Phi(a) = (X_n - \lambda I)^{-1} U_n a$ is a linear bijection from $\ker(I_r + M_n(\lambda))$ to $\ker(Y_n - \lambda I)$, where $M_n(\lambda) = V_n^* (X_n - \lambda I)^{-1} U_n \in \mathbb{C}^{r \times r}$.
- **Kernel localisation (Lemma 3.3):** If $\|I_r + M_n(\lambda) - (I_r - \mu^{-1}\Lambda_n)\| \leq \varepsilon_n$ and $a_n \in \ker(I_r + M_n(\lambda))$, decompose $a_n = (a_\mu, a_{\neq})$; then $\|a_{\neq}\| \leq C\varepsilon_n \|a_\mu\|$.
- **Resolvent bilinear form asymptotics (Lemma 4.1):** For bounded deterministic vectors $u_n, v_n$ and $|z| > 1$: $\langle R_n(z) u_n, v_n \rangle + z^{-1} \langle u_n, v_n \rangle \xrightarrow{P} 0$.

- **Resolvent norm asymptotics (Lemma 4.4):** $\|R_n(z)w_n\|^2 - \|w_n\|^2/(|z|^2-1) \xrightarrow{P} 0$ for $|z| > 1$.
- **Main overlap formula (Theorem 2.6(a)):** $\langle \tilde{u}_{\ell,n}, F_n\rangle^2 \xrightarrow{P} 1 - |\mu|^{-2}$.
- **Cross-spike orthogonality (Theorem 2.6(b)):** $\langle \tilde{u}_{\ell,n}, F_{\ell',n}\rangle^2 \xrightarrow{P} 0$ for distinct spikes (assuming orthogonal eigenspaces).

# 6 Proof Sketch

**Step 1: Finite-rank reduction.** By Corollary 3.2, any unit outlier eigenvector $\tilde{u}_{\ell,n}$ of $Y_n$ can be written as

$$\tilde{u}_{\ell,n} = \frac{R_n(\lambda_{\ell,n})\, U_n\, a_n}{\|R_n(\lambda_{\ell,n})\, U_n\, a_n\|},$$

where $R_n(z) = (X_n - zI)^{-1}$ and $a_n \in \ker(I_r + M_n(\lambda_{\ell,n})) \setminus \{0\}$. This reduces the $n$-dimensional eigenvector problem to understanding the $r$-dimensional kernel vector $a_n$.

**Step 2: Kernel localisation.** Using Corollary 4.2 (bilinear form convergence) and Lemma 4.3 (resolvent continuity), one shows

$$\left\| M_n(\lambda_{\ell,n}) + \frac{1}{\mu}\Lambda_n \right\| \xrightarrow{P} 0.$$

By Lemma 3.3, this forces $a_n$ to concentrate on the $\mu$-block: writing $a_n = (a_{\mu,n},\, a_{\neq,n})$, one gets $\|a_{\neq,n}\| = o_P(1)\|a_{\mu,n}\|$. Define $\tilde{a}_{\mu,n} = (a_{\mu,n}, 0)$; then $U_n\tilde{a}_{\mu,n} \in F_n$ (the spike eigenspace).

**Step 3: Numerator: spike-block projection.** Set $\hat{w}_n = U_n\tilde{a}_{\mu,n}$ and $c_n = Q^*_{\ell,n}\hat{w}_n$ with $\|c_n\| = \|\hat{w}_n\|$. By the resolvent bilinear form estimate (Corollary 4.2),

$$Q^*_{\ell,n} R_n(\mu)\, Q_{\ell,n}\, c_n \;\approx\; -\frac{1}{\mu}\, c_n,$$

so $\|Q^*_{\ell,n} R_n(\mu)\, \hat{w}_n\|^2 \approx |\mu|^{-2}\|c_n\|^2$.

**Step 4: Denominator: resolvent norm.** By Corollary 4.5 (quadratic form estimate derived from Lemma 4.4),

$$\|R_n(\lambda_{\ell,n})\, U_n a_n\|^2 \;\approx\; \frac{\|U_n a_n\|^2}{|\mu|^2 - 1}.$$

**Step 5: Combining.** From Steps 1–4,

$$\langle \tilde{u}_{\ell,n}, F_n\rangle^2 = \frac{\|Q^*_{\ell,n} R_n(\lambda_{\ell,n}) U_n a_n\|^2}{\|R_n(\lambda_{\ell,n}) U_n a_n\|^2} \;\xrightarrow{P}\; \frac{|\mu|^{-2}}{(|\mu|^2-1)^{-1}} = 1 - \frac{1}{|\mu|^2}.$$

**Step 6: Cross-spike orthogonality (part (b)).** The same decomposition gives $\langle \tilde{u}_{\ell,n}, F_{\ell',n}\rangle^2 \approx (|\mu|^2 - 1)\, |\mu|^{-2}\, \|Q^*_{\ell',n} Q_{\ell,n} c_n\|^2/\|c_n\|^2$. Since $F_n \perp F_{\ell',n}$, we have $Q^*_{\ell',n} Q_{\ell,n} = 0$, giving the limit 0.

# Key Assumptions Recap

1. **Sub-Gaussian entries:** $\mathbb{P}(|A_{11}| \geq t) \leq 2\exp(-Ct^2)$.
2. **Sparsity regime:** $K_n \to \infty$ and $\log^9 n/K_n \to 0$ (needed for resolvent norm bounds via universality results of Brailovskaya–van Handel).
3. **Biorthogonal decomposition:** $E_n = P_n\Lambda_n W_n^*$ with $W_n^* P_n = I_r$ and eigenvalues converging to fixed limits outside the unit disk.

## References


[1] W. Hachem, M. Louvaris, J. Najim, *Extreme eigenvalues and eigenvectors for finite rank additive deformations of non-Hermitian sparse random matrices*, arXiv:2602.20956, 2026.

[2] T. Brailovskaya, R. van Handel, *Universality and sharp matrix concentration inequalities*, Geom. Funct. Anal. 34(6):1734–1838, 2024.

[3] F. Benaych-Georges, R. R. Nadakuditi, *The eigenvalues and eigenvectors of finite, low rank perturbations of large random matrices*, Adv. Math. 227(1):494–521, 2011.

[4] T. Tao, *Outliers in the spectrum of i.i.d. matrices with bounded rank perturbations*, Probab. Theory Relat. Fields 155(1):231–263, 2013.

[5] C. Bordenave, M. Capitaine, *Outlier eigenvalues for deformed i.i.d. random matrices*, Comm. Pure Appl. Math. 69(11):2131–2194, 2016.