Reading Report: *Lectures on Open Quantum Systems*

arXiv:2603.11925 — M. Merkli and Á. Neira

Daily arXiv Theory Report

March 13, 2026

1 Core Question

How does *irreversible* dynamics emerge from the fundamentally *unitary* (reversible) evolution of a quantum system coupled to a reservoir, and what is the precise mathematical structure of the generators that govern such dissipative dynamics?

2 Main Statement

Theorem 1 (GKSL Theorem [1, Thm. 3])**.** *Let $\mathcal{H} = \mathbb{C}^d$. A linear superoperator $\mathcal{L}$ on $\mathcal{B}(\mathcal{H})$ generates a completely positive, trace-preserving (CPTP) semigroup $\Phi_t = e^{t\mathcal{L}}$ if and only if there exist*

- *a self-adjoint operator $H = H^\dagger \in \mathcal{B}(\mathcal{H})$,*
- *operators $V_l \in \mathcal{B}(\mathcal{H})$, $l = 1, \ldots, d^2 - 1$,*
- *non-negative numbers $\gamma_l \geq 0$,*

such that for every $X \in \mathcal{B}(\mathcal{H})$,

$$\mathcal{L}X \;=\; -i[H, X] \;+\; \sum_{l=1}^{d^2-1} \gamma_l \Big(V_l \, X \, V_l^\dagger - \tfrac{1}{2}\{V_l^\dagger V_l,\, X\} \Big).$$

3 A Concrete Example: The Dissipative Jaynes–Cummings Model

Consider a two-level atom ($\mathcal{H}_S = \mathbb{C}^2$, ground/excited states $|0\rangle, |1\rangle$) coupled to a bosonic reservoir of harmonic oscillators (the cavity field $\mathcal{H}_R$). Define the raising/lowering operators $\sigma_\pm = |1\rangle\langle 0|, |0\rangle\langle 1|$ and bosonic operators $b_k, b_k^\dagger$ satisfying $[b_k, b_l^\dagger] = \delta_{kl}$. The total Hamiltonian is

$$H = \underbrace{\omega_0 \, \sigma_+ \sigma_-}_{H_S} + \underbrace{\sum_k \omega_k \, b_k^\dagger b_k}_{H_R} + \underbrace{\sigma_+ \otimes B + \sigma_- \otimes B^\dagger}_{H_I}, \qquad B = \sum_k g_k \, b_k.$$

The total excitation number $N = |1\rangle\langle 1| + \sum_k b_k^\dagger b_k$ commutes with H, making the model exactly solvable in the single-excitation sector.

In the **continuous-mode limit** with a Lorentzian spectral density

$$J(\omega) = \frac{g}{\pi} \, \frac{\Gamma}{(\omega - \omega_c)^2 + \Gamma^2},$$

the excited-state amplitude evolves as

$$c_1(t) = c_1(0) \, e^{-(\Gamma - i\Delta)t/2} \Big[\cosh\!\big(\tfrac{Rt}{2}\big) + \frac{\Gamma - i\Delta}{R} \, \sinh\!\big(\tfrac{Rt}{2}\big) \Big], \quad R = \sqrt{(\Gamma - i\Delta)^2 - \frac{4g}{\sqrt{2\pi}}},$$

where $\Delta = \omega_0 - \omega_c$ is the detuning. For large times $c_1(t) \to 0$, so $\rho_S(t) \to |0\rangle\langle 0|$: the atom irreversibly relaxes to its ground state, and the exact master equation takes Lindblad form with a time-dependent decay rate $\gamma(t)$.

4 Intuition and Setup

The paper is a self-contained set of lecture notes structured around a single guiding idea:

> *Start from a concrete, exactly solvable model (the Jaynes–Cummings model), observe the structures that appear (Lindblad generators, Kraus operators, CPTP maps), then develop the general theory.*

Key definitions and objects:

Definition 1 (Density matrix)**.** *A* density matrix *on a Hilbert space $\mathcal{H}$ is an operator $\rho \geq 0$ with $\operatorname{tr}\rho = 1$. If $\rho = |\psi\rangle\langle\psi|$ it is a* pure state*; otherwise it is* mixed*.*

Definition 2 (CPTP map)**.** *A linear map $\Phi : \mathcal{B}(\mathcal{H}) \to \mathcal{B}(\mathcal{H})$ is* completely positive trace-preserving *(CPTP) if it preserves trace and $\Phi \otimes \mathrm{id}_{n\times n}$ is positivity-preserving for every $n \geq 1$.*

Definition 3 (Quantum dynamical semigroup)**.** *A family $\{\Phi_t\}_{t\geq 0}$ of CPTP maps with $\Phi_0 = \mathrm{id}$, $\Phi_{t+s} = \Phi_t \circ \Phi_s$, and $t \mapsto \Phi_t$ continuous, is a* quantum dynamical (Markovian) semigroup*.*

Definition 4 (Reduced dynamics)**.** *For a bipartite system $\mathcal{H}_{SR} = \mathcal{H}_S \otimes \mathcal{H}_R$ evolving under $U = e^{-itH}$, the* reduced system state *is $\rho_S(t) = \operatorname{tr}_R\big(U\,(\rho_S(0) \otimes \rho_R)\,U^\dagger\big)$.*

5 Main Results (Bulleted)

- **Jaynes–Cummings exact master equation.** In the single-excitation sector with Lorentzian spectral density, the atom density matrix satisfies

$$\partial_t \rho_S(t) = -i[H_{\mathrm{eff}}(t),\, \rho_S(t)] + \gamma(t)\Big(\sigma_-\rho_S(t)\,\sigma_+ - \tfrac{1}{2}\{\sigma_+\sigma_-,\, \rho_S(t)\}\Big),$$

  an exact (non-Markovian) equation with time-dependent $\gamma(t)$.

- **Kraus Representation Theorem (Thm. 1).** Φ is CPTP on $\mathcal{B}(\mathbb{C}^d)$ if and only if there exist operators K_α, $\alpha = 1, \ldots, d^2$, with $\sum_\alpha K_\alpha^\dagger K_\alpha = \mathbb{1}$, such that $\Phi(X) = \sum_\alpha K_\alpha\, X\, K_\alpha^\dagger$.

- **Dilation Theorem (Thm. 2).** Φ is CPTP if and only if there exist a Hilbert space $\mathcal{H}_R$ ($\dim \leq d^2$), a unit vector $\Omega \in \mathcal{H}_R$, and a unitary U on $\mathcal{H} \otimes \mathcal{H}_R$ such that $\Phi(X) = \operatorname{tr}_R\big[U(X \otimes |\Omega\rangle\langle\Omega|)U^\dagger\big]$. This links abstract CPTP maps to the physics of open systems.

- **GKSL Theorem (Thm. 3).** The generator $\mathcal{L}$ of a CPTP semigroup must have Lindblad form (see Main Statement above).

6 Proof Sketch

Step 1: Derive irreversibility from the Jaynes–Cummings model

1. Work in the interaction picture $\varphi(t) = e^{iH_0 t}\psi(t)$.
2. Exploit the conservation law $[H, N] = 0$ to reduce to the single-excitation sector; the state is parameterized by amplitudes $c_1(t)$ (atom excited, vacuum field) and $d_k(t)$ (atom ground, one photon in mode k).

3. Eliminate $d_k(t)$ to get an integro-differential equation for $c_1(t)$ involving the vacuum correlation function $f(\tau) = \sum_k |g_k|^2 e^{-i(\omega_k - \omega_0)\tau}$.
4. Pass to the **continuous-mode limit**: replace the discrete sum by an integral against the spectral density $J(\omega) = \sqrt{2\pi}\, \varrho(\omega)|g(\omega)|^2$.
5. With a Lorentzian $J(\omega)$, solve explicitly to get $c_1(t)$ in closed form; show $c_1(t) \to 0$ (irreversibility).
6. Compute $\rho_S(t) = \mathrm{tr}_R\, |\varphi(t)\rangle\langle\varphi(t)|$ and derive the exact master equation (Lindblad form with time-dependent $\gamma(t)$).

Step 2: Kraus Representation (Thm. 1)

1. Start from the "open-system" definition $\Phi(X) = \mathrm{tr}_R[U(X \otimes |\Omega\rangle\langle\Omega|)U^\dagger]$.
2. Expand U in a product basis and take the partial trace to extract operators K_α such that $\Phi(X) = \sum_\alpha K_\alpha X K_\alpha^\dagger$.
3. For the converse, given a CPTP map Φ, use the anti-linear conjugation C and the maximally entangled vector $v = \sum_j e_j \otimes e_j$ to express $\Phi(|\psi\rangle\langle\psi|)$ as a partial trace of $(\Phi \otimes \mathrm{id})(|v\rangle\langle v|)$ contracted against $|C\psi\rangle\langle C\psi|$.
4. Complete positivity guarantees $(\Phi \otimes \mathrm{id})(|v\rangle\langle v|) \geq 0$, so it decomposes as $\sum_\alpha |s_\alpha\rangle\langle s_\alpha|$, yielding operators K_α via the coordinate expansion of each s_α.
5. Extend from rank-one projections to all X by linearity and Hermitian decomposition.

Step 3: Dilation Theorem (Thm. 2)

1. Given Kraus operators K_α, $\alpha = 1, \ldots, N$ ($N \leq d^2$), set $\mathcal{H}_R = \mathbb{C}^N$ with orthonormal basis $\{e_\alpha\}$.
2. Define the partial isometry $\bar{U} : \psi \otimes \Omega \mapsto \sum_\alpha K_\alpha \psi \otimes e_\alpha$.
3. Verify $\bar{U}$ is isometric (using $\sum_\alpha K_\alpha^\dagger K_\alpha = \mathbb{1}$) and extend to a full unitary U on $\mathcal{H} \otimes \mathcal{H}_R$ by Exercise 5 (extending a partial isometry).
4. Check $\mathrm{tr}_R[U(X \otimes |\Omega\rangle\langle\Omega|)U^\dagger] = \Phi(X)$.

Step 4: GKSL Theorem (Thm. 3)

1. Fix an orthonormal basis $\{F_j\}_{j=1}^{d^2}$ of $\mathcal{B}(\mathbb{C}^d)$ with $F_{d^2} = \frac{1}{\sqrt{d}}\mathbb{1}$.
2. Expand the Kraus representation of Φ_t in this basis: $\Phi_t(X) = \sum_{i,j} c_{ij}(t)\, F_i X F_j^\dagger$.
3. Differentiate at $t = 0$ to get $\mathcal{L}X = \lim_{t \to 0^+} \frac{1}{t}(\Phi_t(X) - X)$; the coefficient matrix (a_{ij}) has well-defined limits.
4. Use trace-preservation ($\mathrm{tr}(\mathcal{L}X) = 0$ for all X) to express the "diagonal" terms in terms of the a_{ij}, producing the Hamiltonian part $-i[H, X]$ and the anti-commutator corrections.
5. Show the coefficient matrix $A = (a_{ij})_{i,j=1}^{d^2-1}$ is positive semi-definite (inherited from positivity of the $c_{ij}(t)$ matrix).
6. Diagonalize $A = UDU^\dagger$ with $D = \mathrm{diag}(\gamma_1, \ldots, \gamma_{d^2-1})$, $\gamma_l \geq 0$, and define $V_l = \sum_i U_{il} F_i$ to arrive at Lindblad form.

References

[1] M. Merkli and Á. Neira, *Lectures on Open Quantum Systems*, arXiv:2603.11925, 2026.