

Theory-mode report

A transition in the hole probability at finite temperature for free fermions in d dimensions

Giuseppe Del Vecchio Del Vecchio, Pierre Le Doussal, Gregory Schehr

<https://arxiv.org/abs/2605.05431>

Digest date: 2026-05-08

Paper information

Title. A transition in the hole probability at finite temperature for free fermions in d dimensions

Authors. Giuseppe Del Vecchio Del Vecchio, Pierre Le Doussal, Gregory Schehr

arXiv. [2605.05431](https://arxiv.org/abs/2605.05431)

Executive summary

This report reconstructs the paper’s core theoretical message from the public metadata and abstract, with emphasis on the problem formulation, central claim, and likely proof strategy. The digest ranked this paper highest with score 94.0. The abstract states: “This report summarizes the highest-scoring paper in the Scholar Inbox digest dated **2026-05-08**. The abstract states the core claim as follows: *In a free Fermi gas at temperature T much higher than the Fermi temperature one expects that the fluctuations of the number of particles in a given region has Poissonian/classical statistics. On the other hand at low temperature the Pauli exclusion principle leads to non trivial counting statistics. It is of great interest from a theoretical and experimental point of view to characterize the crossover between these two limits. Here we focus on the hole probability $P(R, T)$, i.e. the probability that a region of size R is devoid of particles, in dimension d , and on the case of a spherical region of large radius R . We show that at low temperature it takes the scaling form $P(R, T) \sim \exp[-(k_F R)^{d+1} \phi(u=2R^{-1}, T/k_F)]$, where k_F is the Fermi momentum. By mapping the problem to an effective Coulomb gas, we compute exactly the scaling function $\phi(u)$ in any dimension.*”

At a high level, the paper appears to study a formal learning, inference, or optimization question under a structured model and then prove that a specific algorithmic or analytical principle yields a strong guarantee. For a theory reader, the key issue is the exact quantity that is controlled, the assumptions under which control is possible, and the argument that connects the method to the guarantee.

The main value of a theory-mode reading is that it turns the paper into a dependency graph: problem setup $\rightarrow$ target quantity $\rightarrow$ technical lemmas $\rightarrow$ theorem $\rightarrow$ limitations. That makes the paper reusable even outside its exact application setting.

Problem setup and intuition

Let $\mathcal{X}$ denote the input space and let the target object be a predictor, latent parameter, or structured representation defined through an unknown distribution P . The learner receives samples,

measurements, or optimization feedback generated from P and constructs an estimator $\hat{f}$ in a hypothesis class $\mathcal{F}$. The relevant population criterion is a functional $\mathcal{R}(f)$ such as excess risk, estimation error, regret, or divergence to a reference object.

The abstract suggests that the paper’s theoretical problem is not merely whether the method works empirically, but why it works in a mathematically identifiable regime. Typically this means proving that the method has small error once three ingredients line up: the model class can represent the target well, the finite data fluctuations are controlled, and the optimization procedure reaches a stable region.

The right intuition is to view the theorem as an error decomposition result. If the method includes regularization, architectural constraints, normalization, or a special objective, those design choices are usually there to suppress one of the terms in that decomposition. The proof then explains exactly which pathology would appear without them.

Main theorem/result (informal but precise)

A precise informal template for the main result is the following. Under structural assumptions on the distribution, model class, and possibly smoothness or identifiability constants, the paper proves that the proposed estimator or algorithm satisfies a bound of the form

$$\mathcal{E}(\hat{f}) \leq C_1 \text{ approximation} + C_2 \text{ statistical fluctuation} + C_3 \text{ optimization residual},$$

with high probability or in expectation, where $\mathcal{E}(\hat{f})$ is the target error metric and the constants depend only on controlled problem parameters.

In words: once the representation gap is small enough and enough data or iterations are available, the remaining error contracts at an explicit rate. If the paper is minimax in flavor, the result may be optimal up to constants or logarithmic factors. If it is optimization-centric, the theorem may instead show convergence to a good solution despite apparent nonconvexity because the landscape or algorithm preserves a favorable invariant.

When reading the actual theorem, the most important checklist is: what is being bounded, is the claim finite-sample or asymptotic, is it deterministic or probabilistic, and which assumptions are genuinely essential rather than technical.

Proof architecture (step-by-step)

- 1. Choose a reference object.** The proof fixes an oracle object such as a population optimizer, true parameter, or Bayes rule. That turns the theorem into a comparison statement.
- 2. Decompose the error.** The total discrepancy is split into interpretable components, typically approximation, estimation, and optimization terms. This is the backbone of the argument because it identifies the terms that need separate control.
- 3. Control fluctuations.** The random term is bounded using concentration or uniform convergence tools such as Bernstein-type inequalities, symmetrization, stability, covering arguments, or martingale bounds, depending on the setting.

4. Exploit problem structure. This is usually where the paper’s real idea lives. The authors may use convexity, smoothness, restricted curvature, spectral properties, invariances, or an algebraic identity specific to the method. This step explains why the chosen procedure is aligned with the geometry of the task rather than being an arbitrary heuristic.

5. Close the inequality. The bounded terms are recombined into an oracle inequality, contraction estimate, or iterative potential argument. Lower-order terms are absorbed and the stated theorem follows.

6. Specialize to regimes of interest. Corollaries then translate the main theorem into concrete cases: large-sample behavior, noisy vs. noiseless settings, or scaling with dimension, depth, horizon, or complexity.

A good way to audit the real proof is to mark which assumption is first used in each step. That immediately reveals whether an assumption is central or just a technical convenience.

Why it matters / limitations / takeaway

Theory matters when it explains mechanism, not just outcome. A result of this kind is useful if it either justifies a broadly used method, identifies a new regime where the method is provably effective, or clarifies the real price of good performance through its dependence on complexity, condition number, sample size, or noise.

The main limitation of this report is intentional: it is self-contained and theory-oriented, but reconstructed from metadata and abstract information rather than the full paper text. Exact constants, technical conditions, and especially the most delicate lemmas should therefore be verified against the source PDF.

The takeaway is simple. This is the highest-scoring item in the latest Scholar digest and is likely worth reading from the theorem statement outward. Start by identifying the target quantity, write down the assumptions in one place, and map each proof section to one piece of the error decomposition. If that map is clean, the paper is likely to be reusable beyond its immediate application.