Reading Report: Quadratic Forms of Heavy-Tailed Self-Normalized Random Vectors and the α-Heavy Marčenko–Pastur Law

arXiv:2603.07132 — Z. Dong, J. Heiny, J. Yao

March 2026

1 Core Question

When the entries of a data matrix have *infinite variance* (regularly varying tails with index $\alpha \in (0, 2)$), the sample correlation matrix has a limiting spectral distribution called the α-heavy Marčenko–Pastur (MP) law $H_{\alpha,\gamma}$. This paper asks: **what are the analytic properties of** $H_{\alpha,\gamma}$—in particular, does it have atoms (point masses), what is its density, and how can it be represented via the resolvent?

2 Setup and Key Definitions

Self-normalized random vectors. Let ξ be a symmetric random variable that is *regularly varying* with index $\alpha \in (0, 2)$: that is,

$$P(|\xi| > x) = x^{-\alpha} L(x), \qquad x > 0,$$

where L is slowly varying at ∞. In particular, $E[\xi^2] = \infty$.

Let $\mathbf{x} = (X_1, \ldots, X_n)^\top$ consist of i.i.d. copies of ξ, and define the *self-normalized* vector

$$\mathbf{y} = \frac{\mathbf{x}}{\|\mathbf{x}\|_2} \in S^{n-1}.$$

Random quadratic forms. Given an $n \times n$ Hermitian matrix $A_n = (a_{ij}^{(n)})$ independent of $\mathbf{y}$, the object of study is the quadratic form

$$Q_n = \mathbf{y}^\top A_n \, \mathbf{y}.$$

Decompose $Q_n = Q_{n,1} + Q_{n,2}$, where $Q_{n,1} = \mathbf{y}^\top \operatorname{diag}(A_n)\mathbf{y}$ (diagonal part) and $Q_{n,2}$ (off-diagonal part). The authors show (Proposition 2.3) that $Q_{n,2} \xrightarrow{P} 0$ whenever $n^{-2} E[\|A_n - \operatorname{diag}(A_n)\|_F^2] \to 0$, so the *diagonal entries alone* govern the limit.

Stieltjes transform. A probability measure μ on $\mathbb{R}$ is characterized by its *Stieltjes transform*

$$s_\mu(z) = \int_{\mathbb{R}} \frac{1}{x - z} \, \mu(dx), \qquad z \in \mathbb{C} \setminus \operatorname{supp}(\mu).$$

Sample correlation matrix. From an $p \times n$ data matrix $X = (X_{ij})$ with i.i.d. entries distributed as ξ, define $S_n = n^{-1} X X^\top$ (sample covariance) and

$$R_n = \{\operatorname{diag}(S_n)\}^{-1/2} \, S_n \, \{\operatorname{diag}(S_n)\}^{-1/2}$$

the *sample correlation matrix*. In the proportional regime $p/n \to \gamma \in (0, \infty)$, the empirical spectral distribution of R_n converges to the α-heavy MP law $H_{\alpha,\gamma}$ (Heiny–Yao, 2022).

3 Concrete Example

Example 1. Let $\nu = \frac{1}{2}\delta_0 + \frac{1}{2}\delta_1$ (the simplest non-degenerate case). Then the limiting measure $\mu_{\nu,\alpha}$ of $Q_{n,1}$ is supported on $[0,1]$ with explicit density

$$f_{\nu,\alpha}(x) = \frac{\sin\frac{\alpha\pi}{2}}{\pi} \frac{(1-x)^{\alpha/2-1}\, x^{\alpha/2-1}}{x^\alpha + (1-x)^\alpha + 2\cos\frac{\alpha\pi}{2}\, x^{\alpha/2}(1-x)^{\alpha/2}}\, \mathbf{1}_{(0,1)}(x).$$

At $\alpha = 1$ this becomes $f_{\nu,1}(x) = \frac{1}{\pi\sqrt{x(1-x)}}$, the arcsine law—a distribution familiar from random walks and beta ensembles. As $\alpha \uparrow 2$, the density concentrates at the mean $x = 1/2$; as $\alpha \downarrow 0$, it converges to ν itself (two atoms).

4 Intuition: Why Diagonals Dominate

In the *light-tailed* regime ($E[\xi^2] < \infty$), mixed moments satisfy $\beta_{2k_1,\ldots,2k_r} = O(n^{-(k_1+\cdots+k_r)})$, which is tiny. The quadratic form Q_n then concentrates around its mean (Hanson–Wright inequality).

In the *heavy-tailed* regime ($\alpha < 2$), Lemma 2.1 shows

$$n^r\, \beta_{2k_1,\ldots,2k_r} \longrightarrow \frac{(\alpha/2)^{r-1}\,\Gamma(r)\prod_{j=1}^r \Gamma(k_j - \alpha/2)}{\Gamma(1-\alpha/2)^r\,\Gamma(k_1+\cdots+k_r)},$$

so mixed moments are $O(n^{-r})$ regardless of the k_j's. This means only the *number of distinct indices* r matters, not the powers k_j. Consequently, the off-diagonal contribution vanishes in probability while the diagonal part retains a *non-degenerate* random limit—a striking departure from the classical setting.

5 Main Results

- **Theorem 2.4 (Limiting law, bounded case).** If $\max_i |a_{ii}^{(n)}| \le M < \infty$ a.s. and $n^{-1}\sum_i \delta_{a_{ii}^{(n)}} \xrightarrow{w} \nu$ a.s. (deterministic), then $Q_{n,1} \xrightarrow{d} \mu_{\nu,\alpha}$ whose Stieltjes transform is

$$\boxed{s_{\mu_{\nu,\alpha}}(z) = -\frac{\displaystyle\int_{\mathbb{R}} (z-x)^{\alpha/2-1}\,\nu(dx)}{\displaystyle\int_{\mathbb{R}} (z-x)^{\alpha/2}\,\nu(dx)}, \qquad z \in \mathbb{C}\setminus \operatorname{supp}(\nu).}$$

- **Theorem 2.10 (Density and no atoms).** If ν is non-degenerate, $\mu_{\nu,\alpha}$ is *absolutely continuous* with an explicit density $f_{\nu,\alpha}$ (formula (2.7)), strictly positive on $(\inf\operatorname{supp}(\nu), \sup\operatorname{supp}(\nu))$ and zero outside.

- **Theorem 2.12 (Unbounded diagonal extension).** Theorem 2.4 extends to unbounded diagonals under the uniform integrability condition $\lim_{T\to\infty}\limsup_n n^{-1}\sum_i |a_{ii}^{(n)}|\mathbf{1}(|a_{ii}^{(n)}| > T) = 0$ a.s.

- **Theorem 3.3 (Stieltjes transform of $H_{\alpha,\gamma}$).** There exists a random holomorphic function $\psi : \mathbb{C}\setminus[0,\infty) \to \mathbb{C}$ (a subsequential limit of the resolvent diagonal $B_n(z)_{11}$) such that

$$\boxed{s_{H_{\alpha,\gamma}}(z) = -\frac{E\big[(1+\psi(z))^{\alpha/2-1}\big]}{z\,E\big[(1+\psi(z))^{\alpha/2}\big]}, \qquad z \in \mathbb{C}\setminus[0,+\infty).}$$

- **Proposition 3.5 (No atoms for $H_{\alpha,\gamma}$).** $H_{\alpha,\gamma}(\{u\}) = 0$ for every $u > 0$. Combined with earlier work showing $H_{\alpha,\gamma}$ may have mass at 0 when $\gamma > 1$, this settles the atom question: the only possible atom is at the origin.

- **Boundary behavior (Proposition 2.7).** $\mu_{\nu,\alpha} \xrightarrow{w} \delta_{\int x\,\nu(dx)}$ as $\alpha \uparrow 2$ (classical concentration) and $\mu_{\nu,\alpha} \xrightarrow{w} \nu$ as $\alpha \downarrow 0$ (atoms emerge).

- **Tail behavior (Example 2.13).** When ν has exponential tails, $\mu_{\nu,\alpha}$ has Gamma-type tails $f_{\nu,\alpha}(x) \sim x^{-\alpha/2}e^{-x}$. When ν has polynomial tails of index s, $\mu_{\nu,\alpha}$ has the *same* polynomial index s, independent of α.

- **Hanson–Wright for self-normalized vectors (Theorem A.1).** When ξ is sub-Gaussian, a Hanson–Wright-type inequality is established for $Q_n = \mathbf{y}^\top A_n \mathbf{y}$, providing exponential concentration in the light-tailed benchmark.

6 Proof Sketch

Step 1. Off-diagonal vanishes. Using $E[|Q_{n,2}|^2] \le 2E[\|A_n - \mathrm{diag}(A_n)\|_F^2]\,\beta_{2,2}$ and $n^2\beta_{2,2} \le 1$, Markov's inequality gives $Q_{n,2} \xrightarrow{P} 0$.

Step 2. Moment computation for $Q_{n,1}$. Expand $E[Q_{n,1}^\ell]$ via the multinomial theorem. The key input is Lemma 2.1 on the asymptotics of the mixed moments $\beta_{2k_1,\dots,2k_r}$. Combined with $n^{-1}\sum_i \delta_{a_{ii}} \xrightarrow{w} \nu$ and boundedness, each moment m_ℓ is computed explicitly.

Step 3. Carleman's condition and moment determinacy. Since $|m_\ell| \le M^\ell$, Carleman's condition $\sum_\ell m_{2\ell}^{-1/(2\ell)} = \infty$ holds, proving that $\{m_\ell\}$ uniquely determines $\mu_{\nu,\alpha}$ and $Q_{n,1} \xrightarrow{d} \mu_{\nu,\alpha}$.

Step 4. Generating-function identification of the Stieltjes transform. Define $K_x(z) = \sum_{j\ge1} \frac{\Gamma(j-\alpha/2)}{\Gamma(1-\alpha/2)\,j!} \int x^j \nu(dx)\, z^j$ and recognize that $\sum m_\ell z^\ell/\ell = -\frac{2}{\alpha}\log\int(1-xz)^{\alpha/2}\nu(dx)$. Differentiating and rearranging yields the Stieltjes transform formula.

Step 5. No-atom proof for $\mu_{\nu,\alpha}$. At a candidate atom a, the non-tangential limit $\lim_{\angle z\to a}(a-z)s_{\mu_{\nu,\alpha}}(z)$ involves $(z-a)^{\alpha/2} \to 0$ in the numerator, while the denominator stays away from zero because ν is non-degenerate. Hence $\mu_{\nu,\alpha}(\{a\}) = 0$.

Step 6. Density via boundary values of Stieltjes transform. Apply $f_{\nu,\alpha}(x) = \frac{1}{\pi}\lim_{v\downarrow0}\mathrm{Im}\, s_{\mu_{\nu,\alpha}}(x+iv)$ and compute the boundary values of $(z-y)^{\alpha/2}$ and $(z-y)^{\alpha/2-1}$ using the principal branch. Verify integrability via the argument-of-F representation $f_{\nu,\alpha}(x) = -\frac{1}{\theta}\frac{d}{dx}\arg F(x)$.

Step 7. Random resolvent and $H_{\alpha,\gamma}$. From the recursive identity (3.2) relating $s_{R_n}(z)$ to $\mathbf{y}_1^\top B_{n,1}(z)\mathbf{y}_1$, extract a subsequential limit $\psi(z)$ of the resolvent diagonal via Prokhorov + diagonal-concentration (Proposition 3.1, martingale bounded-differences). Apply Theorem 2.4 to the quadratic form with $A_n = \mathrm{diag}(B_{n,1}(z))$, yielding the implicit formula (3.1).

Step 8. No atoms for $H_{\alpha,\gamma}$ on $(0,\infty)$. Assume $H_{\alpha,\gamma}(\{u\}) > 0$ for some $u > 0$. At $z = u + iv$, the identity (3.1) forces $E[(1+\psi(z))^{\alpha/2}] \to 0$ as $v \downarrow 0$. But a lower bound on $\mathrm{Im}\,\psi(z)$ (from $\mathrm{Im}\, s_{\widetilde{H}_{\alpha,\gamma}}(z) \sim w/v$) shows $P(\mathrm{Im}\,\psi(z) \ge w/(4v)) \ge p_0 > 0$, giving $E[\mathrm{Im}(1+\psi(z))^{\alpha/2}] \ge Cv^{-\alpha/2} \to \infty$—a contradiction.

References

Z. Dong, J. Heiny, J. Yao, *Quadratic form of heavy-tailed self-normalized random vector with applications in α-heavy Marčenko–Pastur law*, arXiv:2603.07132 (2026).

J. Heiny, J. Yao, *Limiting distributions for eigenvalues of sample correlation matrices from heavy-tailed populations*, Ann. Statist. 50 (2022), 3249–3280.

V. A. Marčenko, L. A. Pastur, *Distribution of eigenvalues of some sets of random matrices*, Mat. Sb. 72 (1967), 507–536.