Reading Report: Upper Tail Large Deviations for Extremal Eigenvalues of the Elliptic Ginibre Matrices

Auto-generated report on `arXiv:2603.16339`
by S.-S. Byun, Y.-W. Lee, and S. Oh

March 18, 2026

1 Core Question

What is the probability that the largest eigenvalue (either the spectral radius $\max|z_j|$ or the rightmost eigenvalue $\max \operatorname{Re} z_j$) of an *elliptic Ginibre matrix* exits the support of the limiting spectral distribution, and how does this probability decay as the matrix size $N \to \infty$? The paper answers this for all three symmetry classes—real ($\beta = 1$), complex ($\beta = 2$), and symplectic ($\beta = 4$)—in a unified framework.

2 Main Result at a Glance

Theorem 1 (Theorem 2.1 of [1])**.** *Let $\tau \in [0, 1)$ be the non-Hermiticity parameter. For $s > 1+\tau$ (i.e. outside the elliptic law support),*

$$\lim_{N\to\infty} \frac{1}{\beta N} \log \mathbb{P}\Big[\max_j |z_j| \geq s\Big] \;=\; \lim_{N\to\infty} \frac{1}{\beta N} \log \mathbb{P}\Big[\max_j \operatorname{Re} z_j \geq s\Big] \;=\; -\Phi_\tau(s),$$

where $\beta \in \{1, 2, 4\}$ is the Dyson index and the rate function is

$$\Phi_\tau(s) \;:=\; \frac{1}{2}\left(\frac{1}{1+\tau} - \frac{1}{2\tau}\right)s^2 \;+\; \frac{s\sqrt{s^2-4\tau}}{4\tau} \;-\; \log \frac{s+\sqrt{s^2-4\tau}}{2}\,.$$

Limiting cases.

- $\tau = 0$ (Ginibre): $\Phi_0(s) = \frac{1}{2}(s^2 - 2\log s - 1)$, recovering the circular-law result of Lacroix-A-Chez-Toine–Majumdar–Schehr and Xu–Zeng.
- $\tau \to 1$ (Hermitian/GOE–GUE–GSE): $\Phi_1(s) = \frac{s\sqrt{s^2-4}}{4} - \log \frac{s+\sqrt{s^2-4}}{2}$, the classical Dean–Majumdar rate function.

3 A Concrete Example

Example 2. Consider a 100×100 complex elliptic Ginibre matrix ($\beta = 2$) with $\tau = 0.3$. The elliptic law is supported on the ellipse with semi-axes $1 + \tau = 1.3$ (real) and $1 - \tau = 0.7$ (imaginary). Take $s = 1.5$. Then

$$\Phi_{0.3}(1.5) \;\approx\; 0.038,$$

so the probability that *any* eigenvalue has modulus ≥ 1.5 is roughly

$$\mathbb{P}\big[\max|z_j| \geq 1.5\big] \;\approx\; e^{-\beta N\,\Phi_\tau(s)} = e^{-2\cdot 100\cdot 0.038} \approx 0.05\%.$$

This illustrates the exponential suppression: even a modest overshoot past the support boundary is extremely unlikely for large N.

4 Setup and Intuition

The elliptic Ginibre ensemble

Let G be an $N \times N$ random matrix with i.i.d. centred Gaussian entries of variance $1/N$ (real, complex, or quaternionic). Define

$$X_\tau \;:=\; \sqrt{\tfrac{1+\tau}{2}}\,(G+G^\dagger) \;+\; \sqrt{\tfrac{1-\tau}{2}}\,(G-G^\dagger), \qquad \tau \in [0,1).$$

- $\tau = 0$: pure Ginibre (maximally non-Hermitian).
- $\tau \to 1$: Hermitian (GOE/GUE/GSE).

The eigenvalues $\{z_j\}_{j=1}^N \subset \mathbb{C}$ and, as $N \to \infty$, their empirical measure converges to the **elliptic law**:

$$\frac{1}{\pi(1-\tau^2)}\,\mathbf{1}_{\mathcal{E}}(z)\,d^2z, \qquad \mathcal{E} = \Big\{(x,y) : \frac{x^2}{(1+\tau)^2} + \frac{y^2}{(1-\tau)^2} \le 1\Big\}.$$

Why spectral radius and rightmost eigenvalue?

The *spectral radius* $\max|z_j|$ measures whether eigenvalues escape the support radially. The *rightmost eigenvalue* $\max \operatorname{Re} z_j$ is the key stability indicator in continuous-time dynamical systems (e.g. May's model of ecosystem stability): if it crosses the imaginary axis, the system becomes unstable.

Key insight: same rate function for both

Despite the different geometry of the two events—$\{|z| \ge s\}$ (annular) vs. $\{\operatorname{Re} z \ge s\}$ (half-plane)—the large deviation rate is the same. This is because the function $\Omega(z)$ (defined below) achieves its minimum on the real axis in both cases, so the "cheapest" way for the system to produce an outlier eigenvalue is always along the real line.

5 Main Results (Detailed)

- **Theorem 2.1** (stated above): Unified rate function Φ_τ for spectral radius *and* rightmost eigenvalue, valid for all $\beta \in \{1,2,4\}$. In particular, the $\beta = 4$ (symplectic) case was *previously unknown* even for $\tau = 0$.

- **Theorem 2.2** (general domain): Let $U \subset \mathbb{C}$ be measurable and bounded away from $\mathcal{E}$. Define

  $$\Omega(z) := \frac{1}{1-\tau^2}\big(|z|^2 - \tau \operatorname{Re} z^2\big) - \operatorname{Re}\frac{2z}{z+\sqrt{z^2-4\tau}} - 2\log\Big|\frac{z+\sqrt{z^2-4\tau}}{2}\Big|.$$

  Then for $\beta = 2,4$:

  $$\lim_{N\to\infty}\frac{1}{\beta N}\log \mathbb{P}\big[\exists\, z_j \in U\big] = -\tfrac{1}{2}\operatorname{ess\,inf}_{z\in U}\,\Omega(z).$$

  For $\beta = 1$ (eGinOE):

  $$\lim_{N\to\infty}\frac{1}{N}\log \mathbb{P}\big[\exists\, z_j \in U\big] = -\min\Big\{\operatorname{ess\,inf}_{z\in U\setminus\mathbb{R}}\Omega(z),\ \tfrac{1}{2}\operatorname{ess\,inf}_{z\in U\cap\mathbb{R}}\Omega(z)\Big\}.$$

  The real eigenvalues (which exist only for $\beta = 1$) have a different exponential weight, and it is *always* a real eigenvalue that is the most likely to cause the large-deviation event.

- **Propositions 4.2–4.4** (one-point function asymptotics): For $z \in \mathcal{E}^c$, the one-point functions satisfy

$$\begin{aligned} R_N^{\mathbb{C}}(z) &= e^{-N\Omega(z)+o(N)} && \text{(eGinUE)}, \\ R_N^{\mathbb{H}}(z) &= e^{-2N\Omega(z)+o(N)} && \text{(eGinSE)}, \\ R_N^{\mathbb{R},r}(x) &= e^{-N\Omega(x)/2+o(N)}, \quad R_N^{\mathbb{R},c}(z) = e^{-N\Omega(z)+o(N)} && \text{(eGinOE)}. \end{aligned}$$

  These are derived with *precise* prefactors (not just leading exponential order), which is of independent interest.

- **Potential-theoretic interpretation**: $\Omega(z) = V(z) - \check{V}(z)$, where V is the external potential and $\check{V}$ is the obstacle function from logarithmic potential theory. The rate function is the "energy cost" of placing a charge at z outside the equilibrium support.

6 Proof Sketch

The proof proceeds through the following steps.

Step 1. Exact finite-N formulas for one-point functions.
Using the determinantal (eGinUE) and Pfaffian (eGinOE, eGinSE) structures of the eigenvalue point processes, the one-point functions are expressed as sums over planar orthogonal/skew-orthogonal polynomials built from Hermite polynomials H_n. Specifically, define monic polynomials

$$\varphi_{N,j}(z) = \Big(\frac{\tau}{2N}\Big)^{j/2} H_j\Big(\sqrt{\frac{N}{2\tau}}\, z\Big).$$

The eGinUE kernel is $K_N^{\mathbb{C}}(z,w) = \sum_{j=0}^{N-1} \varphi_{N,j}(z)\overline{\varphi_{N,j}(w)}/h_{N,j}^{\mathbb{C}}$, and the one-point function is $R_N^{\mathbb{C}}(z) = \omega_N^{\mathbb{C}}(z)\, K_N^{\mathbb{C}}(z,\bar{z})$.

Step 2. Differential-operator reduction (Lee–Riser trick).
Direct asymptotic analysis of the large sum $K_N^{\mathbb{C}}$ is hard. Instead, applying $\partial/\partial z$ to $R_N^{\mathbb{C}}(z)$ collapses the sum to a product of only the *two highest-degree* polynomials $\varphi_{N,N}$ and $\varphi_{N,N-1}$ (Lemma 3.1). Analogous connection formulas reduce eGinSE and eGinOE kernels to the eGinUE kernel plus finite-rank corrections (Lemmas 3.2, 3.3).

Step 3. Hermite polynomial asymptotics.
For z outside the ellipse $\mathcal{E}$, the rescaled Hermite polynomial $\varphi_{N,N+r}(z)$ admits the asymptotic expansion (Lemma 4.1):

$$\varphi_{N,N+r}(z) = \sqrt{h_{N,N+r}^{\mathbb{C}}} \cdot \frac{N^{1/4}}{(1-\tau^2)^{1/4}} \cdot \frac{\psi(z)^r \sqrt{\psi'(z)}}{\sqrt{2\pi^3}} \cdot e^{Ng(z)}\big(1 + O(1/N)\big),$$

where $\psi(z) = \frac{z+\sqrt{z^2-4\tau}}{2}$ is the inverse Joukowsky map and $g(z) = \frac{z}{2}\psi(z) + \log\psi(z)$.

Step 4. Laplace's method to recover $R_N^{\mathbb{C}}$.
Having the derivative $\partial_x R_N^{\mathbb{C}}(z) \sim e^{-N\Omega(z)}$ from Steps 2–3, one integrates in $x = \operatorname{Re} z$ from x to ∞. Since Ω is increasing outward, the integral is dominated by its lower endpoint, yielding $R_N^{\mathbb{C}}(z)$ with a precise prefactor (Proposition 4.2). Analogous arguments give Propositions 4.3 and 4.4 for eGinOE and eGinSE.

Step 5. From one-point functions to large deviations.
Upper bound: By a union bound / first-moment method,

$$\mathbb{P}[\exists\, z_j \in U] \;\le\; \int_U R_N(z)\, d^2z \;\le\; |U| \cdot \sup_{z\in U} R_N(z).$$

The exponential rate is then $-\inf_{z\in U}\Omega(z)$ (up to β-scaling).

Lower bound: One shows that the probability of having *exactly one* eigenvalue in a small ball around the minimiser z^* of Ω in U is at least $e^{-\beta N\,\Omega(z^*)/2+o(N)}$. This uses the one-point function as a lower bound for the inclusion probability, combined with repulsion estimates to control multi-eigenvalue contributions.

For the eGinOE ($\beta = 1$), the real and complex eigenvalue contributions must be handled separately. The key observation is that $R_N^{\mathbb{R},r}(x) \sim e^{-N\Omega(x)/2}$ while $R_N^{\mathbb{R},c}(z) \sim e^{-N\Omega(z)}$, so real eigenvalues dominate the large-deviation event (smaller exponent $\Rightarrow$ higher probability).

Step 6. Minimisation of Ω for specific domains.
For $U = \{|z| \ge s\}$ or $U = \{\operatorname{Re} z \ge s\}$, one verifies (Lemma 4.5) that

$$\inf_{z\in U} \Omega(z) = \Omega(s),$$

i.e. the infimum is achieved at $z = s \in \mathbb{R}$. Combined with Theorem 2.2, this yields Theorem 2.1 with $\Phi_\tau(s) = \frac{1}{2}\Omega(s)$.

References

[1] S.-S. Byun, Y.-W. Lee, and S. Oh, *Upper tail large deviations for extremal eigenvalues of the real, complex and symplectic elliptic Ginibre matrices*, arXiv:2603.16339 (2026).