

Finite Volume Methods for Hyperbolic Problems

Multidimensional Hyperbolic Problems

- Derivation of conservation law
- Hyperbolicity
- Advection
- Gas dynamics and acoustics
- Shear waves

Derivation of conservation law

$$\frac{d}{dt} \iint_{\Omega} q(x, y, t) dx dy = \text{net flux across } \partial\Omega.$$

Net flux is determined by integrating the flux of q normal to $\partial\Omega$ around this boundary.

Derivation of conservation law

$$\frac{d}{dt} \iint_{\Omega} q(x, y, t) dx dy = \text{net flux across } \partial\Omega.$$

Net flux is determined by integrating the flux of q normal to $\partial\Omega$ around this boundary.

$f(q)$ = flux of q in the x -direction,

$g(q)$ = flux of q in the y -direction,

(both per unit length in orthog direction, per unit time),

$$\vec{f}(q) = (f(q), g(q))$$

Derivation of conservation law

$$\frac{d}{dt} \iint_{\Omega} q(x, y, t) dx dy = \text{net flux across } \partial\Omega.$$

Net flux is determined by integrating the flux of q normal to $\partial\Omega$ around this boundary.

$f(q)$ = flux of q in the x -direction,

$g(q)$ = flux of q in the y -direction,

(both per unit length in orthog direction, per unit time),

$$\vec{f}(q) = (f(q), g(q))$$

$\vec{n}(s) = (n^x(s), n^y(s))$ outward-pointing unit normal $(x(s), y(s))$.

Flux at $(x(s), y(s))$ in the direction $\vec{n}(s)$:

$$\vec{n}(s) \cdot \vec{f}(q(x(s), y(s))) = f(q)n^x(s) + g(q)n^y(s),$$

Derivation of conservation law

$$\frac{d}{dt} \iint_{\Omega} q(x, y, t) \, dx \, dy = - \int_{\partial\Omega} \vec{n} \cdot \vec{f}(q) \, ds.$$

Derivation of conservation law

$$\frac{d}{dt} \iint_{\Omega} q(x, y, t) dx dy = - \int_{\partial\Omega} \vec{n} \cdot \vec{f}(q) ds.$$

If q is smooth: divergence theorem \implies

$$\frac{d}{dt} \iint_{\Omega} q(x, y, t) dx dy = - \iint_{\Omega} \vec{\nabla} \cdot \vec{f}(q) dx dy,$$

where the divergence of \vec{f} is

$$\vec{\nabla} \cdot \vec{f}(q) = f(q)_x + g(q)_y.$$

Derivation of conservation law

$$\frac{d}{dt} \iint_{\Omega} q(x, y, t) dx dy = - \int_{\partial\Omega} \vec{n} \cdot \vec{f}(q) ds.$$

If q is smooth: divergence theorem \implies

$$\frac{d}{dt} \iint_{\Omega} q(x, y, t) dx dy = - \iint_{\Omega} \vec{\nabla} \cdot \vec{f}(q) dx dy,$$

where the divergence of \vec{f} is

$$\vec{\nabla} \cdot \vec{f}(q) = f(q)_x + g(q)_y.$$

This leads to

$$\iint_{\Omega} \left[q_t + \vec{\nabla} \cdot \vec{f}(q) \right] dx dy = 0.$$

Derivation of conservation law

$$\frac{d}{dt} \iint_{\Omega} q(x, y, t) dx dy = - \int_{\partial\Omega} \vec{n} \cdot \vec{f}(q) ds.$$

If q is smooth: divergence theorem \implies

$$\frac{d}{dt} \iint_{\Omega} q(x, y, t) dx dy = - \iint_{\Omega} \vec{\nabla} \cdot \vec{f}(q) dx dy,$$

where the divergence of \vec{f} is

$$\vec{\nabla} \cdot \vec{f}(q) = f(q)_x + g(q)_y.$$

This leads to

$$\iint_{\Omega} \left[q_t + \vec{\nabla} \cdot \vec{f}(q) \right] dx dy = 0.$$

True for any $\Omega \implies q_t + \vec{\nabla} \cdot \vec{f}(q) = 0.$ (PDE form)

First order hyperbolic PDE in 2 space dimensions

General conservation law: $q_t + f(q)_x + g(q)_y = 0$

Quasi-linear form: $q_t + f'(q)q_x + g'(q)q_y = 0$

First order hyperbolic PDE in 2 space dimensions

General conservation law: $q_t + f(q)_x + g(q)_y = 0$

Quasi-linear form: $q_t + f'(q)q_x + g'(q)q_y = 0$

Constant coefficient linear system: $q_t + Aq_x + Bq_y = 0$

where $q \in \mathbb{R}^m$, $f(q) = Aq$, $g(q) = Bq$ and $A, B \in \mathbb{R}^{m \times m}$.

First order hyperbolic PDE in 2 space dimensions

General conservation law: $q_t + f(q)_x + g(q)_y = 0$

Quasi-linear form: $q_t + f'(q)q_x + g'(q)q_y = 0$

Constant coefficient linear system: $q_t + Aq_x + Bq_y = 0$

where $q \in \mathbb{R}^m$, $f(q) = Aq$, $g(q) = Bq$ and $A, B \in \mathbb{R}^{m \times m}$.

Advection equation: $q_t + uq_x + vq_y = 0$

First order hyperbolic PDE in 2 space dimensions

General conservation law: $q_t + f(q)_x + g(q)_y = 0$

Quasi-linear form: $q_t + f'(q)q_x + g'(q)q_y = 0$

Constant coefficient linear system: $q_t + Aq_x + Bq_y = 0$

where $q \in \mathbb{R}^m$, $f(q) = Aq$, $g(q) = Bq$ and $A, B \in \mathbb{R}^{m \times m}$.

Advection equation: $q_t + uq_x + vq_y = 0$

Hyperbolic if $\cos(\theta)f'(q) + \sin(\theta)g'(q)$ is diagonalizable with real eigenvalues, for all angles θ .

First order hyperbolic PDE in 2 space dimensions

General conservation law: $q_t + f(q)_x + g(q)_y = 0$

Quasi-linear form: $q_t + f'(q)q_x + g'(q)q_y = 0$

Constant coefficient linear system: $q_t + Aq_x + Bq_y = 0$

where $q \in \mathbb{R}^m$, $f(q) = Aq$, $g(q) = Bq$ and $A, B \in \mathbb{R}^{m \times m}$.

Advection equation: $q_t + uq_x + vq_y = 0$

Hyperbolic if $\cos(\theta)f'(q) + \sin(\theta)g'(q)$ is diagonalizable with real eigenvalues, for all angles θ .

Then plane wave propagating in any direction satisfies 1D hyperbolic equation.

Plane wave solutions

Suppose

$$\begin{aligned} q(x, y, t) &= \check{q}(x \cos \theta + y \sin \theta, t) \\ &= \check{q}(\xi, t). \end{aligned}$$

Plane wave solutions

Suppose

$$\begin{aligned}q(x, y, t) &= \check{q}(x \cos \theta + y \sin \theta, t) \\ &= \check{q}(\xi, t).\end{aligned}$$

Then:

$$\begin{aligned}q_x(x, y, t) &= \cos \theta \check{q}_\xi(\xi, t) \\ q_y(x, y, t) &= \sin \theta \check{q}_\xi(\xi, t)\end{aligned}$$

so

$$q_t + Aq_x + Bq_y = \check{q}_t + (A \cos \theta + B \sin \theta) \check{q}_\xi$$

and the 2d problem reduces to the 1d hyperbolic equation

$$\check{q}_t(\xi, t) + (A \cos \theta + B \sin \theta) \check{q}_\xi(\xi, t) = 0.$$

Advection in 2 dimensions

Constant coefficient: $q_t + uq_x + vq_y = 0$

In this case solution for arbitrary initial data is easy:

$$q(x, y, t) = q(x - ut, y - vt, 0).$$

Data simply shifts at constant velocity (u, v) in x - y plane.

Advection in 2 dimensions

Constant coefficient: $q_t + uq_x + vq_y = 0$

In this case solution for **arbitrary** initial data is easy:

$$q(x, y, t) = q(x - ut, y - vt, 0).$$

Data simply shifts at constant velocity (u, v) in x - y plane.

Variable coefficient:

Conservation form: $q_t + (u(x, y, t)q)_x + (v(x, y, t)q)_y = 0$

Advective form (color eqn): $q_t + u(x, y, t)q_x + v(x, y, t)q_y = 0$

Advection in 2 dimensions

Constant coefficient: $q_t + uq_x + vq_y = 0$

In this case solution for **arbitrary** initial data is easy:

$$q(x, y, t) = q(x - ut, y - vt, 0).$$

Data simply shifts at constant velocity (u, v) in x - y plane.

Variable coefficient:

Conservation form: $q_t + (u(x, y, t)q)_x + (v(x, y, t)q)_y = 0$

Advective form (color eqn): $q_t + u(x, y, t)q_x + v(x, y, t)q_y = 0$

Equivalent only if flow is divergence-free (**incompressible**):

$$\nabla \cdot \vec{u} = u_x(x, y, t) + v_y(x, y, t) = 0 \quad \forall t \geq 0.$$

Gas dynamics in 2D

$\rho(x, y, t)$ = mass density

$\rho(x, y, t)u(x, y, t)$ = x -momentum density

$\rho(x, y, t)v(x, y, t)$ = y -momentum density

Gas dynamics in 2D

$\rho(x, y, t)$ = mass density

$\rho(x, y, t)u(x, y, t)$ = x -momentum density

$\rho(x, y, t)v(x, y, t)$ = y -momentum density

If pressure = $P(\rho)$, e.g. **isothermal or isentropic**:

$$\rho_t + (\rho u)_x + (\rho v)_y = 0$$

$$(\rho u)_t + (\rho u^2 + p)_x + (\rho uv)_y = 0$$

$$(\rho v)_t + (\rho uv)_x + (\rho v^2 + p)_y = 0$$

Gas dynamics in 2D

$\rho(x, y, t)$ = mass density

$\rho(x, y, t)u(x, y, t)$ = x -momentum density

$\rho(x, y, t)v(x, y, t)$ = y -momentum density

If pressure = $P(\rho)$, e.g. **isothermal or isentropic**:

$$\rho_t + (\rho u)_x + (\rho v)_y = 0$$

$$(\rho u)_t + (\rho u^2 + p)_x + (\rho uv)_y = 0$$

$$(\rho v)_t + (\rho uv)_x + (\rho v^2 + p)_y = 0$$

For any θ , the matrix $f'(q) \cos \theta + g'(q) \sin \theta$ has **eigenvalues**

$$\check{u} - c, \check{u}, \check{u} + c$$

where $c = \sqrt{P'(\rho)}$ and $\check{u} = u \cos \theta + v \sin \theta$.

Gas dynamics in 2D

$\rho(x, y, t)$ = mass density

$\rho(x, y, t)u(x, y, t)$ = x -momentum density

$\rho(x, y, t)v(x, y, t)$ = y -momentum density

If pressure = $P(\rho)$, e.g. **isothermal** or **isentropic**:

$$\rho_t + (\rho u)_x + (\rho v)_y = 0$$

$$(\rho u)_t + (\rho u^2 + p)_x + (\rho uv)_y = 0$$

$$(\rho v)_t + (\rho uv)_x + (\rho v^2 + p)_y = 0$$


Full Euler equations: 1 more equation for Energy

For any θ , the matrix $f'(q) \cos \theta + g'(q) \sin \theta$ has **eigenvalues**

$$\check{u} - c, \check{u}, \check{u} + c \quad \text{Euler: another wave with } \lambda = \check{u}$$

where $c = \sqrt{P'(\rho)}$ and $\check{u} = u \cos \theta + v \sin \theta$.

Solution of plane wave Riemann problem in 2D



Jump in v from v_l to v_r propagates with the contact discontinuity

Acoustics in 2 dimensions

Linearize about $u = 0$, $v = 0$ and $p =$ perturbation in pressure:

$$p_t + K_0(u_x + v_y) = 0$$

$$\rho_0 u_t + p_x = 0$$

$$\rho_0 v_t + p_y = 0$$

Note: pressure responds to compression or expansion and so p_t is proportional to divergence of velocity.

Second and third equations are $F = ma$.

Acoustics in 2 dimensions

Linearize about $u = 0$, $v = 0$ and $p =$ perturbation in pressure:

$$p_t + K_0(u_x + v_y) = 0$$

$$\rho_0 u_t + p_x = 0$$

$$\rho_0 v_t + p_y = 0$$

Note: pressure responds to compression or expansion and so p_t is proportional to divergence of velocity.

Second and third equations are $F = ma$.

Gives hyperbolic system $q_t + Aq_x + Bq_y = 0$ with

$$q = \begin{bmatrix} p \\ u \\ v \end{bmatrix}, \quad A = \begin{bmatrix} 0 & K_0 & 0 \\ 1/\rho_0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \quad B = \begin{bmatrix} 0 & 0 & K_0 \\ 0 & 0 & 0 \\ 1/\rho_0 & 0 & 0 \end{bmatrix}.$$

Acoustics in 2 dimensions

$$q = \begin{bmatrix} p \\ u \\ v \end{bmatrix}, \quad A = \begin{bmatrix} 0 & K_0 & 0 \\ 1/\rho_0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \quad B = \begin{bmatrix} 0 & 0 & K_0 \\ 0 & 0 & 0 \\ 1/\rho_0 & 0 & 0 \end{bmatrix}.$$

Plane waves:

$$A \cos \theta + B \sin \theta = \begin{bmatrix} 0 & K_0 \cos \theta & K_0 \sin \theta \\ \cos \theta / \rho_0 & 0 & 0 \\ \sin \theta / \rho_0 & 0 & 0 \end{bmatrix}.$$

Acoustics in 2 dimensions

$$q = \begin{bmatrix} p \\ u \\ v \end{bmatrix}, \quad A = \begin{bmatrix} 0 & K_0 & 0 \\ 1/\rho_0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \quad B = \begin{bmatrix} 0 & 0 & K_0 \\ 0 & 0 & 0 \\ 1/\rho_0 & 0 & 0 \end{bmatrix}.$$

Plane waves:

$$A \cos \theta + B \sin \theta = \begin{bmatrix} 0 & K_0 \cos \theta & K_0 \sin \theta \\ \cos \theta / \rho_0 & 0 & 0 \\ \sin \theta / \rho_0 & 0 & 0 \end{bmatrix}.$$

Eigenvalues: $\lambda^1 = -c_0$, $\lambda^2 = 0$, $\lambda^3 = +c_0$

where $c_0 = \sqrt{K_0/\rho_0}$ is **independent** of angle θ .

Isotropic: sound propagates at same speed in any direction.

Acoustics in 2 dimensions

$$q = \begin{bmatrix} p \\ u \\ v \end{bmatrix}, \quad A = \begin{bmatrix} 0 & K_0 & 0 \\ 1/\rho_0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \quad B = \begin{bmatrix} 0 & 0 & K_0 \\ 0 & 0 & 0 \\ 1/\rho_0 & 0 & 0 \end{bmatrix}.$$

Plane waves:

$$A \cos \theta + B \sin \theta = \begin{bmatrix} 0 & K_0 \cos \theta & K_0 \sin \theta \\ \cos \theta / \rho_0 & 0 & 0 \\ \sin \theta / \rho_0 & 0 & 0 \end{bmatrix}.$$

Eigenvalues: $\lambda^1 = -c_0$, $\lambda^2 = 0$, $\lambda^3 = +c_0$

where $c_0 = \sqrt{K_0/\rho_0}$ is **independent** of angle θ .

Isotropic: sound propagates at same speed in any direction.

Note: Zero wave speed for “shear wave” with variation only in velocity in direction $(-\sin \theta, \cos \theta)$.

Diagonalization 2 dimensions

Can we diagonalize system $q_t + Aq_x + Bq_y = 0$ to decouple?

Diagonalization 2 dimensions

Can we diagonalize system $q_t + Aq_x + Bq_y = 0$ to decouple?

Only if A and B have the same eigenvectors!

If $A = R\Lambda R^{-1}$ and $B = RMR^{-1}$, then let $w = R^{-1}q$ and

$$w_t + \Lambda w_x + Mw_y = 0$$

Diagonalization 2 dimensions

Can we diagonalize system $q_t + Aq_x + Bq_y = 0$ to decouple?

Only if A and B have the same eigenvectors!

If $A = R\Lambda R^{-1}$ and $B = RM R^{-1}$, then let $w = R^{-1}q$ and

$$w_t + \Lambda w_x + M w_y = 0$$

In this case, decouples into scalar advection equation
for each component of w :

$$w_t^p + \lambda^p w_x^p + \mu^p w_y^p = 0 \implies w^p(x, y, t) = w^p(x - \lambda^p t, y - \mu^p t, 0).$$

Note: In this case information propagates only in a finite
number of directions (λ^p, μ^p) for $p = 1, \dots, m$.

Diagonalization 2 dimensions

Can we diagonalize system $q_t + Aq_x + Bq_y = 0$ to decouple?

Only if A and B have the same eigenvectors!

If $A = R\Lambda R^{-1}$ and $B = RM R^{-1}$, then let $w = R^{-1}q$ and

$$w_t + \Lambda w_x + M w_y = 0$$

In this case, decouples into scalar advection equation
for each component of w :

$$w_t^p + \lambda^p w_x^p + \mu^p w_y^p = 0 \implies w^p(x, y, t) = w^p(x - \lambda^p t, y - \mu^p t, 0).$$

Note: In this case information propagates only in a finite
number of directions (λ^p, μ^p) for $p = 1, \dots, m$.

This is not true for most coupled systems, e.g. acoustics.

Acoustics in 2 dimensions

$$p_t + K_0(u_x + v_y) = 0$$

$$\rho_0 u_t + p_x = 0$$

$$\rho_0 v_t + p_y = 0$$

$$A = \begin{bmatrix} 0 & K_0 & 0 \\ 1/\rho_0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \quad R^x = \begin{bmatrix} -Z_0 & 0 & Z_0 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$$

Solving $q_t + Aq_x = 0$ gives pressure waves in (p, u) .

Acoustics in 2 dimensions

$$p_t + K_0(u_x + v_y) = 0$$

$$\rho_0 u_t + p_x = 0$$

$$\rho_0 v_t + p_y = 0$$

$$A = \begin{bmatrix} 0 & K_0 & 0 \\ 1/\rho_0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \quad R^x = \begin{bmatrix} -Z_0 & 0 & Z_0 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$$

Solving $q_t + Aq_x = 0$ gives pressure waves in (p, u) .

$$B = \begin{bmatrix} 0 & 0 & K_0 \\ 0 & 0 & 0 \\ 1/\rho_0 & 0 & 0 \end{bmatrix} \quad R^y = \begin{bmatrix} -Z_0 & 0 & Z_0 \\ 0 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix}$$

Solving $q_t + Bq_y = 0$ gives pressure waves in (p, v) .