

In this document, I explore the methods used by GEMs for generating offspring.

Here is the code for generating new individuals.

```
## Code for generating new individuals
pick_individuals <- function(N0, traitmean, traitsd) {
  mu <- log(traitmean^2 / sqrt((traitsd)^2+traitmean^2))
  sigma <- sqrt(log(traitsd^2/traitmean^2 + 1))
  ## record this initial distribution in the output
  return(rlnorm(N0, meanlog=mu, sdlog=sigma))
}
```

I am going to generate an initial population of “parents” with identical traits. To do this, I am using the `pick_individuals` function with  $N0=1000$ ,  $\text{traitmean}=1$ , and  $\text{traitsd}=0.3$ . Each of these parents will give birth to a single offspring. Offspring generation in GEMs is a bit complex. In particular, given a parent with trait  $p$  and a population of parents with mean trait  $\bar{p}$ , and a heritability of  $h^2$ , the value of `traitmean` that will be plugged into `pick_individuals` is  $(1 - h^2)\bar{p} + h^2p$ . At the extremes, if  $h^2 = 1$  then the mean trait in the offspring generation function is identical to their parent trait; if  $h^2 = 0$ , then the mean trait in the offspring generation function is equal to the population-level mean. The value of `traitsd` is calculated using  $\sqrt{1 - (h^2)^2}((1 - h^2) * \sigma_0 + h^2\sigma_p)$ , where  $\sigma_0$  was the initial amount of trait variation and  $\sigma_p$  is the trait variation in the current population. Starting out, then,  $\sigma_0 = \sigma_p$ .

What I am going to do is generating one offspring per parent for 10 different levels of heritability ranging from 0.1 to 1.0. That gives me data upon which I calculate a parent-offspring regression, which is a standard way of estimating heritability in real data: heritability is the slope of the parent-offspring regression. I repeat this process, using the offspring from the first generation as parents for a second generation, and then again performing a parent-offspring regression. Finally, just for completeness, I do this for a third generation, using the offspring of the second generation as parents. The three figures show the estimates of heritability across the generations.




FIGURE 1. Parent-offspring regressions for the first generation.




FIGURE 2. Parent-offspring regressions for the second generation.




FIGURE 3. Parent-offspring regressions for the third generation.

Let's compare that to the case where I'm actually running a GEM to see what happens. In particular, I want to study how the parent-offspring regression changes at different timepoints. You can see that, if you examine the regression over the entire simulation run, the heritability appears to be very high, examined over any smaller interval, it is exactly where you expect it to be!




FIGURE 4. Parent-offspring regressions from a GEM simulation.