Estrogen


73 languages

- Article
- Talk
- Read
- <u>Edit</u>
- View history

Tools

From Wikipedia, the free encyclopedia

This article is about estrogens as hormones. For their use as medications, see <u>Estrogen</u> (medication).





Estrogen or oestrogen (see spelling differences) is a category of sex hormone responsible for the development and regulation of the female reproductive system and secondary sex characteristics. There are three major endogenous estrogens that have estrogenic hormonal activity: estrone (E1), estradiol (E2), and estriol (E3). Estradiol, an estrane, is the most potent and prevalent. Another estrogen called estetrol (E4) is produced only during pregnancy.

Estrogens are synthesized in all vertebrates^[4] and some insects.^[5] Their presence in both vertebrates and insects suggests that estrogenic sex hormones have an ancient evolutionary history. Quantitatively, estrogens circulate at lower levels than <u>androgens</u> in both men and women.^[6] While estrogen levels are significantly lower in males than in females, estrogens nevertheless have important physiological roles in males.^[7]

Like all <u>steroid hormones</u>, estrogens readily <u>diffuse</u> across the <u>cell membrane</u>. Once inside the cell, they bind to and activate <u>estrogen receptors</u> (ERs) which in turn <u>modulate</u> the <u>expression</u> of many <u>genes</u>. Additionally, estrogens bind to and activate rapid-signaling <u>membrane estrogen receptors</u> (mERs), such as <u>GPER</u> (GPR30).

In addition to their role as natural hormones, estrogens are used as <u>medications</u>, for instance in <u>menopausal hormone therapy</u>, <u>hormonal birth control</u> and <u>feminizing</u> hormone therapy for transgender women and nonbinary people.

Synthetic and natural estrogens have been found in the environment and are referred to as <u>xenoestrogens</u>. Estrogens are among the wide range of endocrine-disrupting compounds (EDCs) and can cause health issues and reproductive disfunction in both wildlife and humans.[12][13]

Types and examples[edit]



Structures of major endogenous estrogens



Note the <u>hydroxyl</u> (–OH) <u>groups</u>: estrone (E1) has one, estradiol (E2) has two, estriol (E3) has three, and estetrol (E4) has four.

Estriol (E3)


Estetrol (E4)

The four major naturally occurring estrogens in women are <u>estrone</u> (E1), <u>estradiol</u> (E2), <u>estriol</u> (E3), and <u>estetrol</u> (E4). Estradiol (E2) is the predominant estrogen during reproductive years both in terms of absolute serum levels as well as in terms of estrogenic activity. During <u>menopause</u>, estrone is the predominant circulating estrogen and during pregnancy estriol is the predominant circulating estrogen in terms of serum levels. Given by <u>subcutaneous injection</u> in mice, estradiol is about 10-fold more potent than estrone and about 100-fold more potent than estriol. Thus, estradiol is the most important estrogen in non-pregnant females who are between the <u>menarche</u> and menopause stages of life. However, during <u>pregnancy</u> this role shifts to estriol, and in postmenopausal women estrone becomes the primary form of estrogen in the body. Another type of estrogen called <u>estetrol</u> (E4) is produced only during pregnancy. All of the different forms of estrogen are synthesized from <u>androgens</u>, specifically <u>testosterone</u> and <u>androstenedione</u>, by the <u>enzyme aromatase</u>.

Minor endogenous estrogens, the biosyntheses of which do not involve <u>aromatase</u>, include <u>27-hydroxycholesterol</u>, <u>dehydroepiandrosterone</u> (DHEA), <u>7-oxo-DHEA</u>, <u>7α-hydroxy-DHEA</u>, <u>16α-hydroxy-DHEA</u>, <u>7β-hydroxyepiandrosterone</u>, androstenedione (A4), <u>androstenediol</u> (A5), <u>3α-</u>

androstanediol, and <u>3β-androstanediol</u>. Some estrogen metabolites, such as the <u>catechol estrogens 2-hydroxyestradiol</u>, <u>2-hydroxyestrone</u>, <u>4-hydroxyestradiol</u>, and <u>4-hydroxyestrone</u>, as well as <u>16α-hydroxyestrone</u>, are also estrogens with varying degrees of activity. The biological importance of these minor estrogens is not entirely clear.

Biological function[edit]



Reference ranges for the blood content of estradiol, the primary type of estrogen, during the menstrual cycle.[18]

The actions of estrogen are mediated by the <u>estrogen receptor</u> (ER), a dimeric nuclear protein that binds to DNA and controls <u>gene expression</u>. Like other steroid hormones, estrogen enters passively into the cell where it binds to and activates the estrogen receptor. The estrogen:ER complex binds to specific DNA sequences called a <u>hormone response element</u> to activate the transcription of target genes (in a study using an estrogen-dependent breast cancer cell line as model, 89 such genes were identified). Since estrogen enters all cells, its actions are dependent on the presence of the ER in the cell. The ER is expressed in specific tissues including the ovary, uterus and breast. The metabolic effects of estrogen in postmenopausal women have been linked to the genetic polymorphism of the ER.

While estrogens are present in both <u>men</u> and <u>women</u>, they are usually present at significantly higher levels in women of reproductive age. They promote the development of female <u>secondary sexual characteristics</u>, such as <u>breasts</u>, and are also involved in the thickening of the <u>endometrium</u> and other aspects of regulating the menstrual cycle. In males, estrogen regulates certain functions of the <u>reproductive system</u> important to the maturation of sperm^{[21][22][23]} and may be necessary for a healthy libido.^[24]

```
• <u>v</u>
• <u>t</u>
• <u>e</u>
```

Affinities of estrogen receptor ligands for the ER α and ER β show

<u>v</u> <u>t</u>

Relative affinities of estrogens for steroid hormone receptors and blood proteins show

```
• <u>v</u>
• <u>t</u>
• <u>e</u>
```

Affinities and estrogenic potencies of estrogen esters and ethers at the estrogen receptors show

<u>v</u> <u>t</u> e

Selected biological properties of endogenous estrogens in rats show

Overview of actions[edit]



This section is in <u>list</u> format but may read better as <u>prose</u>. You can help by converting this section, if appropriate. Editing help is available. (*October 2019*)

- Structural
 - Anabolic: Increases <u>muscle mass</u> and strength, speed of muscle regeneration, and <u>bone density</u>, increased sensitivity to exercise, protection against muscle damage, stronger <u>collagen</u> synthesis, increases the collagen content of <u>connective tissues</u>, <u>tendons</u>, and <u>ligaments</u>, but also decreases stiffness of <u>tendons</u> and <u>ligaments</u> (especially during <u>menstruation</u>). Decreased stiffness of tendons gives women much lower predisposition to muscle strains but soft ligaments are much more prone to injuries (<u>ACL</u> tears are 2-8x more common among women than men).
 - Anti-inflammatory properties
 - o Mediate formation of female <u>secondary sex characteristics</u>
 - Accelerate metabolism
 - Increased <u>fat storage</u> in some body parts such as breasts, buttocks, and legs but decreased abdominal and <u>visceral fat</u> (androgenic obesity). [29][30][31] <u>Estradiol</u> also regulates energy expenditure, body weight <u>homeostasis</u>, and seems to have much stronger anti-obesity effects than testosterone in general. [32]
- Women tend to have lower base strength but on average have about the same increases of muscle mass in responses to resistance training as men and far faster relative increases in strength.[33][34]
 - Stimulate endometrial growth
 - o Increase uterine growth
 - Increase vaginal lubrication
 - Thicken the vaginal wall
 - Maintenance of vessel and skin
 - Reduce bone resorption, increase bone formation
- Protein synthesis
 - Increase hepatic production of binding proteins
- Coagulation

0

- o Increase circulating level of <u>factors 2, 7, 9, 10, plasminogen</u>
- Decrease antithrombin III
- o Increase <u>platelet</u> adhesiveness
- o Increase <u>vWF</u> (estrogen -> <u>Angiotensin II -> Vasopressin</u>)

- o Increase PAI-1 and PAI-2 also through Angiotensin II
- Lipid

0

0

0

- o Increase HDL, triglyceride
 - Decrease LDL, fat deposition
- Fluid balance
 - Salt (sodium) and water retention
 - o Increase cortisol, SHBG
- Gastrointestinal tract
 - Reduce bowel motility
 - Increase cholesterol in bile
- Melanin
 - Increase pheomelanin, reduce eumelanin
- Cancer
 - Support hormone-sensitive breast cancers (see section below)
- Lung function
 - Promotes lung function by supporting <u>alveoli</u> (in rodents but probably in humans).
- <u>Uterus</u> lining
 - Estrogen together with <u>progesterone</u> promotes and maintains the uterus lining in preparation for implantation of fertilized egg and maintenance of uterus function during gestation period, also upregulates <u>oxytocin</u> receptor in myometrium
- Ovulation
 - Surge in estrogen level induces the release of <u>luteinizing hormone</u>, which then triggers ovulation by releasing the egg from the <u>Graafian follicle</u> in the <u>ovary</u>.
- Sexual behavior
 - Estrogen is required for female mammals to engage in <u>lordosis</u> <u>behavior</u> during <u>estrus</u> (when animals are "in heat"). This behavior is required for sexual receptivity in these mammals and is regulated by the <u>ventromedial nucleus</u> of the hypothalamus. [38]
 - Sex drive is dependent on androgen levels only in the presence of estrogen, but without estrogen, free testosterone level actually decreases sexual desire (instead of increases sex drive), as demonstrated for those women who have hypoactive sexual desire disorder, and the sexual desire in these women can be restored by administration of estrogen (using oral contraceptive).[40]

Female pubertal development[edit]

Estrogens are responsible for the development of female <u>secondary sexual</u> <u>characteristics</u> during <u>puberty</u>, including <u>breast development</u>, widening of the <u>hips</u>, and female <u>fat distribution</u>. Conversely, <u>androgens</u> are responsible for <u>pubic</u> and <u>body</u> <u>hair growth</u>, as well as <u>acne</u> and <u>axillary odor</u>.

Breast development[edit]

See also: Breast development § Biochemistry

Estrogen, in conjunction with <u>growth hormone</u> (GH) and its secretory product <u>insulin-like</u> <u>growth factor 1</u> (IGF-1), is critical in mediating breast development during <u>puberty</u>, as well as breast maturation during <u>pregnancy</u> in preparation of <u>lactation</u> and <u>breastfeeding</u>. [41][42] Estrogen is primarily and directly responsible for

inducing the ductal component of breast development, [43][44][45] as well as for causing fat deposition and connective tissue growth. [43][44] It is also indirectly involved in the lobuloalveolar component, by increasing progesterone receptor expression in the breasts [43][45][46] and by inducing the secretion of prolactin. [47][48] Allowed for by estrogen, progesterone and prolactin work together to complete lobuloalveolar development during pregnancy. [44][49]

<u>Androgens</u> such as testosterone powerfully oppose estrogen action in the breasts, such as by reducing <u>estrogen receptor</u> expression in them. [50][51]

Female reproductive system[edit]

Estrogens are responsible for maturation and maintenance of the <u>vagina</u> and <u>uterus</u>, and are also involved in <u>ovarian</u> function, such as maturation of <u>ovarian follicles</u>. In addition, estrogens play an important role in regulation of <u>gonadotropin secretion</u>. For these reasons, estrogens are required for female fertility.

Neuroprotection and DNA repair[edit]

Estrogen regulated <u>DNA repair</u> mechanisms in the <u>brain</u> have neuroprotective effects. Estrogen regulates the <u>transcription</u> of DNA <u>base excision repair</u> genes as well as the translocation of the base excision repair enzymes between different subcellular compartments.

Brain and behavior[edit]

Sex drive[edit]

See also: <u>Sexual motivation and hormones</u>

Estrogens are involved in <u>libido</u> (sex drive) in both women and men.

Cognition[edit]

<u>Verbal memory</u> scores are frequently used as one measure of higher level <u>cognition</u>. These scores vary in direct proportion to estrogen levels throughout the menstrual cycle, pregnancy, and menopause. Furthermore, estrogens when administered shortly after natural or surgical menopause prevents decreases in verbal memory. In contrast, estrogens have little effect on verbal memory if first administered years after menopause. [53] Estrogens also have positive influences on other measures of cognitive function. [54] However the effect of estrogens on cognition is not uniformly favorable and is dependent on the timing of the dose and the type of cognitive skill being measured. [55]

The protective effects of estrogens on cognition may be mediated by estrogen's antiinflammatory effects in the brain. Studies have also shown that the Met allele gene and level of estrogen mediates the efficiency of <u>prefrontal cortex</u> dependent working memory tasks. Researchers have urged for further research to illuminate the role of estrogen and its potential for improvement on cognitive function.

Mental health[edit]

Estrogen is considered to play a significant role in women's <u>mental health</u>. Sudden estrogen withdrawal, fluctuating estrogen, and <u>periods</u> of sustained low estrogen levels

correlate with a significant lowering of mood. Clinical recovery from <u>postpartum</u>, <u>perimenopause</u>, and <u>postmenopause</u> depression has been shown to be effective after levels of estrogen were stabilized and/or restored. Menstrual <u>exacerbation (including menstrual psychosis)</u> is typically triggered by low estrogen levels, and is often mistaken for premenstrual dysphoric disorder.

Compulsions in male lab mice, such as those in obsessive-compulsive disorder (OCD), may be caused by low estrogen levels. When estrogen levels were raised through the increased activity of the enzyme <u>aromatase</u> in male lab mice, OCD rituals were dramatically decreased. <u>Hypothalamic</u> protein levels in the gene <u>COMT</u> are enhanced by increasing estrogen levels which are believed to return mice that displayed OCD rituals to normal activity. Aromatase deficiency is ultimately suspected which is involved in the synthesis of estrogen in humans and has therapeutic implications in humans having obsessive-compulsive disorder. [65]

Local application of estrogen in the rat hippocampus has been shown to inhibit the reuptake of <u>serotonin</u>. Contrarily, local application of estrogen has been shown to block the ability of <u>fluvoxamine</u> to slow serotonin clearance, suggesting that the same pathways which are involved in SSRI efficacy may also be affected by components of local estrogen signaling pathways. [66]

Parenthood[edit]

Studies have also found that fathers had lower levels of cortisol and testosterone but higher levels of estrogen (estradiol) than did non-fathers. [67]

Binge eating[edit]

Estrogen may play a role in suppressing <u>binge eating</u>. Hormone replacement therapy using estrogen may be a possible treatment for binge eating behaviors in females. Estrogen replacement has been shown to suppress binge eating behaviors in female mice. The mechanism by which estrogen replacement inhibits binge-like eating involves the replacement of <u>serotonin</u> (5-HT) neurons. Women exhibiting binge eating behaviors are found to have increased brain uptake of neuron 5-HT, and therefore less of the neurotransmitter serotonin in the cerebrospinal fluid. Estrogen works to activate 5-HT neurons, leading to suppression of binge like eating behaviors.

It is also suggested that there is an interaction between hormone levels and eating at different points in the female <u>menstrual cycle</u>. Research has predicted increased emotional eating during hormonal flux, which is characterized by high <u>progesterone</u> and <u>estradiol</u> levels that occur during the mid-<u>luteal phase</u>. It is hypothesized that these changes occur due to brain changes across the menstrual cycle that are likely a genomic effect of hormones. These effects produce menstrual cycle changes, which result in hormone release leading to behavioral changes, notably binge and emotional eating. These occur especially prominently among women who are genetically vulnerable to binge eating phenotypes.^[70]

Binge eating is associated with decreased estradiol and increased progesterone. [71] Klump et al. [72] Progesterone may moderate the effects of low estradiol (such as during dysregulated eating behavior), but that this may only be true in women

who have had clinically diagnosed binge episodes (BEs). Dysregulated eating is more strongly associated with such ovarian hormones in women with BEs than in women without BEs.[72]

The implantation of 17β-estradiol pellets in ovariectomized mice significantly reduced binge eating behaviors and injections of GLP-1 in ovariectomized mice decreased binge-eating behaviors. [68]

The associations between binge eating, menstrual-cycle phase and ovarian hormones correlated. [71][73][74]

Masculinization in rodents[edit]

In rodents, estrogens (which are locally aromatized from androgens in the brain) play an important role in psychosexual differentiation, for example, by masculinizing territorial behavior;^[75] the same is not true in humans.^[76] In humans, the masculinizing effects of prenatal androgens on behavior (and other tissues, with the possible exception of effects on bone) appear to act exclusively through the androgen receptor.^[77] Consequently, the utility of rodent models for studying human psychosexual differentiation has been questioned.^[78]

Skeletal system[edit]

Estrogens are responsible for both the pubertal growth spurt, which causes an acceleration in linear growth, and <u>epiphyseal closure</u>, which limits <u>height</u> and <u>limb</u> length, in both females and males. In addition, estrogens are responsible for bone maturation and maintenance of <u>bone mineral density</u> throughout life. Due to hypoestrogenism, the risk of <u>osteoporosis</u> increases during <u>menopause</u>.

Cardiovascular system[edit]

Women are less impacted by heart disease due to vasculo-protective action of estrogen which helps in preventing atherosclerosis. It also helps in maintaining the delicate balance between fighting infections and protecting arteries from damage thus lowering the risk of cardiovascular disease. During pregnancy, high levels of estrogens increase coagulation and the risk of venous thromboembolism. Estrogen has been shown to upregulate the peptide hormone adropin. Estrogen has been

• <u>v</u>
• <u>t</u>
• e

Absolute and relative incidence of venous thromboembolism (VTE) during pregnancy and the postpartum period show

Immune system[edit]

The effect of estrogen one the <u>immune system</u> is in general described as <u>Th2</u> favoring, rather than suppressive, as is the case of the effect of male sex hormone - testosterone. Indeed women respond better to <u>vaccines</u>, <u>infections</u> and are generally less likely to develop <u>cancer</u>, the tradeoff of this is that they are more likely do develop an <u>autoimmune disease</u>. The <u>Th2</u> shift manifests itself in a decrease of cellular

immunity and increase in humoral immunity (antibody production) shifts it from cellular to humoral by downregulating cell-mediated immunity and enhancing Th2 immune response by stimulating IL-4 production and Th2 differentiation. Type 1 and type 17 immune responses are downregulated, likely to be at least partially due to IL-4, which inhibits Th1. Effect of estrogen on different immune cells' cell types is in line with its Th2 bias. Activity of basophils, eosinophils, M2 macrophages and is enhanced, whereas activity of NK cells is downregulated. Conventional dendritic cells are biased towards Th2 under the influence of estrogen, whereas plasmacytoid dendritic cells, key players in antiviral defence, have increased IFN-g secretion. Estrogen also influences B cells by increasing their survival, proliferation, differentiation and function, which corresponds with higher antibody and B cell count generally detected in women.

On a molecular level estrogen induces the above mentioned effects on cell via acting on intracellular <u>receptors</u> termed ER α and ER β, which upon ligation form either homo or heterodimers. The genetic and nongenetic targets of the receptors differ between homo and heterodimers. Ligation of these receptors allows them to translocate to the <u>nucleus</u> and act as <u>transcription factors</u> either by binding estrogen response elements (ERE) on <u>DNA</u> or binding DNA together with other transcriptional factors e.g. <u>Nf-kB</u> or <u>AP-1</u>, both of which result in <u>RNA polymerase</u> recruitment and further <u>chromatin remodelation</u>. A non-transcriptional response to oestrogen stimulation was also documented (termed membrane-initiated steroid signalling, MISS). This pathway stimulates the ERK and PI3K/AKT pathways, which are known to increase cellular proliferation and affect chromatin remodelation.

Associated conditions[edit]


Researchers have implicated estrogens in various <u>estrogen-dependent conditions</u>, such as ER-positive <u>breast cancer</u>, as well as a number of <u>genetic conditions</u> involving estrogen signaling or metabolism, such as <u>estrogen insensitivity syndrome</u>, <u>aromatase deficiency</u>, and <u>aromatase excess syndrome</u>.

High estrogen can amplify stress-hormone responses in stressful situations.[88]

Biochemistry[edit]

See also: Estradiol § Biochemistry

Biosynthesis[edit]



Estrogens, in females, are produced primarily by the <u>ovaries</u>, and during pregnancy, the <u>placenta</u>. [90] Follicle-stimulating hormone (FSH) stimulates the ovarian production of estrogens by the <u>granulosa cells</u> of the <u>ovarian follicles</u> and <u>corpora lutea</u>. Some estrogens are also produced in smaller amounts by other tissues such as the <u>liver</u>, <u>pancreas</u>, <u>bone</u>, <u>adrenal glands</u>, <u>skin</u>, <u>brain</u>, <u>adipose tissue</u>, [91] and the <u>breasts</u>. [92] These secondary sources of estrogens are especially important in postmenopausal women. [93] The pathway of estrogen biosynthesis in extragonadal tissues is different. These tissues are not able to synthesize C19 steroids, and therefore depend on C19 supplies from other tissues [93] and the level of aromatase. [94]

In females, synthesis of estrogens starts in theca internation cells in the ovary, by the synthesis of androstenedione from cholesterol. Androstenedione is a substance of weak androgenic activity which serves predominantly as a precursor for more potent androgens such as testosterone as well as estrogen. This compound crosses the basal membrane into the surrounding granulosa cells, where it is converted either immediately into estrone, or into testosterone and then estradiol in an additional step. The conversion of androstenedione to testosterone is catalyzed by 17β -hydroxysteroid dehydrogenase (17 β -HSD), whereas the conversion of androstenedione and testosterone into estrone and estradiol, respectively is catalyzed by aromatase, enzymes which are both expressed in granulosa cells. In contrast, granulosa cells lack 17α -hydroxylase and 17,20-lyase, whereas theca cells express these enzymes and 17β -HSD but lack aromatase. Hence, both granulosa and theca cells are essential for the production of estrogen in the ovaries.

Estrogen levels vary through the <u>menstrual cycle</u>, with levels highest near the end of the <u>follicular phase</u> just before <u>ovulation</u>.

Note that in males, estrogen is also produced by the <u>Sertoli cells</u> when FSH binds to their FSH receptors.

```
• <u>v</u>
• <u>t</u>
• <u>e</u>
```

Production rates, secretion rates, clearance rates, and blood levels of major sex hormones show

Distribution[edit]

Estrogens are <u>plasma protein bound</u> to <u>albumin</u> and/or <u>sex hormone-binding globulin</u> in the circulation.

Metabolism[edit]

See also: Estradiol § Metabolism, and Estradiol (medication) § Metabolism

Estrogens are <u>metabolized</u> via <u>hydroxylation</u> by <u>cytochrome P450 enzymes</u> such as <u>CYP1A1</u> and <u>CYP3A4</u> and via <u>conjugation</u> by <u>estrogen sulfotransferases</u> (<u>sulfation</u>) and <u>UDP-glucuronyltransferases</u> (<u>glucuronidation</u>). In addition, estradiol is <u>dehydrogenated</u> by <u>17 β -hydroxysteroid dehydrogenase</u> into the much less potent estrogen estrone. These reactions occur primarily in the <u>liver</u>, but also in other <u>tissues</u>.

• <u>v</u> • <u>t</u>

Estrogen metabolism in humans

Estradiol

Estrone sulfate

Estrone glucuronide

2-Hydroxyestrone

Estrone

4-Hydroxyestrone

2-Methoxyestrone

 $\underline{16\alpha\text{-Hydroxyestrone}}$

4-Methoxyestrone

17-Epiestriol

Estriol

16-Epiestriol

EST

STS

UGT1A3 UGT1A9

CYP450

CYP450

COMT

CYP450

COMT

unidentified

17β-HSD

unidentified



Description: The metabolic pathways involved in the metabolism of estradiol and other natural estrogens (e.g., estrone, estriol) in humans. In addition to the metabolic transformations shown in the diagram, conjugation (e.g., sulfation and glucuronidation) occurs in the case of estradiol and metabolites of estradiol that have one or more available hydroxyl (-OH) groups. Sources: See template page.

Excretion[edit]

Estrogens are <u>excreted</u> primarily by the <u>kidneys</u> as <u>conjugates</u> via the <u>urine</u>.

Medical use[edit]

Main article: Estrogen (medication)

Estrogens are used as <u>medications</u>, mainly in <u>hormonal contraception</u>, <u>hormone</u> <u>replacement therapy</u>, and to treat gender dysphoria in <u>transgender women</u> and other <u>transfeminine individuals</u> as part of feminizing hormone therapy. [96]

Chemistry[edit]

See also: <u>List of estrogens</u>

The estrogen steroid hormones are estrane steroids.

History[edit]

See also: Estradiol § History, Estrone § History, and Estrogen (medication) § History

In 1929, Adolf Butenandt and Edward Adelbert Doisy independently isolated and purified estrone, the first estrogen to be discovered. Then, estriol and estradiol were discovered in 1930 and 1933, respectively. Shortly following their discovery, estrogens, both natural and synthetic, were introduced for medical use. Examples include estriol glucuronide (Emmenin, Progynon), estradiol benzoate, conjugated estrogens (Premarin), diethylstilbestrol, and ethinylestradiol.

The word estrogen derives from Ancient Greek. It is derived from "oestros" [18] (a periodic state of sexual activity in female mammals), and genos (generating). [18] It was first published in the early 1920s and referenced as "oestrin". [19] With the years, American English adapted the spelling of estrogen to fit with its phonetic pronunciation. Nevertheless, both estrogen and oestrogen are used nowadays, yet some still wish to maintain its original spelling as it reflects the origin of the word.

Society and culture[edit]

Etymology[edit]

The name *estrogen* is derived from the <u>Greek</u> οἶστρος (*oîstros*), literally meaning "verve" or "inspiration" but figuratively sexual passion or desire, and the suffix *-gen*, meaning "producer of".

Environment[edit]

A range of synthetic and natural substances that possess estrogenic activity have been identified in the <u>environment</u> and are referred to <u>xenoestrogens</u>.[101]

- Synthetic substances such as <u>bisphenol A</u> as well as <u>metalloestrogens</u> (e.g., <u>cadmium</u>).
- Plant products with estrogenic activity are called phytoestrogens (e.g., coumestrol, daidzein, genistein, miroestrol).
- Those produced by fungi are known as mycoestrogens (e.g., zearalenone).

Estrogens are among the wide range of <u>endocrine-disrupting compounds</u> (EDCs) because they have high estrogenic potency. When an EDC makes its way into the environment, it may cause male reproductive dysfunction to wildlife and humans. The estrogen excreted from farm animals makes its way into fresh water systems. During the germination period of reproduction the fish are exposed to low levels of estrogen which may cause reproductive dysfunction to male fish.

Cosmetics[edit]

Some hair <u>shampoos</u> on the market include estrogens and placental extracts; others contain <u>phytoestrogens</u>. In 1998, there were case reports of four prepubescent African-American girls developing breasts after exposure to these shampoos. In 1993, the FDA determined that not all <u>over-the-counter</u> topically applied hormone-containing drug products for human use are <u>generally recognized as safe and effective</u> and are misbranded. An accompanying proposed rule deals with cosmetics, concluding that any use of natural estrogens in a cosmetic product makes the product an unapproved new drug and that any cosmetic using the term "hormone" in the text of its labeling or in its ingredient statement makes an implied drug claim, subjecting such a product to regulatory action.

In addition to being considered misbranded drugs, products claiming to contain placental extract may also be deemed to be misbranded cosmetics if the extract has been prepared from placentas from which the hormones and other biologically active substances have been removed and the extracted substance consists principally of protein. The FDA recommends that this substance be identified by a name other than "placental extract" and describing its composition more accurately because consumers associate the name "placental extract" with a therapeutic use of some biological activity. [106]

See also[edit]

- <u>List of steroid abbreviations</u>
- Breastfeeding and fertility

References[edit]

- 1. ^ Jump up to: ^{a b c} Huether SE, McCance KL (2019). <u>Understanding Pathophysiology</u>. Elsevier Health Sciences. p. 767. <u>ISBN 978-0-32-367281-8</u>. Estrogen is a generic term for any of three similar hormones derived from cholesterol: estradiol, estrone, and estriol.
- 2. <u>^</u> Satoskar RS, Rege N, Bhandarkar SD (2017). <u>Pharmacology and Pharmacotherapeutics</u>. Elsevier Health Sciences. p. 943. <u>ISBN 978-8-13-124941-3</u>. The natural estrogens are steroids. However, typical estrogenic activity is also shown by chemicals which are not steroids. Hence, the term 'estrogen' is used as a generic term to describe all the compounds having estrogenic activity.
- 3. ___ Delgado BJ, Lopez-Ojeda W (20 December 2021). <u>"Estrogen"</u>. StatPearls [Internet]. StatPearls Publishing. <u>PMID 30855848</u>. Estrogen is a steroid hormone associated with the female reproductive organs and is responsible for the development of female sexual characteristics. Estrogen is often referred to as estrone, estradiol, and estriol. ... Synthetic estrogen is also available for clinical use, designed to increase absorption and effectiveness by altering the estrogen chemical structure for topical or oral administration. Synthetic steroid estrogens include ethinyl estradiol, estradiol valerate, estropipate, conjugate esterified estrogen, and quinestrol.
- 4. A Ryan KJ (August 1982). "Biochemistry of aromatase: significance to female reproductive physiology". Cancer Research. **42** (8 Suppl): 3342s–3344s. PMID 7083198.
- 5. <u>^</u> Mechoulam R, Brueggemeier RW, Denlinger DL (September 2005). "Estrogens in insects". Cellular and Molecular Life Sciences. **40** (9): 942–944. doi:10.1007/BF01946450. S2CID 31950471.
- 6. <u>^ Burger HG (April 2002). "Androgen production in women"</u>. Fertility and Sterility. **77** (Suppl 4): S3–S5. doi:10.1016/S0015-0282(02)02985-0. PMID 12007895.
- 7. <u>^</u> Lombardi G, Zarrilli S, Colao A, Paesano L, Di Somma C, Rossi F, De Rosa M (June 2001). "Estrogens and health in males". Molecular and Cellular Endocrinology. **178** (1–2): 51–55. <u>doi:10.1016/S0303-7207(01)00420-8</u>. <u>PMID</u> <u>11403894</u>. <u>S2CID</u> <u>36834775</u>.

- 8. <u>^ Whitehead SA, Nussey S (2001). Endocrinology: an integrated approach.</u> Oxford: BIOS: Taylor & Francis. ISBN 978-1-85996-252-7. PMID 20821847.
- 9. <u>^</u> Soltysik K, Czekaj P (April 2013). "Membrane estrogen receptors is it an alternative way of estrogen action?". Journal of Physiology and Pharmacology. **64** (2): 129–142. PMID 23756388.
- 10. <u>^ Micevych PE, Kelly MJ (2012). "Membrane estrogen receptor regulation of hypothalamic function"</u>. Neuroendocrinology. **96** (2): 103–110. doi:10.1159/000338400. PMC 3496782. PMID 22538318.
- 11. <u>^</u> Prossnitz ER, Arterburn JB, Sklar LA (February 2007). <u>"GPR30: A G protein-coupled receptor for estrogen"</u>. Molecular and Cellular Endocrinology. 265–266: 138–142. doi:10.1016/j.mce.2006.12.010. PMC 1847610. PMID 17222505.
- 12. ^ Jump up to: ^a ½ Wang S, Huang W, Fang G, Zhang Y, Qiao H (2008). "Analysis of steroidal estrogen residues in food and environmental samples". International Journal of Environmental Analytical Chemistry. **88** (1): 1–25. doi:10.1080/03067310701597293. S2CID 93975613.
- 13. ^ Jump up to: ^a ½ Korach KD (1998). <u>Reproductive and developmental toxicology</u>. New York: Marcel Dekker. <u>ISBN 0-585-15807-X</u>. <u>OCLC 44957536</u>.
- 14. <u>^</u> A. Labhart (6 December 2012). <u>Clinical Endocrinology: Theory and Practice</u>. Springer Science & Business Media. pp. 548–. ISBN 978-3-642-96158-8.
- 15. <u>^</u> Baker ME (March 2013). "What are the physiological estrogens?". Steroids. **78** (3): 337–340. doi:10.1016/j.steroids.2012.12.011. PMID 23313336. S2CID 11803629.
- 16. <u>^</u> Miller KK, Al-Rayyan N, Ivanova MM, Mattingly KA, Ripp SL, Klinge CM, Prough RA (January 2013). "DHEA metabolites activate estrogen receptors alpha and beta". Steroids. **78** (1): 15–25. doi:10.1016/j.steroids.2012.10.002. PMC 3529809. PMID 23123738.
- 17. <u>^</u> Bhavnani BR, Nisker JA, Martin J, Aletebi F, Watson L, Milne JK (2000). "Comparison of pharmacokinetics of a conjugated equine estrogen preparation (premarin) and a synthetic mixture of estrogens (C.E.S.) in postmenopausal women". Journal of the Society for Gynecologic Investigation. **7** (3): 175–183. doi:10.1016/s1071-5576(00)00049-6. PMID 10865186.
- 18. <u>^ Häggström M (2014). "Reference ranges for estradiol, progesterone, luteinizing hormone and follicle-stimulating hormone during the menstrual cycle"</u>. WikiJournal of Medicine. **1** (1). doi:10.15347/wjm/2014.001. ISSN 2002-4436.
- 19. <u>^</u> Lin CY, Ström A, Vega VB, Kong SL, Yeo AL, Thomsen JS, et al. (2004). <u>"Discovery of estrogen receptor alpha target genes and response elements in breast tumor cells"</u>. Genome Biology. **5** (9): R66. doi:10.1186/gb-2004-5-9-r66. PMC 522873. PMID 15345050.
- 20. __ Darabi M, Ani M, Panjehpour M, Rabbani M, Movahedian A, Zarean E (2011). "Effect of estrogen receptor β A1730G polymorphism on ABCA1 gene expression response to postmenopausal hormone replacement therapy". Genetic Testing and Molecular Biomarkers. 15 (1–2): 11–15. doi:10.1089/gtmb.2010.0106. PMID 21117950.
- 21. <u>^</u> Raloff J (6 December 1997). "Science News Online (12/6/97): Estrogen's Emerging Manly Alter Ego". Science News. Retrieved 4 March 2008.
- 22. <u>^</u> Hess RA, Bunick D, Lee KH, Bahr J, Taylor JA, Korach KS, Lubahn DB (December 1997). <u>"A role for oestrogens in the male reproductive system"</u>. Nature. **390** (6659): 509–512. <u>Bibcode:1997Natur.390..509H. doi:10.1038/37352. PMC 5719867. PMID 9393999</u>.
- 23. <u>^ "Estrogen Linked To Sperm Count, Male Fertility"</u>. Science Blog. Archived from the original on 7 May 2007. Retrieved 4 March 2008.
- 24. <u>^</u> Hill RA, Pompolo S, Jones ME, Simpson ER, Boon WC (December 2004). "Estrogen deficiency leads to apoptosis in dopaminergic neurons in the medial preoptic area and arcuate nucleus of male mice". Molecular and Cellular Neurosciences. **27** (4): 466–476. doi:10.1016/j.mcn.2004.04.012. PMID 15555924. S2CID 25280077.
- 25. <u>^ Chidi-Ogbolu N, Baar K (2018). "Effect of Estrogen on Musculoskeletal Performance and Injury Risk"</u>. Frontiers in Physiology. **9**: 1834. doi:10.3389/fphys.2018.01834. PMC 6341375. PMID 30697162.
- 26. <u>^</u> Lowe DA, Baltgalvis KA, Greising SM (April 2010). <u>"Mechanisms behind estrogen's beneficial effect on muscle strength in females"</u>. Exercise and Sport Sciences Reviews. **38** (2): 61–67. doi:10.1097/JES.0b013e3181d496bc. PMC 2873087. PMID 20335737.
- 27. <u>^</u> Max SR (December 1984). "Androgen-estrogen synergy in rat levator ani muscle: glucose-6-phosphate dehydrogenase". Molecular and Cellular Endocrinology. **38** (2–3): 103–107. doi:10.1016/0303-7207(84)90108-4. PMID 6510548. S2CID 24198956.

- 28. _^ Koot RW, Amelink GJ, Blankenstein MA, Bär PR (1991). "Tamoxifen and oestrogen both protect the rat muscle against physiological damage". The Journal of Steroid Biochemistry and Molecular Biology. 40 (4–6): 689–695. doi:10.1016/0960-0760(91)90292-d. PMID 1958566. S2CID 44446541.
- 29. <u>^</u> Frank AP, de Souza Santos R, Palmer BF, Clegg DJ (October 2019). <u>"Determinants of body fat distribution in humans may provide insight about obesity-related health risks"</u>. Journal of Lipid Research. **60** (10): 1710–1719. <u>doi:10.1194/jlr.R086975</u>. <u>PMC 6795075</u>. <u>PMID 30097511</u>.
- Brown LM, Gent L, Davis K, Clegg DJ (September 2010). "Metabolic impact of sex hormones on obesity". Brain Research. 1350: 77–85. doi:10.1016/j.brainres.2010.04.056. PMC 2924463. PMID 20441773.
- 31. _____ Janssen I, Powell LH, Kazlauskaite R, Dugan SA (March 2010). <u>"Testosterone and visceral fat in midlife women: the Study of Women's Health Across the Nation (SWAN) fat patterning study"</u>. Obesity. **18** (3): 604–610. doi:10.1038/oby.2009.251. PMC 2866448. PMID 19696765.
- 32. <u>^</u> Rubinow KB (2017). <u>"Estrogens and Body Weight Regulation in Men"</u>. Advances in Experimental Medicine and Biology. **1043**: 285–313. <u>doi:10.1007/978-3-319-70178-3_14</u>. <u>ISBN 978-3-319-70177-6</u>. PMC 5835337. PMID 29224100.
- 33. <u>^</u> O'Hagan FT, Sale DG, MacDougall JD, Garner SH (July 1995). "Response to resistance training in young women and men". International Journal of Sports Medicine. **16** (5): 314–321. doi:10.1055/s-2007-973012. PMID 7558529.
- 34. A Roth SM, Ivey FM, Martel GF, Lemmer JT, Hurlbut DE, Siegel EL, et al. (November 2001). "Muscle size responses to strength training in young and older men and women". Journal of the American Geriatrics Society. 49 (11): 1428–1433. doi:10.1046/j.1532-5415.2001.4911233.x. PMID 11890579. S2CID 31872193.
- 35. __^ Massaro D, Massaro GD (December 2004). <u>"Estrogen regulates pulmonary alveolar formation, loss, and regeneration in mice"</u> (PDF). American Journal of Physiology. Lung Cellular and Molecular Physiology. **287** (6): L1154–L1159. <u>doi:10.1152/ajplung.00228.2004</u>. <u>PMID 15298854</u>. <u>S2CID 24642944</u>. Archived from <u>the original (PDF)</u> on 25 February 2019.
- 36. <u>^</u> Christensen A, Bentley GE, Cabrera R, Ortega HH, Perfito N, Wu TJ, Micevych P (July 2012). <u>"Hormonal regulation of female reproduction"</u>. Hormone and Metabolic Research. **44** (8): 587–591. <u>doi:10.1055/s-0032-1306301</u>. <u>PMC</u> <u>3647363</u>. <u>PMID</u> <u>22438212</u>.
- 37. <u>^</u> Handa RJ, Ogawa S, Wang JM, Herbison AE (January 2012). <u>"Roles for oestrogen receptor β in adult brain function"</u>. Journal of Neuroendocrinology. **24** (1): 160–173. <u>doi:10.1111/j.1365-2826.2011.02206.x. PMC</u> <u>3348521. PMID</u> <u>21851428</u>.
- 38. ____ Kow LM, Pfaff DW (May 1998). "Mapping of neural and signal transduction pathways for lordosis in the search for estrogen actions on the central nervous system". Behavioural Brain Research. 92 (2): 169–180. doi:10.1016/S0166-4328(97)00189-7. PMID 9638959. S2CID 28276218.
- 39. <u>^</u> Warnock JK, Swanson SG, Borel RW, Zipfel LM, Brennan JJ (2005). "Combined esterified estrogens and methyltestosterone versus esterified estrogens alone in the treatment of loss of sexual interest in surgically menopausal women". Menopause. **12** (4): 374–384. doi:10.1097/01.GME.0000153933.50860.FD. PMID 16037752. S2CID 24557071.
- 40. <u>^</u> Heiman JR, Rupp H, Janssen E, Newhouse SK, Brauer M, Laan E (May 2011). "Sexual desire, sexual arousal and hormonal differences in premenopausal US and Dutch women with and without low sexual desire". Hormones and Behavior. **59** (5): 772–779. doi:10.1016/j.yhbeh.2011.03.013. PMID 21514299. S2CID 20807391.
- 41. <u>^</u> Brisken C, O'Malley B (December 2010). <u>"Hormone action in the mammary gland"</u>. Cold Spring Harbor Perspectives in Biology. **2** (12): a003178. <u>doi:10.1101/cshperspect.a003178</u>. <u>PMC 2982168</u>. <u>PMID 20739412</u>.
- 42. <u>^</u> Kleinberg DL (February 1998). "Role of IGF-I in normal mammary development". Breast Cancer Research and Treatment. **47** (3): 201–208. doi:10.1023/a:1005998832636. PMID 9516076. S2CID 30440069.
- 43. ^ Jump up to: ^{a b c} Johnson LR (2003). <u>Essential Medical Physiology</u>. Academic Press. p. 770. <u>ISBN 978-0-12-387584-6</u>.
- 44. ^ Jump up to: ^{a b c} Norman AW, Henry HL (30 July 2014). <u>Hormones</u>. Academic Press. p. 311. ISBN 978-0-08-091906-5.

- 45. ^ Jump up to: ^a Coad J, Dunstall M (2011). <u>Anatomy and Physiology for Midwives, with Pageburst online access, 3: Anatomy and Physiology for Midwives</u>. Elsevier Health Sciences. p. 413. <u>ISBN 978-0-7020-3489-3</u>.
- 46. <u>^ Haslam SZ</u>, Osuch JR (1 January 2006). <u>Hormones and Breast Cancer in Post-Menopausal Women</u>. IOS Press. p. 69. <u>ISBN 978-1-58603-653-9</u>.
- 47. <u>^</u> Silbernagl S, Despopoulos A (1 January 2011). <u>Color Atlas of Physiology</u>. Thieme. pp. 305–. <u>ISBN 978-3-13-149521-1</u>.
- 48. <u>^ Fadem B (2007). High-yield Comprehensive USMLE Step 1 Review</u>. Lippincott Williams & Wilkins. pp. 445–. <u>ISBN 978-0-7817-7427-7</u>.
- 49. <u>^</u> Blackburn S (14 April 2014). <u>Maternal, Fetal, & Neonatal Physiology</u>. Elsevier Health Sciences. pp. 146–. <u>ISBN 978-0-323-29296-2</u>.
- 50. <u>^ Strauss JF, Barbieri RL (13 September 2013). Yen and Jaffe's Reproductive Endocrinology</u>. Elsevier Health Sciences. pp. 236–. <u>ISBN 978-1-4557-2758-2</u>.
- 51. <u>^</u> Wilson CB, Nizet V, Maldonado Y, Remington JS, Klein JO (24 February 2015). <u>Remington and Klein's Infectious Diseases of the Fetus and Newborn Infant</u>. Elsevier Health Sciences. pp. 190–. <u>ISBN 978-0-323-24147-2</u>.
- 52. <u>^ Zárate S, Stevnsner T, Gredilla R (2017). "Role of Estrogen and Other Sex Hormones in Brain Aging. Neuroprotection and DNA Repair". Frontiers in Aging Neuroscience. 9: 430. doi:10.3389/fnagi.2017.00430. PMC 5743731. PMID 29311911.</u>
- 53. <u>^ Sherwin BB (February 2012). "Estrogen and cognitive functioning in women: lessons we have learned"</u>. Behavioral Neuroscience. **126** (1): 123–127. doi:10.1037/a0025539. PMC 4838456. PMID 22004260.
- 54. <u>^</u> Hara Y, Waters EM, McEwen BS, Morrison JH (July 2015). <u>"Estrogen Effects on Cognitive and Synaptic Health Over the Lifecourse"</u>. Physiological Reviews. **95** (3): 785–807. doi:10.1152/physrev.00036.2014. PMC 4491541. PMID 26109339.
- 55. <u>^</u> Korol DL, Pisani SL (August 2015). <u>"Estrogens and cognition: Friends or foes?: An evaluation of the opposing effects of estrogens on learning and memory"</u>. Hormones and Behavior. **74**: 105–115. <u>doi:10.1016/j.yhbeh.2015.06.017</u>. <u>PMC</u> 4573330. <u>PMID</u> 26149525.
- 56. <u>^</u> Au A, Feher A, McPhee L, Jessa A, Oh S, Einstein G (January 2016). <u>"Estrogens, inflammation and cognition"</u>. Frontiers in Neuroendocrinology. **40**: 87–100. <u>doi:10.1016/j.yfrne.2016.01.002</u>. <u>PMID 26774208</u>.
- 57. <u>^</u> Jacobs E, D'Esposito M (April 2011). <u>"Estrogen shapes dopamine-dependent cognitive processes: implications for women's health"</u>. The Journal of Neuroscience. **31** (14): 5286–5293. doi:10.1523/JNEUROSCI.6394-10.2011. PMC 3089976. PMID 21471363.
- Colzato LS, Hommel B (1 January 2014). "Effects of estrogen on higher-order cognitive functions in unstressed human females may depend on individual variation in dopamine baseline levels". Frontiers in Neuroscience. 8:

 65. doi:10.3389/fnins.2014.00065. PMC 3985021. PMID 24778605.
- 59. <u>^</u> Hogervorst E (March 2013). "Estrogen and the brain: does estrogen treatment improve cognitive function?". Menopause International. **19** (1): 6–19. doi:10.1177/1754045312473873. PMID 27951525. S2CID 10122688.
- 60. <u>^</u> Douma SL, Husband C, O'Donnell ME, Barwin BN, Woodend AK (2005). "Estrogen-related mood disorders: reproductive life cycle factors". ANS. Advances in Nursing Science. **28** (4): 364–375. doi:10.1097/00012272-200510000-00008. PMID 16292022. S2CID 9172877.
- 61. __Osterlund MK, Witt MR, Gustafsson JA (December 2005). "Estrogen action in mood and neurodegenerative disorders: estrogenic compounds with selective properties-the next generation of therapeutics". Endocrine. 28 (3): 235–242. doi:10.1385/ENDO:28:3:235. PMID 16388113. S2CID 8205014.
- 62. <u>^</u> Lasiuk GC, Hegadoren KM (October 2007). "The effects of estradiol on central serotonergic systems and its relationship to mood in women". Biological Research for Nursing. **9** (2): 147–160. doi:10.1177/1099800407305600. PMID 17909167. S2CID 37965502.
- 63. <u>^</u> Grigoriadis S, Seeman MV (June 2002). <u>"The role of estrogen in schizophrenia: implications for schizophrenia practice guidelines for women"</u>. Canadian Journal of Psychiatry. **47** (5): 437–442. <u>doi:10.1177/070674370204700504</u>. <u>PMID</u> <u>12085678</u>.
- 64. <u>^ "PMDD/PMS"</u>. The Massachusetts General Hospital Center for Women's Mental Health. Retrieved 12 January 2019.

- 65. <u>^</u> Hill RA, McInnes KJ, Gong EC, Jones ME, Simpson ER, Boon WC (February 2007). "Estrogen deficient male mice develop compulsive behavior". Biological Psychiatry. **61** (3): 359–366. doi:10.1016/j.biopsych.2006.01.012. PMID 16566897. S2CID 22669945.
- 66. <u>^</u> Benmansour S, Weaver RS, Barton AK, Adeniji OS, Frazer A (April 2012). <u>"Comparison of the effects of estradiol and progesterone on serotonergic function"</u>. Biological Psychiatry. **71** (7): 633–641. doi:10.1016/j.biopsych.2011.11.023. <u>PMC</u> 3307822. <u>PMID</u> 22225849.
- 67. <u>^</u> Berg SJ, Wynne-Edwards KE (June 2001). "Changes in testosterone, cortisol, and estradiol levels in men becoming fathers". Mayo Clinic Proceedings. **76** (6): 582–592. doi:10.4065/76.6.582. PMID 11393496.
- 68. ^ Jump up to: ^{a b c} Cao X, Xu P, Oyola MG, Xia Y, Yan X, Saito K, et al. (October 2014). <u>"Estrogens stimulate serotonin neurons to inhibit binge-like eating in mice"</u>. The Journal of Clinical Investigation. **124** (10): 4351–4362. <u>doi:10.1172/JCI74726</u>. <u>PMC</u> 4191033. <u>PMID</u> 25157819.
- 69. <u>^</u> Jimerson DC, Lesem MD, Kaye WH, Hegg AP, Brewerton TD (September 1990). "Eating disorders and depression: is there a serotonin connection?". Biological Psychiatry. **28** (5): 443–454. doi:10.1016/0006-3223(90)90412-u. PMID 2207221. S2CID 31058047.
- 70. <u>^ Klump KL, Keel PK, Racine SE, Burt SA, Burt AS, Neale M, et al. (February 2013). "The interactive effects of estrogen and progesterone on changes in emotional eating across the menstrual cycle"</u>. Journal of Abnormal Psychology. **122** (1): 131–137. doi:10.1037/a0029524. PMC 3570621. PMID 22889242.
- 71. ^ Jump up to: ^a <u>b</u> Edler C, Lipson SF, Keel PK (January 2007). "Ovarian hormones and binge eating in bulimia nervosa". Psychological Medicine. **37** (1): 131–141. doi:10.1017/S0033291706008956. PMID 17038206. S2CID 36609028.
- 72. ^ Jump up to: ^{a b} Klump KL, Racine SE, Hildebrandt B, Burt SA, Neale M, Sisk CL, et al. (September 2014). <u>"Ovarian Hormone Influences on Dysregulated Eating: A Comparison of Associations in Women with versus without Binge Episodes"</u>. Clinical Psychological Science. **2** (4): 545–559. doi:10.1177/2167702614521794. PMC 4203460. PMID 25343062.
- 73. <u>^ Klump KL, Keel PK, Culbert KM, Edler C (December 2008). "Ovarian hormones and binge eating: exploring associations in community samples"</u>. Psychological Medicine. **38** (12): 1749–1757. doi:10.1017/S0033291708002997. PMC 2885896. PMID 18307829.
- 74. <u>^</u> Lester NA, Keel PK, Lipson SF (January 2003). "Symptom fluctuation in bulimia nervosa: relation to menstrual-cycle phase and cortisol levels". Psychological Medicine. **33** (1): 51–60. doi:10.1017/s0033291702006815. PMID 12537036. S2CID 21497515.
- 75. <u>^</u> Wu MV, Manoli DS, Fraser EJ, Coats JK, Tollkuhn J, Honda S, et al. (October 2009). <u>"Estrogen masculinizes neural pathways and sex-specific behaviors"</u>. Cell. **139** (1): 61–72. <u>doi:10.1016/j.cell.2009.07.036</u>. <u>PMC</u> <u>2851224</u>. <u>PMID</u> <u>19804754</u>.
- 76. <u>^</u> Rochira V, Carani C (October 2009). <u>"Aromatase deficiency in men: a clinical perspective"</u>. Nature Reviews. Endocrinology. **5** (10): 559–568. doi:10.1038/nrendo.2009.176. PMID 19707181. S2CID 22116130.
- 77. <u>^</u> Wilson JD (September 2001). <u>"Androgens, androgen receptors, and male gender role behavior"</u> (PDF). Hormones and Behavior. **40** (2): 358–366. <u>doi:10.1006/hbeh.2001.1684</u>. <u>PMID 11534997</u>. <u>S2CID 20480423</u>. Archived from <u>the original</u> (PDF) on 26 February 2019.
- 78. <u>^</u> Baum MJ (November 2006). "Mammalian animal models of psychosexual differentiation: when is 'translation' to the human situation possible?". Hormones and Behavior. **50** (4): 579–588. doi:10.1016/j.yhbeh.2006.06.003. PMID 16876166. S2CID 7465192.
- 79. <u>^</u> Rosano GM, Panina G (1999). "Oestrogens and the heart". Therapie. **54** (3): 381–385. PMID 10500455.
- 80. <u>^</u> Nadkarni S, Cooper D, Brancaleone V, Bena S, Perretti M (November 2011). <u>"Activation of the annexin A1 pathway underlies the protective effects exerted by estrogen in polymorphonuclear leukocytes"</u>. Arteriosclerosis, Thrombosis, and Vascular Biology. **31** (11): 2749–2759. doi:10.1161/ATVBAHA.111.235176. PMC 3357483. PMID 21836070.
- 81. <u>^</u> Stokar J, Gurt I, Cohen-Kfir E, Yakubovsky O, Hallak N, Benyamini H, et al. (June 2022). <u>"Hepatic adropin is regulated by estrogen and contributes to adverse metabolic phenotypes in ovariectomized mice"</u>. Molecular Metabolism. **60**: 101482. doi:10.1016/j.molmet.2022.101482. PMC 9044006. PMID 35364299.
- 82. <u>^</u> Abdul Sultan A, West J, Stephansson O, Grainge MJ, Tata LJ, Fleming KM, Humes D, Ludvigsson JF (November 2015). <u>"Defining venous thromboembolism and measuring its incidence</u>

- using Swedish health registries: a nationwide pregnancy cohort study". BMJ Open. 5 (11): e008864. doi:10.1136/bmjopen-2015-008864. PMC 4654387. PMID 26560059.
- 83. ^ Jump up to: ^a ^b Foo YZ, Nakagawa S, Rhodes G, Simmons LW (February 2017). "The effects of sex hormones on immune function: a meta-analysis". Biological Reviews of the Cambridge Philosophical Society. **92** (1): 551–571. doi:10.1111/brv.12243. PMID 26800512.
- 84. <u>^</u> Taneja V (27 August 2018). "Sex Hormones Determine Immune Response". Frontiers in Immunology. **9**: 1931. doi:10.3389/fimmu.2018.01931. PMID 30210492.
- 85. ^ Jump up to: ^{a b} Roved J, Westerdahl H, Hasselquist D (February 2017). "Sex differences in immune responses: Hormonal effects, antagonistic selection, and evolutionary consequences". Hormones and Behavior. 88: 95–105. doi:10.1016/j.yhbeh.2016.11.017. PMID 27956226.
- 86. <u>^</u> Khan D, Ansar Ahmed S (6 January 2016). "The Immune System Is a Natural Target for Estrogen Action: Opposing Effects of Estrogen in Two Prototypical Autoimmune Diseases". Frontiers in Immunology. **6**: 635. doi:10.3389/fimmu.2015.00635. PMID 26779182.
- 87. ^ Jump up to: ^{a b c} Kovats S (April 2015). "Estrogen receptors regulate innate immune cells and signaling pathways". Cellular Immunology. **294** (2): 63–69. doi:10.1016/j.cellimm.2015.01.018. PMID 25682174.
- 88. <u>^ Prior JC</u> (2018). <u>Estrogen's Storm Season: stories of perimenopause</u>. Vancouver, British Columbia: CeMCOR (Centre for Menstrual Cycle and Ovulation Research). <u>ISBN</u> <u>9780973827521</u>. Retrieved 24 July 2021. [...] high estrogen amplifies your stress hormone responses to stressful things [...]
- 89. <u>^ Häggström M, Richfield D (2014). "Diagram of the pathways of human</u> steroidogenesis". WikiJournal of Medicine. **1** (1). doi:10.15347/wjm/2014.005. ISSN 2002-4436.
- 90. <u>^</u> Marieb E (2013). Anatomy & physiology. Benjamin-Cummings. p. 903. <u>ISBN</u> <u>978-0-321-88760-3</u>.
- 91. <u>^</u> Hemsell DL, Grodin JM, Brenner PF, Siiteri PK, MacDonald PC (March 1974). "Plasma precursors of estrogen. II. Correlation of the extent of conversion of plasma androstenedione to estrone with age". The Journal of Clinical Endocrinology and Metabolism. **38** (3): 476–479. doi:10.1210/jcem-38-3-476. PMID 4815174.
- 92. <u>^</u> Barakat R, Oakley O, Kim H, Jin J, Ko CJ (September 2016). <u>"Extra-gonadal sites of estrogen biosynthesis and function"</u>. BMB Reports. **49** (9): 488–496. <u>doi:10.5483/BMBRep.2016.49.9.141</u>. <u>PMC 5227141</u>. <u>PMID 27530684</u>.
- 93. ^ Jump up to: ^a b Nelson LR, Bulun SE (September 2001). "Estrogen production and action". Journal of the American Academy of Dermatology. **45** (3 Suppl): S116–S124. doi:10.1067/mjd.2001.117432. PMID 11511861.
- 94. <u>^</u> Labrie F, Bélanger A, Luu-The V, Labrie C, Simard J, Cusan L, et al. (1998). "DHEA and the intracrine formation of androgens and estrogens in peripheral target tissues: its role during aging". Steroids. **63** (5–6): 322–328. doi:10.1016/S0039-128X(98)00007-5. PMID 9618795. S2CID 37344052.
- 95. <u>^</u> Kuhl H (August 2005). "Pharmacology of estrogens and progestogens: influence of different routes of administration". Climacteric. **8** (Suppl 1): 3–63. doi:10.1080/13697130500148875. PMID 16112947. S2CID 24616324.
- 96. <u>^</u> Wesp LM, Deutsch MB (March 2017). "Hormonal and Surgical Treatment Options for Transgender Women and Transfeminine Spectrum Persons". The Psychiatric Clinics of North America. **40** (1): 99–111. doi:10.1016/j.psc.2016.10.006. PMID 28159148.
- 97. <u>^ Tata JR (June 2005)</u>. <u>"One hundred years of hormones"</u>. EMBO Reports. **6** (6): 490–496. <u>doi:10.1038/sj.embor.7400444</u>. <u>PMC 1369102</u>. <u>PMID 15940278</u>.
- 98. ^ Jump up to: ^a <u>b</u> "Origin in Biomedical Terms: oestrogen or oestrogen". Bioetymology. Retrieved 24 January 2018.
- 99. <u>^</u> "Council on Pharmacy and Chemistry". Journal of the American Medical Association. **107** (15): 1221–3. 1936. <u>doi:10.1001/jama.1936.02770410043011</u>.
- 100. <u>^ "Greek Word Study Tool: oistros". Perseus Digital Library. Retrieved 28 December 2011.</u>
- 101. <u>^</u> Fang H, Tong W, Shi LM, Blair R, Perkins R, Branham W, et al. (March 2001). "Structure-activity relationships for a large diverse set of natural, synthetic, and environmental estrogens". Chemical Research in Toxicology. **14** (3): 280–294. CiteSeerX 10.1.1.460.20. doi:10.1021/tx000208y. PMID 11258977.

- 102. <u>^</u> Wise A, O'Brien K, Woodruff T (January 2011). "Are oral contraceptives a significant contributor to the estrogenicity of drinking water?". Environmental Science & Technology. **45** (1): 51–60. <u>doi:10.1021/es1014482</u>. <u>PMID 20977246</u>. <u>Lay summary</u> Chemical & Engineering News. {{cite journal}}: Cite uses deprecated parameter |lay-url=(help)
- 103. <u>^</u> Liney KE, Jobling S, Shears JA, Simpson P, Tyler CR (October 2005). <u>"Assessing the sensitivity of different life stages for sexual disruption in roach (Rutilus rutilus) exposed to effluents from wastewater treatment works"</u>. Environmental Health Perspectives. 113 (10): 1299–1307. doi:10.1289/ehp.7921. <u>PMC</u> 1281270. <u>PMID</u> 16203238.
- 104. ___ Jobling S, Williams R, Johnson A, Taylor A, Gross-Sorokin M, Nolan M, et al. (April 2006). <u>"Predicted exposures to steroid estrogens in U.K. rivers correlate with widespread sexual disruption in wild fish populations"</u>. Environmental Health Perspectives. **114** (Suppl 1): 32–39. <u>doi:10.1289/ehp.8050</u>. <u>PMC</u> 1874167. <u>PMID</u> 16818244.
- 105. <u>^ Sanghavi DM (17 October 2006). "Preschool Puberty, and a Search for the Causes"</u>. <u>The New York Times</u>. Retrieved 4 June 2008.
- 106. ^ Jump up to: ^a ^b FDA (February 1995). <u>"Products containing estrogenic hormones, placental extract or vitamins"</u>. Guide to Inspections of Cosmetic Product Manufacturers. Archived from <u>the original</u> on 14 October 2007. Retrieved 24 October 2006.

External links