

Endocrine system

- ☐ 101 languages
- [Article](#)
 - [Talk](#)
 - [Read](#)
 - [Edit](#)
 - [View history](#)

- ☐ Tools

From Wikipedia, the free encyclopedia

Endocrine system

The diagram illustrates the human endocrine system. It features two human figures, one male and one female, with callouts to various glands. The male figure shows the pineal gland (in the brain), pituitary gland (below the brain), parathyroid glands (four spots on the thyroid gland in the neck), thyroid gland (in the neck), adrenal glands (two glands, each sitting atop a kidney), pancreas (in the abdominal region), ovary (in the female pelvic region), and testis (in the male pelvic region). The female figure shows the same glands except for the testis. The glands are color-coded: the brain glands are pink, the thyroid and adrenal glands are yellow, the pancreas is orange, and the reproductive glands are red.

Main [glands](#) of the human endocrine system

Details

Identifiers

Latin	<i>Systema endocrinum</i>
MeSH	D004703
FMA	9668
	Anatomical terminology

[\[edit on Wikidata\]](#)

The **endocrine system**^[1] is a messenger system comprising feedback loops of the [hormones](#) released by internal [glands](#) of an [organism](#) directly into the [circulatory system](#), regulating distant target organs. In [vertebrates](#), the [hypothalamus](#) is the neural control center for all endocrine systems. In [humans](#), the major [endocrine glands](#) are the [thyroid gland](#), [parathyroid gland](#), [pituitary gland](#), [pineal gland](#), the [testes](#) (male), [ovaries](#) (female), and the [adrenal glands](#).

The [hypothalamus](#), [pancreas](#), and [thymus](#) also function as endocrine glands, among other functions. Other organs, such as the [kidneys](#), also have roles within the endocrine system by secreting certain hormones. The study of the endocrine system and its disorders is known as [endocrinology](#).^[2] It is one of the most important systems of the human body.

Glands that signal each other in sequence are often referred to as an axis, such as the [hypothalamic-pituitary-adrenal axis](#). In addition to the specialized endocrine organs mentioned above, many other organs that are part of other body systems have secondary endocrine functions, including [bone](#), [kidneys](#), [liver](#), [heart](#) and [gonads](#). For example, the kidney secretes the endocrine hormone [erythropoietin](#). Hormones can be amino acid complexes, [steroids](#), [eicosanoids](#), [leukotrienes](#), or [prostaglandins](#).^[3]

The endocrine system can be contrasted to both [exocrine glands](#), which secrete hormones to the outside of the body, and [paracrine signalling](#) between cells over a relatively short distance. Endocrine glands have no [ducts](#), are vascular, and commonly have intracellular vacuoles or granules that store their hormones. In contrast, exocrine glands, such as [salivary glands](#), [sweat glands](#), and glands within the [gastrointestinal tract](#), tend to be much less vascular and have ducts or a hollow [lumen](#). Endocrinology is a branch of [internal medicine](#).^[3]

Structure^[edit]

Major endocrine systems^[edit]

The human endocrine system consists of several systems that operate via [feedback loops](#). Several important feedback systems are mediated via the hypothalamus and pituitary.^[4]

- [TRH – TSH – T3/T4](#)
- [GnRH – LH/FSH – sex hormones](#)
- [CRH – ACTH – cortisol](#)
- [Renin – angiotensin – aldosterone](#)
- [leptin vs. Ghrelin](#)

Glands^[edit]

Main article: [Endocrine gland](#)

Endocrine glands are [glands](#) of the endocrine system that secrete their products, [hormones](#), directly into interstitial spaces where they are absorbed into blood

rather than through a duct. The major glands of the endocrine system include the [pineal gland](#), [pituitary gland](#), [pancreas](#), [ovaries](#), [testes](#), [thyroid gland](#), [parathyroid gland](#), [hypothalamus](#) and [adrenal glands](#). The hypothalamus and pituitary gland are [neuroendocrine organs](#).

The hypothalamus and the anterior pituitary are two out of the three endocrine glands that are important in cell signaling. They are both part of the HPA axis which is known to play a role in cell signaling in the nervous system.

Hypothalamus: The hypothalamus is a key regulator of the autonomic nervous system. The endocrine system has three sets of endocrine outputs^[6] which include the magnocellular system, the parvocellular system, and autonomic intervention. The magnocellular is involved in the expression of oxytocin or vasopressin. The parvocellular is involved in controlling the secretion of hormones from the anterior pituitary.

Anterior Pituitary: The main role of the anterior pituitary gland is to produce and secrete tropic hormones.^[6] Some examples of tropic hormones secreted by the anterior pituitary gland include TSH, ACTH, GH, LH, and FSH.

Cells[\[edit\]](#)

There are many types of cells that make up the endocrine system and these cells typically make up larger tissues and organs that function within and outside of the endocrine system.

- [Hypothalamus](#)
- [Anterior pituitary gland](#)
- [Pineal gland](#)
- [Posterior pituitary gland](#)
 - The posterior pituitary gland is a section of the pituitary gland. This organ does not produce any hormone but stores and secretes hormones such as antidiuretic hormone (ADH) which is synthesized by supraoptic nucleus of hypothalamus and oxytocin which is synthesized by paraventricular nucleus of hypothalamus. ADH functions to help the body to retain water; this is important in maintaining a homeostatic balance between blood solutions and water. Oxytocin functions to induce uterine contractions, stimulate lactation, and allows for ejaculation.^{[7][8]}
- [Thyroid gland](#)
 - [follicular cells](#) of the thyroid gland produce and secrete T_3 and T_4 in response to elevated levels of [TRH](#), produced by the [hypothalamus](#), and subsequent elevated levels of [TSH](#), produced by the [anterior pituitary gland](#), which further regulates the [metabolic activity](#) and rate of all cells, including [cell growth](#) and [tissue differentiation](#).
- [Parathyroid gland](#)
 - [Epithelial](#) cells of the parathyroid glands are richly supplied with blood from the [inferior](#) and [superior thyroid arteries](#) and secrete [parathyroid hormone](#) (PTH). PTH acts on bone, the kidneys, and the [GI tract](#) to

increase [calcium reabsorption](#) and phosphate excretion. In addition, PTH stimulates the conversion of [Vitamin D](#) to its most active variant, [1,25-dihydroxyvitamin D₃](#), which further stimulates [calcium](#) absorption in the GI tract.^[3]

- [Thymus Gland](#)
- [Adrenal glands](#)
 - [Adrenal cortex](#)
 - [Adrenal medulla](#)
- [Pancreas](#)
 - Pancreas contain nearly 1 to 2 million islets of Langerhans (a tissue which consists cells that secrete hormones) and acini. Acini secretes digestive enzymes.^[9]
 - [Alpha cells](#)
 - The alpha cells of the pancreas secrete hormones to maintain homeostatic blood sugar. Insulin is produced and excreted to lower blood sugar to normal levels. Glucagon, another hormone produced by alpha cells, is secreted in response to low blood sugar levels; glucagon stimulates glycogen stores in the liver to release sugar into the bloodstream to raise blood sugar to normal levels.^[10]
 - [Beta cells](#)
 - 60% of the cells present in [islet of Langerhans](#) are beta cells. Beta cells secrete [insulin](#). Along with glucagon, insulin helps in maintaining glucose levels in our body. Insulin decreases blood glucose level (a hypoglycemic hormone) whereas glucagon increases blood glucose level.^[9]
 - [Delta cells](#)
 - [F Cells](#)
- [Ovaries](#)
 - [Granulosa cells](#)
- [Testis](#)
 - [Leydig cells](#)^[11]

Development^[edit]

Main article: [Development of the endocrine system](#)

The **fetal endocrine system** is one of the first systems to develop during [prenatal development](#).

Adrenal glands^[edit]

The fetal [adrenal cortex](#) can be identified within four weeks of [gestation](#). The adrenal cortex originates from the thickening of the intermediate [mesoderm](#). At five to six weeks of gestation, the [mesonephros](#) differentiates into a tissue known as the genital ridge. The genital ridge produces the steroidogenic cells for both the gonads and the adrenal cortex. The adrenal medulla is derived from [ectodermal cells](#). Cells that will become adrenal tissue move retroperitoneally to the upper portion of the mesonephros. At seven weeks of gestation, the adrenal cells are joined by sympathetic cells that originate from

the neural crest to form the [adrenal medulla](#). At the end of the eighth week, the adrenal glands have been encapsulated and have formed a distinct organ above the developing kidneys. At birth, the adrenal glands weigh approximately eight to nine grams (twice that of the adult adrenal glands) and are 0.5% of the total body weight. At 25 weeks, the adult adrenal cortex zone develops and is responsible for the primary synthesis of steroids during the early postnatal weeks.

Thyroid gland[\[edit\]](#)

The [thyroid gland](#) develops from two different clusterings of embryonic cells. One part is from the thickening of the pharyngeal floor, which serves as the precursor of the thyroxine (T_4) producing follicular cells. The other part is from the caudal extensions of the fourth pharyngobranchial pouches which results in the parafollicular calcitonin-secreting cells. These two structures are apparent by 16 to 17 days of gestation. Around the 24th day of gestation, the foramen cecum, a thin, flask-like diverticulum of the median [anlage](#) develops. At approximately 24 to 32 days of gestation the median anlage develops into a bilobed structure. By 50 days of gestation, the medial and lateral anlage have fused together. At 12 weeks of gestation, the fetal thyroid is capable of storing iodine for the production of [TRH](#), [TSH](#), and free thyroid hormone. At 20 weeks, the fetus is able to implement feedback mechanisms for the production of thyroid hormones. During fetal development, T_4 is the major thyroid hormone being produced while triiodothyronine (T_3) and its inactive derivative, reverse T_3 , are not detected until the third trimester.

Parathyroid glands[\[edit\]](#)

A lateral and ventral view of an embryo showing the third (inferior) and fourth (superior) parathyroid glands during the 6th week of embryogenesis

Once the embryo reaches four weeks of gestation, the [parathyroid glands](#) begins to develop. The human embryo forms five sets of [endoderm](#)-lined pharyngeal pouches. The third and fourth pouch are responsible for developing into the inferior and superior parathyroid glands, respectively. The third pharyngeal pouch encounters the developing thyroid gland and they migrate down to the lower poles of the thyroid lobes. The fourth pharyngeal pouch later encounters the developing thyroid gland and migrates to the upper poles of the thyroid lobes. At 14 weeks of gestation, the parathyroid glands begin to enlarge from 0.1 mm in diameter to approximately 1 – 2 mm at birth. The developing parathyroid glands are physiologically functional beginning in the second trimester.

Studies in mice have shown that interfering with the [HOX15](#) gene can cause parathyroid gland [aplasia](#), which suggests the gene plays an important role in the development of the parathyroid gland. The genes, [TBX1](#), [CRKL](#), [GATA3](#), [GCM2](#), and [SOX3](#) have also been shown to play a crucial role in the formation of the parathyroid gland. Mutations in TBX1 and CRKL genes are correlated with [DiGeorge syndrome](#), while mutations in GATA3 have also resulted in a DiGeorge-like syndrome. Malformations in the GCM2 gene have resulted in hypoparathyroidism. Studies on SOX3 gene mutations have demonstrated that it plays a role in parathyroid development. These mutations also lead to varying degrees of hypopituitarism.

Pancreas[\[edit\]](#)

The human fetal [pancreas](#) begins to develop by the fourth week of gestation. Five weeks later, the pancreatic [alpha](#) and [beta cells](#) have begun to emerge. Reaching eight to ten weeks into development, the pancreas starts producing [insulin](#), [glucagon](#), [somatostatin](#), and [pancreatic polypeptide](#). During the early stages of fetal development, the number of pancreatic alpha cells outnumbers the number of pancreatic beta cells. The alpha cells reach their peak in the middle stage of gestation. From the middle stage until term, the beta cells continue to increase in number until they reach an approximate 1:1 ratio with the alpha cells. The insulin concentration within the fetal pancreas is 3.6 pmol/g at seven to ten weeks, which rises to 30 pmol/g at 16–25 weeks of gestation. Near term, the insulin concentration increases to 93 pmol/g. The endocrine cells have dispersed throughout the body within 10 weeks. At 31 weeks of development, the [islets of Langerhans](#) have differentiated.

While the fetal pancreas has functional beta cells by 14 to 24 weeks of gestation, the amount of insulin that is released into the bloodstream is relatively low. In a study of pregnant women carrying fetuses in the mid-gestation and near term stages of development, the fetuses did not have an increase in plasma insulin levels in response to injections of high levels of glucose. In contrast to insulin, the fetal plasma glucagon levels are relatively high and continue to increase during development. At the mid-stage of gestation, the glucagon concentration is 6 µg/g, compared to 2 µg/g in adult humans. Just like insulin, fetal glucagon plasma levels do not change in response to an infusion of glucose. However, a study of an infusion of alanine into pregnant women was shown to increase the cord blood and maternal glucagon concentrations, demonstrating a fetal response to amino acid exposure.

As such, while the fetal pancreatic alpha and beta islet cells have fully developed and are capable of hormone synthesis during the remaining fetal maturation, the islet cells are relatively immature in their capacity to produce glucagon and insulin. This is thought to be a result of the relatively stable levels of fetal [serum glucose](#) concentrations achieved via maternal transfer of glucose through the placenta. On the other hand, the stable fetal serum glucose levels could be attributed to the absence of pancreatic signaling initiated by incretins during feeding. In addition, the fetal pancreatic islets cells are unable to sufficiently produce [cAMP](#) and rapidly degrade cAMP by [phosphodiesterase](#) necessary to secrete glucagon and insulin.

During fetal development, the storage of glycogen is controlled by fetal [glucocorticoids](#) and [placental lactogen](#). Fetal insulin is responsible for increasing glucose uptake and lipogenesis during the stages leading up to birth. Fetal cells contain a higher amount of insulin receptors in comparison to adults cells and fetal insulin receptors are not downregulated in cases of [hyperinsulinemia](#). In comparison, fetal haptic glucagon receptors are lowered in comparison to adult cells and the glycemic effect of glucagon is blunted. This temporary physiological change aids the increased rate of fetal development during the final trimester. Poorly managed maternal [diabetes mellitus](#) is linked to [fetal macrosomia](#), increased risk of miscarriage, and defects in fetal development. Maternal hyperglycemia is also linked to increased insulin levels and beta cell hyperplasia in the post-term infant. Children of diabetic mothers are at an increased risk for conditions such as: [polycythemia](#), [renal vein thrombosis](#), [hypocalcemia](#), [respiratory distress](#)

[syndrome](#), [jaundice](#), [cardiomyopathy](#), [congenital heart disease](#), and improper organ development.

Gonads[\[edit\]](#)

Main article: [Development of the gonads](#)

The reproductive system begins development at four to five weeks of gestation with germ cell migration. The bipotential gonad results from the collection of the medioventral region of the urogenital ridge. At the five-week point, the developing gonads break away from the adrenal primordium. Gonadal differentiation begins 42 days following conception.

Male gonadal development[\[edit\]](#)

For males, the [testes](#) form at six fetal weeks and the sertoli cells begin developing by the eight week of gestation. [SRY](#), the sex-determining locus, serves to differentiate the [Sertoli cells](#). The Sertoli cells are the point of origin for [anti-Müllerian hormone](#). Once synthesized, the anti-Müllerian hormone initiates the ipsilateral regression of the Müllerian tract and inhibits the development of female internal features. At 10 weeks of gestation, the Leydig cells begin to produce androgen hormones. The androgen hormone dihydrotestosterone is responsible for the development of the male external genitalia.

The testicles descend during prenatal development in a two-stage process that begins at eight weeks of gestation and continues through the middle of the third trimester. During the transabdominal stage (8 to 15 weeks of gestation), the [gubernacular ligament](#) contracts and begins to thicken. The craniosuspensory ligament begins to break down. This stage is regulated by the secretion of [insulin-like 3](#) (INSL3), a relaxin-like factor produced by the testicles, and the INSL3 G-coupled receptor, LGR8. During the transinguinal phase (25 to 35 weeks of gestation), the testicles descend into the scrotum. This stage is regulated by androgens, the genitofemoral nerve, and calcitonin gene-related peptide. During the second and third trimester, testicular development concludes with the diminution of the fetal Leydig cells and the lengthening and coiling of the [seminiferous cords](#).

Female gonadal development[\[edit\]](#)

For females, the [ovaries](#) become morphologically visible by the 8th week of gestation. The absence of testosterone results in the diminution of the Wolffian structures. The Müllerian structures remain and develop into the fallopian tubes, uterus, and the upper region of the vagina. The [urogenital sinus](#) develops into the urethra and lower region of the vagina, the genital tubercle develops into the clitoris, the urogenital folds develop into the labia minora, and the urogenital swellings develop into the labia majora. At 16 weeks of gestation, the ovaries produce [FSH](#) and [LH/hCG receptors](#). At 20 weeks of gestation, the theca cell precursors are present and oogonia mitosis is occurring. At 25 weeks of gestation, the ovary is morphologically defined and [folliculogenesis](#) can begin.

Studies of gene expression show that a specific complement of genes, such as follistatin and multiple cyclin kinase inhibitors are involved in ovarian development. An assortment of genes and proteins - such as WNT4, RSPO1, FOXL2, and various

estrogen receptors - have been shown to prevent the development of testicles or the lineage of male-type cells.

Pituitary gland[\[edit\]](#)

The [pituitary gland](#) is formed within the rostral neural plate. The Rathke's pouch, a cavity of ectodermal cells of the [oropharynx](#), forms between the fourth and fifth week of gestation and upon full development, it gives rise to the anterior pituitary gland. By seven weeks of gestation, the anterior pituitary vascular system begins to develop. During the first 12 weeks of gestation, the anterior pituitary undergoes cellular differentiation. At 20 weeks of gestation, the [hypophyseal portal system](#) has developed. The Rathke's pouch grows towards the third ventricle and fuses with the diverticulum. This eliminates the lumen and the structure becomes Rathke's cleft. The posterior pituitary lobe is formed from the diverticulum. Portions of the pituitary tissue may remain in the nasopharyngeal midline. In rare cases this results in functioning ectopic hormone-secreting tumors in the nasopharynx.

The functional development of the anterior pituitary involves spatiotemporal regulation of transcription factors expressed in pituitary stem cells and dynamic gradients of local soluble factors. The coordination of the dorsal gradient of pituitary morphogenesis is dependent on neuroectodermal signals from the infundibular bone morphogenetic protein 4 (BMP4). This protein is responsible for the development of the initial invagination of the Rathke's pouch. Other essential proteins necessary for pituitary cell proliferation are [Fibroblast growth factor 8](#) (FGF8), Wnt4, and Wnt5. Ventral developmental patterning and the expression of transcription factors is influenced by the gradients of BMP2 and [sonic hedgehog protein](#) (SHH). These factors are essential for coordinating early patterns of cell proliferation.

Six weeks into gestation, the [corticotroph cells](#) can be identified. By seven weeks of gestation, the anterior pituitary is capable of secreting ACTH. Within eight weeks of gestation, somatotroph cells begin to develop with cytoplasmic expression of human growth hormone. Once a fetus reaches 12 weeks of development, the thyrotrophs begin expression of Beta subunits for TSH, while [gonadotrophs](#) begin to express beta-subunits for LH and FSH. Male fetuses predominately produced LH-expressing gonadotrophs, while female fetuses produce an equal expression of LH and FSH expressing gonadotrophs. At 24 weeks of gestation, prolactin-expressing [lactotrophs](#) begin to emerge.

Function[\[edit\]](#)

See also: [List of human endocrine organs and actions](#)

Hormones[\[edit\]](#)

Main article: [Hormone](#)

A [hormone](#) is any of a class of [signaling molecules](#) produced by cells in [glands](#) in [multicellular organisms](#) that are transported by the [circulatory system](#) to target distant organs to regulate [physiology](#) and [behaviour](#). Hormones have diverse chemical structures, mainly of 3 classes: [eicosanoids](#), [steroids](#), and [amino acid/protein](#) derivatives ([amines](#), [peptides](#), and [proteins](#)). The glands that secrete

hormones comprise the endocrine system. The term hormone is sometimes extended to include chemicals produced by cells that affect the same cell ([autocrine](#) or [intracrine signalling](#)) or nearby cells ([paracrine signalling](#)).

Hormones are used to communicate between [organs](#) and tissues for [physiological](#) regulation and [behavioral](#) activities, such as digestion, [metabolism](#), [respiration](#), [tissue](#) function, [sensory perception](#), [sleep](#), [excretion](#), [lactation](#), [stress](#), [growth and development](#), [movement](#), [reproduction](#), and [mood](#).^{[12][13]}

Hormones affect distant cells by binding to specific [receptor](#) proteins in the target cell resulting in a change in cell function. This may lead to cell type-specific responses that include rapid changes to the activity of existing proteins, or slower changes in the [expression](#) of target genes. Amino acid-based hormones ([amines](#) and [peptide or protein hormones](#)) are water-soluble and act on the surface of target cells via [signal transduction](#) pathways; [steroid hormones](#), being lipid-soluble, move through the [plasma membranes](#) of target cells to act within their [nuclei](#).

Cell signalling^[edit]

The typical mode of [cell signalling](#) in the endocrine system is endocrine signaling, that is, using the circulatory system to reach distant target organs. However, there are also other modes, i.e., paracrine, autocrine, and [neuroendocrine](#) signaling. Purely neurocrine signaling between [neurons](#), on the other hand, belongs completely to the [nervous system](#).

Autocrine^[edit]

Main article: [Autocrine signalling](#)

Autocrine signaling is a form of signaling in which a cell secretes a hormone or chemical messenger (called the autocrine agent) that binds to autocrine receptors on the same cell, leading to changes in the cells.

Paracrine^[edit]

Main article: [Paracrine signalling](#)

Some endocrinologists and clinicians include the paracrine system as part of the endocrine system, but there is not consensus. Paracrine are slower acting, targeting cells in the same tissue or organ. An example of this is [somatostatin](#) which is released by some pancreatic cells and targets other pancreatic cells.^[3]

Juxtacrine^[edit]

Main article: [Juxtacrine signalling](#)

Juxtacrine signaling is a type of intercellular communication that is transmitted via oligosaccharide, lipid, or protein components of a cell membrane, and may affect either the emitting cell or the immediately adjacent cells.^[14]


It occurs between adjacent cells that possess broad patches of closely opposed plasma membrane linked by transmembrane channels known as [connexons](#). The gap between the cells can usually be between only 2 and 4 nm.^[15]

Clinical significance^{[[edit](#)]}

Disease^{[[edit](#)]}



[Disability-adjusted life year](#) for endocrine disorders per 100,000 inhabitants in 2002.^[16]



Main article: [Endocrine diseases](#)

[Diseases of the endocrine system](#) are common,^[17] including conditions such as [diabetes mellitus](#), [thyroid](#) disease, and [obesity](#). Endocrine disease is characterized by misregulated hormone release (a productive [pituitary adenoma](#)), inappropriate response to signaling ([hypothyroidism](#)), lack of a gland ([diabetes mellitus type 1](#), diminished [erythropoiesis](#) in [chronic kidney failure](#)), or structural enlargement in a critical site such as the thyroid ([toxic multinodular goitre](#)). Hypofunction of endocrine glands can occur as a result of loss of reserve, hyposecretion, [agenesis](#), atrophy, or active destruction. Hyperfunction can occur as a result of hypersecretion, loss of suppression, [hyperplastic](#) or [neoplastic](#) change, or hyperstimulation.

Endocrinopathies are classified as primary, secondary, or tertiary. Primary endocrine disease inhibits the action of downstream glands. Secondary endocrine disease is indicative of a problem with the pituitary gland. Tertiary endocrine disease is associated with dysfunction of the hypothalamus and its releasing hormones.^[18]

As the [thyroid](#), and hormones have been implicated in signaling distant tissues to proliferate, for example, the [estrogen receptor](#) has been shown to be involved in certain [breast cancers](#). Endocrine, paracrine, and autocrine signaling have all been implicated in proliferation, one of the required steps of [oncogenesis](#).^[19]

Other common diseases that result from endocrine dysfunction include [Addison's disease](#), [Cushing's disease](#) and [Graves' disease](#). Cushing's disease and Addison's disease are pathologies involving the dysfunction of the adrenal gland. Dysfunction in the adrenal gland could be due to primary or secondary factors and can result in hypercortisolism or hypocortisolism. Cushing's disease is characterized by the hypersecretion of the adrenocorticotrophic hormone (ACTH) due to a pituitary adenoma that ultimately causes endogenous hypercortisolism by stimulating the adrenal glands.^[20] Some clinical signs of Cushing's disease include obesity, moon face, and hirsutism.^[21] Addison's disease is an endocrine disease that results from hypocortisolism caused by adrenal gland insufficiency. Adrenal insufficiency is significant because it is correlated with decreased ability to maintain blood pressure and blood sugar, a defect that can prove to be fatal.^[22]

Graves' disease involves the hyperactivity of the thyroid gland which produces the T3 and T4 hormones.^[21] [Graves' disease](#) effects range from excess sweating, fatigue, heat intolerance and [high blood pressure](#) to swelling of the eyes that causes redness, puffiness and in rare cases reduced or double vision.^[15]

Other animals^[edit]

A neuroendocrine system has been observed in all [animals](#) with a nervous system and all [vertebrates](#) have a hypothalamus-pituitary axis.^[23] All vertebrates have a thyroid, which in [amphibians](#) is also crucial for transformation of larvae into adult form.^{[24][25]} All vertebrates have adrenal gland tissue, with mammals unique in having it organized into layers.^[26] All vertebrates have some form of a renin–angiotensin axis, and all [tetrapods](#) have aldosterone as a primary [mineralocorticoid](#).^{[27][28]}

Additional images^[edit]



Female endocrine system



Male endocrine system

See also^[edit]

- [Endocrine disease](#)
- [Endocrinology](#)
- [List of human endocrine organs and actions](#)
- [Neuroendocrinology](#)
- [Nervous system](#)
- [Paracrine signalling](#)
- [Releasing hormones](#)
- [Tropic hormone](#)

References[[edit](#)]

- ¹. [^] ["Anatomy of the Endocrine System"](#). www.hopkinsmedicine.org. November 19, 2019. Retrieved June 14, 2022.
- ². [^] Gardner, Shoback (2017). *Greenspan's Basic and Clinical Endocrinology* (10th ed.). McGraw Hill / Medical. pp. 49–68. [ISBN 978-1259589287](#).
- ³. [^] ^{Jump up to:[ⓘ] [Ⓘ] [Ⓝ]} Marieb E (2014). *Anatomy & physiology*. Glenview, IL: Pearson Education, Inc. [ISBN 978-0-321-86158-0](#).
- ⁴. [^] Sherwood, L. (1997). [Human Physiology: From Cells to Systems](#). Wadsworth Pub Co. [ISBN 978-0-495-39184-5](#).
- ⁵. [^] Clifford B. Saper; Bradford B. Lowell (December 1, 2014). ["The Hypothalamus"](#). *Current Biology*. **24** (23): R1111-6. [doi:10.1016/j.cub.2014.10.023](#). [PMID 25465326](#). [S2CID 18782796](#).
- ⁶. [^] "Chapter 3. Anterior Pituitary Gland". [Endocrine Physiology](#) (4 ed.). McGraw Hill. 2013.
- ⁷. [^] Quesada, Ivan (2008). ["Physiology of the pancreatic \$\alpha\$ -cell and glucagon secretion: role in glucose homeostasis and diabetes"](#). *Journal of Endocrinology*. **199** (1): 5–19. [doi:10.1677/JOE-08-0290](#). [PMID 18669612](#).
- ⁸. [^] Patel, H.; Jessu, R.; Tiwari, V. (2021). ["Physiology, Posterior Pituitary"](#). How Does The Pancreas Work?. Informed Health. StatPearls. [PMID 30252386](#).
- ⁹. [^] ^{Jump up to:[ⓘ] [Ⓘ] [Ⓝ]} ["267 Endocrine System Facts"](#). Facts Legend. September 19, 2018.
- ¹⁰. [^] Patel, H.; Jessu, R.; Tiwari, V. (2020). ["Physiology, Posterior Pituitary"](#). NCBI. StatPearls. [PMID 30252386](#).
- ¹¹. [^] ["Leydig cell | anatomy | Britannica"](#). www.britannica.com. Retrieved June 14, 2022.
- ¹². [^] Neave N (2008). *Hormones and behaviour: a psychological approach*. Cambridge: Cambridge Univ. Press. [ISBN 978-0-521-69201-4](#).
 - [■] Claire L. Gibson (Winter 2010). ["Hormones and Behaviour: A Psychological Approach"](#). *Perspectives in Biology and Medicine* (Review). **53** (1): 152–155. [doi:10.1353/pbm.0.0141](#). [S2CID 72100830](#).
- ¹³. [^] ["Hormones"](#). MedlinePlus. U.S. National Library of Medicine.
- ¹⁴. [^] Gilbert F (2000). "Juxtacrine Signaling". [Developmental Biology](#) (6th ed.). Sunderland, MA: Sinauer Associates. [ISBN 0-87893-243-7](#).
- ¹⁵. [^] ^{Jump up to:[ⓘ] [Ⓘ] [Ⓝ]} Vander A (2008). [Vander's Human Physiology: the mechanisms of body function](#). Boston: McGraw-Hill Higher Education. pp. 332–333. [ISBN 978-0-07-304962-5](#).
- ¹⁶. [^] ["Mortality and Burden of Disease Estimates for WHO Member States in 2002"](#) (xls). World Health Organization. 2002.
- ¹⁷. [^] Kasper DL, Harrison TR (2005). [Harrison's Principles of Internal Medicine](#). McGraw Hill. pp. 2074. [ISBN 978-0-07-139140-5](#).
- ¹⁸. [^] Macksey LF (2012). *Surgical procedures and anesthetic implications: a handbook for nurse anesthesia practice*. Sudbury, MA: Jones & Bartlett Learning. p. 479. [ISBN 978-0-7637-8057-9](#). [OCLC 632070527](#).
- ¹⁹. [^] Bhowmick NA, Chytil A, Plieth D, Gorska AE, Dumont N, Shappell S, Washington MK, Neilson EG, Moses HL (February 2004). "TGF-beta signaling in fibroblasts modulates the oncogenic potential of adjacent epithelia". *Science*. **303** (5659): 848–51. [Bibcode:2004Sci...303..848B](#). [doi:10.1126/science.1090922](#). [PMID 14764882](#). [S2CID 1703215](#).

20. [^] Buliman A, Tataranu LG, Paun DL, Mirica A, Dumitrache C (2016). "[Cushing's disease: a multidisciplinary overview of the clinical features, diagnosis, and treatment](#)". *Journal of Medicine and Life*. **9** (1): 12–18. [PMC 5152600](#). [PMID 27974908](#).
21. [^] ^{Jump up to:}[^] Vander A (2008). *Vander's Human Physiology: the mechanisms of body function*. Boston: McGraw-Hill Higher Education. pp. 345–347. [ISBN 978-0-07-304962-5](#).
22. [^] Inder WJ, Meyer C, Hunt PJ (June 2015). "Management of hypertension and heart failure in patients with Addison's disease". *Clinical Endocrinology*. **82** (6): 789–92. [doi:10.1111/cen.12592](#). [PMID 25138826](#). [S2CID 13552007](#).
23. [^] Hartenstein V (September 2006). "[The neuroendocrine system of invertebrates: a developmental and evolutionary perspective](#)". *The Journal of Endocrinology*. **190** (3): 555–70. [doi:10.1677/joe.1.06964](#). [PMID 17003257](#).
24. [^] Dickhoff WW, Darling DS (1983). "[Evolution of Thyroid Function and Its Control in Lower Vertebrates](#)". *American Zoologist*. **23** (3): 697–707. [doi:10.1093/icb/23.3.697](#). [JSTOR 3882951](#).
25. [^] Galton VA (January 1, 1988). "[The Role of Thyroid Hormone in Amphibian Development](#)". *Integrative and Comparative Biology*. **28** (2): 309–18. [doi:10.1093/icb/28.2.309](#). [JSTOR 3883279](#).
26. [^] Pohorecky LA, Wurtman RJ (March 1971). "Adrenocortical control of epinephrine synthesis". *Pharmacological Reviews*. **23** (1): 1–35. [PMID 4941407](#).
27. [^] Wilson JX (1984). "The renin-angiotensin system in nonmammalian vertebrates". *Endocrine Reviews*. **5** (1): 45–61. [doi:10.1210/edrv-5-1-45](#). [PMID 6368215](#).
28. [^] Colombo L, Dalla Valle L, Fiore C, Armanini D, Belvedere P (April 2006). "Aldosterone and the conquest of land". *Journal of Endocrinological Investigation*. **29** (4): 373–9. [doi:10.1007/bf03344112](#). [PMID 16699307](#). [S2CID 25316873](#).

External links
