Intracrine

3 languages

Article

Talk

• Read

• <u>Edit</u>

<u>View history</u>

Tools

From Wikipedia, the free encyclopedia



Illustrations of intracrine, paracrine, autocrine and endocrine

Intracrine refers to a hormone that acts inside a cell, regulating intracellular events. In simple terms it means that the cell stimulates itself by cellular production of a factor that acts within the cell. Steroid hormones act through intracellular (mostly nuclear) receptors and, thus, may be considered to be intracrines. In contrast, peptide or protein hormones, in general, act as endocrines, autocrines, or paracrines by binding to their receptors present on the cell surface. Several peptide/protein hormones or their isoforms also act inside the cell through different mechanisms. These peptide/protein hormones, which have intracellular functions, are also called intracrines. The term 'intracrine' is thought to have been coined to represent peptide/protein hormones that also have intracellular actions. To better understand intracrine, we can compare it to paracrine, autocrine and endocrine. The autocrine system deals with the autocrine receptors of a cell allowing for the hormones to bind. which have been secreted from that same cell. The paracrine system is one where nearby cells get hormones from a cell, and change the functioning of those nearby cells. The endocrine system refers to when the hormones from a cell affect another cell that is very distant from the one that released the hormone.

Paracrine physiology has been understood for decades now and the effects of paracrine hormones have been observed when for example, an obesity associate tumor will face the effects of local adipocytes, even if it is not in direct contact with the fat pads in concern. Endocrine physiology on the other hand is a growing field and has had a new area explored, called intracrinology. In intracrinology, the sex steroids produced locally, exert their action in the same cell where they are produced.

The biological effects produced by intracellular actions are referred as intracrine effects, whereas those produced by binding to cell surface receptors are called endocrine, <u>autocrine</u>, or <u>paracrine</u> effects, depending on the origin of the hormone. The intracrine effect of some of the peptide/protein hormones are similar to their endocrine, autocrine, or paracrine effects; however, these effects are different for some other hormones.

Intracrine can also refer to a hormone acting within the cell that synthesizes it.

Examples of intracrine peptide hormones: There are several protein/peptide hormones that are also intracrines. Notable examples that have been described in the references include:

- Peptides of the renin-angiotensin system: angiotensin II and angiotensin (1-7)
- Fibroblast growth factor 2
- Parathyroid hormone-related protein

See also [edit]

- Local hormone
- Autocrine signalling

References[edit]

- 1. Park, Jiyoung; Euhus, David M.; Scherer, Philipp E. (August 2011). "Paracrine and Endocrine Effects of Adipose Tissue on Cancer Development and Progression". Endocrine Reviews. **32** (4): 550–570. doi:10.1210/er.2010-0030.
- 2. Labrie, Fernand; Luu-The, Van; Labrie, Claude; Bélanger, Alain; Simard, Jacques; Lin, Sheng-Xiang; Pelletier, Georges (April 2003). "Endocrine and Intracrine Sources of Androgens in Women: Inhibition of Breast Cancer and Other Roles of Androgens and Their Precursor Dehydroepiandrosterone". Endocrine Reviews. 24 (2): 152–182. doi:10.1210/er.2001-0031.
- 3. Re, RN (Apr 2003). "The intracrine hypothesis and intracellular peptide hormone action". BioEssays. **25** (4): 401–9. doi:10.1002/bies.10248. PMID 12655647.
- 4. Re RN (Jan 2002). "The origins of intracrine hormone action". American Journal of the Medical Sciences. **323** (1): 43–8. doi:10.1097/00000441-200201000-00008. PMID 11814142.
- 5. Kumar R, Singh VP, Baker KM (July 2007). "The intracellular renin–angiotensin system: a new paradigm". Trends in Endocrinology & Metabolism. **18** (5): 208–214. doi:10.1016/j.tem.2007.05.001. PMID 17509892. S2CID 24041932.
- 6. Fiaschi-Taesch NM, Stewart AF (Feb 2003). "Minireview: parathyroid hormone-related protein as an intracrine factor—trafficking mechanisms and functional consequences". Endocrinology. **144** (2): 407–11. doi:10.1210/en.2002-220818. PMID 12538599.
- 7. Sorensen V, Nilsen T, Wiedlocha A (May 2006). "Functional diversity of FGF-2 isoforms by intracellular sorting". BioEssays. **28** (5): 504–14. doi:10.1002/bies.20405. PMID 16615083. Singh VP, Baker KM, Kumar R (Apr 2008). "Activation of the intracellular renin—angiotensin system in cardiac fibroblasts by high

glucose: role in extracellular matrix production". American Journal of Physiology. Heart and Circulatory Physiology. **294** (4): 1675–84. doi:10.1152/ajpheart.91493.2007. PMID 18296558.

Specific

- 1. <u>^ Park, Jiyoung; Euhus, David M.; Scherer, Philipp E. (August 2011). "Paracrine and Endocrine Effects of Adipose Tissue on Cancer Development and Progression"</u>. Endocrine Reviews. **32** (4): 550–570. doi:10.1210/er.2010-0030. PMC 3369575. PMID 21642230.
- Labrie, Fernand; Luu-The, Van; Labrie, Claude; Bélanger, Alain; Simard, Jacques; Lin, Sheng-Xiang; Pelletier, Georges (April 2003). "Endocrine and Intracrine Sources of Androgens in Women: Inhibition of Breast Cancer and Other Roles of Androgens and Their Precursor
 <u>Dehydroepiandrosterone</u>". Endocrine Reviews. 24 (2): 152–182. doi:10.1210/er.2001-0031. PMID 12700178.