

a) **General 2 2014 Q26c**

Solve the equation $\frac{5x+1}{3} - 4 = 5 - 7x$. 3

b) **General 2 2018 Q28b**

Solve the equation $\frac{2x}{5} + 1 = \frac{3x+1}{2}$, leaving your answer as a fraction. 3

a) **General 2 2014 Q26c**

Solve the equation $\frac{5x+1}{3} - 4 = 5 - 7x$. 3

b) **General 2 2018 Q28b**

Solve the equation $\frac{2x}{5} + 1 = \frac{3x+1}{2}$, leaving your answer as a fraction. 3

Solution

a) General 2 2014 Q26c

Solve the equation $\frac{5x+1}{3} - 4 = 5 - 7x$. 3

b) General 2 2018 Q28b

Solve the equation $\frac{2x}{5} + 1 = \frac{3x+1}{2}$, leaving your answer as a fraction. 3

Solution

a) $x = 1$

a) General 2 2014 Q26c

Solve the equation $\frac{5x+1}{3} - 4 = 5 - 7x$. 3

b) General 2 2018 Q28b

Solve the equation $\frac{2x}{5} + 1 = \frac{3x+1}{2}$, leaving your answer as a fraction. 3

Solution

a) $x = 1$

b) $x = \frac{5}{11}$

Formulae

Standard

MS-A1 Formulae and Equations

updated: 2021-01-21

Learning Outcome

Topic:

Formulae

Syllabus:

- review evaluating the subject of a formula, given the value of other pronumerals in the formula

Activities/Tasks:

- Cambridge Ex 3D Q1-27

Equations

Definition 1

A **formula** is an equation which connects two or more variables. The plural of formula is **formulae** or **formulas**.

Equations

Definition 1

A **formula** is an equation which connects two or more variables. The plural of formula is **formulae** or **formulas**.

Generally a formula is written with a single variable, the **subject** on the LHS and the rest on the right.

$$\underset{\substack{\uparrow \\ \text{subject}}}{A} = \frac{1}{2} ab \cos C$$

Formula substitution

If the formula contains two or more variables and we know the value of all but one of them, we can solve an equation to find the remaining variable.

Formula substitution

If the formula contains two or more variables and we know the value of all but one of them, we can solve an equation to find the remaining variable.

Step 1: State the variables in the formula and the known values.

Formula substitution

If the formula contains two or more variables and we know the value of all but one of them, we can solve an equation to find the remaining variable.

Step 1: State the variables in the formula and the known values.

Step 2: Write down the formula and Substitute the known values into the formula.

Formula substitution

If the formula contains two or more variables and we know the value of all but one of them, we can solve an equation to find the remaining variable.

Step 1: State the variables in the formula and the known values.

Step 2: Write down the formula and Substitute the known values into the formula.

Step 3: Solve the one variable equation to find the unknown value.

Example 1

The area of a trapezium is given by $A = \frac{h}{2}(a + b)$ where a and b are the lengths of the parallel sides and h is the height.

Calculate the height of a 20 cm^2 trapezium with parallel lengths of 3 cm and 7 cm.

Example 1

The area of a trapezium is given by $A = \frac{h}{2}(a + b)$ where a and b are the lengths of the parallel sides and h is the height.

Calculate the height of a 20 cm^2 trapezium with parallel lengths of 3 cm and 7 cm.

Solution

$$A =$$

$$h =$$

$$a =$$

$$b =$$

Example 1

The area of a trapezium is given by $A = \frac{h}{2}(a + b)$ where a and b are the lengths of the parallel sides and h is the height.

Calculate the height of a 20 cm^2 trapezium with parallel lengths of 3 cm and 7 cm.

Solution

$$A = 20 \text{ cm}^2$$

$$h =$$

$$a =$$

$$b =$$

Example 1

The area of a trapezium is given by $A = \frac{h}{2}(a + b)$ where a and b are the lengths of the parallel sides and h is the height.

Calculate the height of a 20 cm^2 trapezium with parallel lengths of 3 cm and 7 cm.

Solution

$$A = 20 \text{ cm}^2$$

$$h = ?$$

$$a =$$

$$b =$$

Example 1

The area of a trapezium is given by $A = \frac{h}{2}(a + b)$ where a and b are the lengths of the parallel sides and h is the height.

Calculate the height of a 20 cm^2 trapezium with parallel lengths of 3 cm and 7 cm.

Solution

$$A = 20 \text{ cm}^2$$

$$h = ?$$

$$a = 3 \text{ cm}$$

$$b =$$

Example 1

The area of a trapezium is given by $A = \frac{h}{2}(a + b)$ where a and b are the lengths of the parallel sides and h is the height.

Calculate the height of a 20 cm^2 trapezium with parallel lengths of 3 cm and 7 cm.

Solution

$$A = 20 \text{ cm}^2$$

$$h = ?$$

$$a = 3 \text{ cm}$$

$$b = 7 \text{ cm}$$

Example 1

The area of a trapezium is given by $A = \frac{h}{2}(a + b)$ where a and b are the lengths of the parallel sides and h is the height.

Calculate the height of a 20 cm^2 trapezium with parallel lengths of 3 cm and 7 cm.

Solution

$$A = 20 \text{ cm}^2$$

$$h = ?$$

$$a = 3 \text{ cm}$$

$$b = 7 \text{ cm}$$

$$A = \frac{h}{2}(a + b)$$

Example 1

The area of a trapezium is given by $A = \frac{h}{2}(a + b)$ where a and b are the lengths of the parallel sides and h is the height.

Calculate the height of a 20 cm^2 trapezium with parallel lengths of 3 cm and 7 cm.

Solution

$$A = 20 \text{ cm}^2$$

$$h = ?$$

$$a = 3 \text{ cm}$$

$$b = 7 \text{ cm}$$

$$A = \frac{h}{2}(a + b)$$

$$20 = \frac{h}{2}(3 + 7)$$

Example 1

The area of a trapezium is given by $A = \frac{h}{2}(a + b)$ where a and b are the lengths of the parallel sides and h is the height.

Calculate the height of a 20 cm^2 trapezium with parallel lengths of 3 cm and 7 cm.

Solution

$$A = 20 \text{ cm}^2$$

$$h = ?$$

$$a = 3 \text{ cm}$$

$$b = 7 \text{ cm}$$

$$A = \frac{h}{2}(a + b)$$

$$20 = \frac{h}{2}(3 + 7)$$

$$20 = \frac{h}{2}(10)$$

Example 1

The area of a trapezium is given by $A = \frac{h}{2}(a + b)$ where a and b are the lengths of the parallel sides and h is the height.

Calculate the height of a 20 cm^2 trapezium with parallel lengths of 3 cm and 7 cm.

Solution

$$A = 20 \text{ cm}^2$$

$$h = ?$$

$$a = 3 \text{ cm}$$

$$b = 7 \text{ cm}$$

$$A = \frac{h}{2}(a + b)$$

$$20 = \frac{h}{2}(3 + 7)$$

$$20 = \frac{h}{2}(10)$$

$$20 = 5h$$

Example 1

The area of a trapezium is given by $A = \frac{h}{2}(a + b)$ where a and b are the lengths of the parallel sides and h is the height.

Calculate the height of a 20 cm^2 trapezium with parallel lengths of 3 cm and 7 cm.

Solution

$$A = 20 \text{ cm}^2$$

$$h = ?$$

$$a = 3 \text{ cm}$$

$$b = 7 \text{ cm}$$

$$A = \frac{h}{2}(a + b)$$

$$20 = \frac{h}{2}(3 + 7)$$

$$20 = \frac{h}{2}(10)$$

$$20 = 5h$$

$$20 \div 5 = 5h \div 5$$

Example 1

The area of a trapezium is given by $A = \frac{h}{2}(a + b)$ where a and b are the lengths of the parallel sides and h is the height.

Calculate the height of a 20 cm^2 trapezium with parallel lengths of 3 cm and 7 cm.

Solution

$$A = 20 \text{ cm}^2$$

$$h = ?$$

$$a = 3 \text{ cm}$$

$$b = 7 \text{ cm}$$

$$A = \frac{h}{2}(a + b)$$

$$20 = \frac{h}{2}(3 + 7)$$

$$20 = \frac{h}{2}(10)$$

$$20 = 5h$$

$$20 \div 5 = 5h \div 5$$

$$4 = h$$

Example 1

The area of a trapezium is given by $A = \frac{h}{2}(a + b)$ where a and b are the lengths of the parallel sides and h is the height.

Calculate the height of a 20 cm^2 trapezium with parallel lengths of 3 cm and 7 cm.

Solution

$$A = 20 \text{ cm}^2$$

$$h = ?$$

$$a = 3 \text{ cm}$$

$$b = 7 \text{ cm}$$

$$A = \frac{h}{2}(a + b)$$

$$20 = \frac{h}{2}(3 + 7)$$

$$20 = \frac{h}{2}(10)$$

$$20 = 5h$$

$$20 \div 5 = 5h \div 5$$

$$4 = h$$

$$h = 4$$

Example 1

The area of a trapezium is given by $A = \frac{h}{2}(a + b)$ where a and b are the lengths of the parallel sides and h is the height.

Calculate the height of a 20 cm^2 trapezium with parallel lengths of 3 cm and 7 cm.

Solution

$$A = 20 \text{ cm}^2$$

$$h = ?$$

$$a = 3 \text{ cm}$$

$$b = 7 \text{ cm}$$

$$A = \frac{h}{2}(a + b)$$

$$20 = \frac{h}{2}(3 + 7)$$

$$20 = \frac{h}{2}(10)$$

$$20 = 5h$$

$$20 \div 5 = 5h \div 5$$

$$4 = h$$

$$h = 4 \text{ cm}$$

The height of the trapezium is equal to 4 cm.

Today's work

- Cambridge Ex 3D Q1-27