

General 2 2018 Q28e

Sophie is driving at 70 km/h. She notices a branch on the road ahead and decides to apply the brakes. Her reaction time is 1.5 seconds. Her braking distance (D metres) is given by $D = 0.01v^2$, where v is speed in km/h.

What is Sophie's stopping distance, to the nearest metre?

General 2 2018 Q28e

Sophie is driving at 70 km/h. She notices a branch on the road ahead and decides to apply the brakes. Her reaction time is 1.5 seconds. Her braking distance (D metres) is given by $D = 0.01v^2$, where v is speed in km/h.

What is Sophie's stopping distance, to the nearest metre?

3

Solution

General 2 2018 Q28e

Sophie is driving at 70 km/h. She notices a branch on the road ahead and decides to apply the brakes. Her reaction time is 1.5 seconds. Her braking distance (D metres) is given by $D = 0.01v^2$, where v is speed in km/h.

What is Sophie's stopping distance, to the nearest metre?

3

Solution

78 m

Blood Alcohol Content

Standard

MS-A1 Formulae and Equations

updated: 2021-01-26

Learning Outcome

Topic:

Blood Alcohol Content

Syllabus:

- calculate and interpret blood alcohol content (BAC) based on drink consumption and body weight
 - use formulae, both in word form and algebraic form, to calculate an estimate for blood alcohol content (BAC), including $BAC_{Male} = \frac{10N7.5H}{6.8M}$ and $BAC_{Female} = \frac{10N7.5H}{5.5M}$ where N is the number of standard drinks consumed, H is the number of hours of drinking, and M is the person's weight in kilograms.
 - determine the number of hours required for a person to stop consuming alcohol in order to reach zero BAC, eg using the formula $time = \frac{BAC}{0.015}$
 - describe the limitations of methods estimating BAC

Activities/Tasks:

- Cambridge Ex 3F Q1-11

Blood Alcohol Content

Blood Alcohol Content

Definition 1

Blood alcohol content (BAC) is a measure of the amount of alcohol in your blood.

Blood Alcohol Content

Definition 1

Blood alcohol content (BAC) is a measure of the amount of alcohol in your blood.

BAC tells you the number of grams of alcohol per 100 mL of blood.

Blood Alcohol Content

Definition 1

Blood alcohol content (BAC) is a measure of the amount of alcohol in your blood.

BAC tells you the number of grams of alcohol per 100 mL of blood.

For example a BAC of 0.10 means there is 0.10 g of alcohol for every 100 mL of blood, or 1 g for every 1 L of blood.

Estimating BAC

BAC can be estimated using the following formulae:

Formula

$$BAC_{\text{Male}} = \frac{10N - 7.5H}{6.8M} \quad \text{or} \quad BAC_{\text{Female}} = \frac{10N - 7.5H}{5.5M}$$

Where:

BAC - Blood alcohol content

Estimating BAC

BAC can be estimated using the following formulae:

Formula

$$BAC_{\text{Male}} = \frac{10N - 7.5H}{6.8M} \quad \text{or} \quad BAC_{\text{Female}} = \frac{10N - 7.5H}{5.5M}$$

Where:

BAC - Blood alcohol content

N - Number of standard
drinks consumed

Estimating BAC

BAC can be estimated using the following formulae:

Formula

$$BAC_{\text{Male}} = \frac{10N - 7.5H}{6.8M} \quad \text{or} \quad BAC_{\text{Female}} = \frac{10N - 7.5H}{5.5M}$$

Where:

BAC - Blood alcohol content

N - Number of standard
drinks consumed

H - Hours drinking

Estimating BAC

BAC can be estimated using the following formulae:

Formula

$$BAC_{\text{Male}} = \frac{10N - 7.5H}{6.8M} \quad \text{or} \quad BAC_{\text{Female}} = \frac{10N - 7.5H}{5.5M}$$

Where:

BAC - Blood alcohol content

N - Number of standard
drinks consumed

H - Hours drinking

M - Mass in kilograms

Important Note

These formulae are **not** on the reference sheet.

You are **not** expected to remember these formulae.

Limitations

These formulae are only good for calculating *estimates* of a person BAC.

Limitations

These formulae are only good for calculating *estimates* of a person BAC.

The formulae for estimating BAC only takes into account the number of standard drinks consumed, sex, the amount of time, and mass.

Limitations

These formulae are only good for calculating *estimates* of a person BAC.

The formulae for estimating BAC only takes into account the number of standard drinks consumed, sex, the amount of time, and mass. There are other factors that influence BAC including:

- Fitness
- Health
- Liver function
- Food in the stomach
- Medications

Example 1

Osman is 87 kg and has consumed 3 standard drinks in the past hour. Estimate Osman's BAC to 3 decimal places using the following formula where N represents the number of standard drinks, H is the number of hours drinking and M is his mass in kilograms.

$$BAC_{\text{Male}} = \frac{10N - 7.5H}{6.8M}$$

Example 1

Osman is 87 kg and has consumed 3 standard drinks in the past hour. Estimate Osman's BAC to 3 decimal places using the following formula where N represents the number of standard drinks, H is the number of hours drinking and M is his mass in kilograms.

$$BAC_{\text{Male}} = \frac{10N - 7.5H}{6.8M}$$

Solution

$$BAC_{\text{Male}} = \frac{10N - 7.5H}{6.8M}$$

Example 1

Osman is 87 kg and has consumed 3 standard drinks in the past hour. Estimate Osman's BAC to 3 decimal places using the following formula where N represents the number of standard drinks, H is the number of hours drinking and M is his mass in kilograms.

$$BAC_{\text{Male}} = \frac{10N - 7.5H}{6.8M}$$

Solution

$$\begin{aligned} BAC_{\text{Male}} &= \frac{10N - 7.5H}{6.8M} \\ &= \frac{10(3) - 7.5(1)}{6.8(87)} \end{aligned}$$

Example 1

Osman is 87 kg and has consumed 3 standard drinks in the past hour. Estimate Osman's BAC to 3 decimal places using the following formula where N represents the number of standard drinks, H is the number of hours drinking and M is his mass in kilograms.

$$BAC_{\text{Male}} = \frac{10N - 7.5H}{6.8M}$$

Solution

$$\begin{aligned} BAC_{\text{Male}} &= \frac{10N - 7.5H}{6.8M} \\ &= \frac{10(3) - 7.5(1)}{6.8(87)} \\ &= 0.038032... \end{aligned}$$

Example 1

Osman is 87 kg and has consumed 3 standard drinks in the past hour. Estimate Osman's BAC to 3 decimal places using the following formula where N represents the number of standard drinks, H is the number of hours drinking and M is his mass in kilograms.

$$BAC_{\text{Male}} = \frac{10N - 7.5H}{6.8M}$$

Solution

$$\begin{aligned} BAC_{\text{Male}} &= \frac{10N - 7.5H}{6.8M} \\ &= \frac{10(3) - 7.5(1)}{6.8(87)} \\ &= 0.038032... \\ &\approx 0.038 \end{aligned}$$

Osman's BAC is estimated to be 0.038.

Reaching zero BAC

The number of hours required for a person to reach zero BAC after they stop consuming alcohol is given by the following formula:

Reaching zero BAC

The number of hours required for a person to reach zero BAC after they stop consuming alcohol is given by the following formula:

Formula

$$\text{Time (h)} = \frac{BAC}{0.015}$$

Reaching zero BAC

The number of hours required for a person to reach zero BAC after they stop consuming alcohol is given by the following formula:

Formula

$$\text{Time (h)} = \frac{BAC}{0.015}$$

Important Note

This formula is **not** on the reference sheet.
You are **not** expected to remember this formula.

Example 2

Fiona has a BAC of 0.027. Use the following formula to estimate the time she will need for her BAC to reach zero. Answer to the nearest minute.

$$\text{Time} = \frac{BAC}{0.015}$$

Example 2

Fiona has a BAC of 0.027. Use the following formula to estimate the time she will need for her BAC to reach zero. Answer to the nearest minute.

$$\text{Time} = \frac{BAC}{0.015}$$

Solution

$$\text{Time} = \frac{BAC}{0.015}$$

Example 2

Fiona has a BAC of 0.027. Use the following formula to estimate the time she will need for her BAC to reach zero. Answer to the nearest minute.

$$\text{Time} = \frac{BAC}{0.015}$$

Solution

$$\begin{aligned}\text{Time} &= \frac{BAC}{0.015} \\ &= \frac{(0.027)}{0.015}\end{aligned}$$

Example 2

Fiona has a BAC of 0.027. Use the following formula to estimate the time she will need for her BAC to reach zero. Answer to the nearest minute.

$$\text{Time} = \frac{BAC}{0.015}$$

Solution

$$\begin{aligned}\text{Time} &= \frac{BAC}{0.015} \\ &= \frac{(0.027)}{0.015} \\ &= 1.8\end{aligned}$$

Example 2

Fiona has a BAC of 0.027. Use the following formula to estimate the time she will need for her BAC to reach zero. Answer to the nearest minute.

$$\text{Time} = \frac{BAC}{0.015}$$

Solution

$$\begin{aligned}\text{Time} &= \frac{BAC}{0.015} \\ &= \frac{(0.027)}{0.015} \\ &= 1.8\end{aligned}$$



Example 2

Fiona has a BAC of 0.027. Use the following formula to estimate the time she will need for her BAC to reach zero. Answer to the nearest minute.

$$\text{Time} = \frac{BAC}{0.015}$$

Solution

$$\begin{aligned}\text{Time} &= \frac{BAC}{0.015} \\ &= \frac{(0.027)}{0.015} \\ &= 1.8 \\ &\quad \boxed{0.999} \\ &= 1^{\circ}48'0''\end{aligned}$$

Example 2

Fiona has a BAC of 0.027. Use the following formula to estimate the time she will need for her BAC to reach zero. Answer to the nearest minute.

$$\text{Time} = \frac{BAC}{0.015}$$

Solution

$$\begin{aligned}\text{Time} &= \frac{BAC}{0.015} \\ &= \frac{(0.027)}{0.015} \\ &= 1.8 \\ &\quad \boxed{0.999} \\ &= 1^{\circ}48'0''\end{aligned}$$

Fiona needs to wait 1 hour and 48 minutes.

Today's work

- Cambridge Ex 3F Q1-11