

Busemann Functions in the Wasserstein Space: Existence, Closed-Forms and Applications

Clément Bonet¹, Elsa Cazelles², Lucas Drumetz³, Nicolas Courty⁴

¹Ecole Polytechnique, CMAP, Institut Polytechnique de Paris

²CNRS, Université de Toulouse, IRIT

³IMT Atlantique, Lab-STICC

⁴Université Bretagne Sud, IRISA

Journée Titouan

17/12/2025



Motivation

Busemann function:

- Major tool in geometry (Busemann, 1955; Bridson and Haefliger, 2013)
- Many applications in Machine Learning (Chami et al., 2021; Ghadimi Atigh et al., 2021; Bonet et al., 2023; Berg et al., 2024)
 - PCA, Classification, Projections...
- Applications restricted to finite dimensional spaces

Space of probability distributions:

- Datasets of Distributions
 - Documents (Kusner et al., 2015), point clouds (Geuter et al., 2025), images (Rubner et al., 2000), single-cells (Bellazzi et al., 2021)
 - Labeled distributions (Alvarez-Melis and Fusi, 2020), Gaussian mixtures (Delon and Desolneux, 2020)...
- Rich geometry with Optimal Transport (Ambrosio et al., 2008)

Table of Contents

Busemann Function in Metric Spaces

Busemann Functions in Wasserstein Space

Comparison of Labeled Datasets

Geodesic Metric space (Bridson and Haefliger, 2013)



Let (X, d) be a metric space.

- Let $x, y \in X$. A continuous map $\gamma : [0, 1] \rightarrow X$ is a (constant-speed) geodesic between x and y if

$$\begin{cases} \gamma(0) = x \\ \gamma(1) = y \\ \forall t, s \in [0, 1], d(\gamma(t), \gamma(s)) = |t - s|d(x, y) \end{cases}$$


→ γ minimizes the length between x and y

- (X, d) is a geodesic metric space if any two points are joined by a geodesic



$$X = \mathbb{R}^d, d(x, y) = \|x - y\|_2, \quad \gamma(t) = (1 - t)x + ty$$

$$X = S^{d-1}$$



$$X = \mathcal{M}, \gamma(t) = \exp_x(tv)$$

Geodesic Lines and Rays


Let $\gamma : [0, 1] \rightarrow X$ be a geodesic, $\kappa = d(\gamma(0), \gamma(1))$ its speed.

- **Geodesic line:** extension of γ to \mathbb{R} such that


$$\forall t, s \in \mathbb{R}, d(\gamma(t), \gamma(s)) = \kappa|t - s|$$

- **Geodesic ray:** extension of γ to $[0, +\infty[$ such that

$$\forall t, s \in [0, +\infty[, d(\gamma(t), \gamma(s)) = \kappa|t - s|$$



$$\forall s, t \in \mathbb{R}, \gamma(t) = (1-t)x + ty$$



No geodesic ray or line

Sufficient Conditions for Geodesic Rays?

- Curvature $\leq 0 \iff$ for all $x \in X, t \in [0, 1]$,


$$d^2(x, \gamma(t)) \leq (1-t)d^2(x, \gamma(0)) + td^2(x, \gamma(1)) - t(1-t)d^2(\gamma(0), \gamma(1))$$

→ geodesics are **geodesics lines**


- Curvature $> 0 \iff$ for all $x \in X, t \in [0, 1]$,

$$d^2(x, \gamma(t)) \geq (1-t)d^2(x, \gamma(0)) + td^2(x, \gamma(1)) - t(1-t)d^2(\gamma(0), \gamma(1))$$


→ **no guarantees**



Negatively curved



No curvature



Positively curved

Examples of Non-Positively Curved Spaces


- **Euclidean spaces** $(\mathbb{R}^d, \|\cdot\|_2)$: $\forall x \in \mathbb{R}^d$, (parallelogram rule)

$$\|x - \gamma(t)\|_2^2 = (1-t)\|x - \gamma(0)\|_2^2 + t\|x - \gamma(1)\|_2^2 - t(1-t)\|\gamma(0) - \gamma(1)\|_2^2$$

- **Hyperbolic spaces**: $\mathbb{L}^d = \{(x_0, \dots, x_d) \in \mathbb{R}^{d+1}, \langle x, x \rangle_{\mathbb{L}} = -1, x_0 > 0\}$,

$$d_{\mathbb{L}}(x, y) = \operatorname{arccosh}(-\langle x, y \rangle_{\mathbb{L}}), \quad \langle x, y \rangle_{\mathbb{L}} = -x_0y_0 + \sum_{i=1}^d x_iy_i$$

Geodesic: $\forall t \in \mathbb{R}, \gamma(t) = \cosh(\|v\|_{\mathbb{L}})x + \sinh(t\|v\|_{\mathbb{L}})\frac{v}{\|v\|_{\mathbb{L}}}$
 \rightarrow (constant) negative curvature



Busemann Function

Let (X, d) be a geodesic metric space and $\gamma : [0, +\infty[\rightarrow X$ be a geodesic ray.

Busemann function associated to γ :

$$\begin{aligned}\forall x \in X, \quad B^\gamma(x) &= \lim_{t \rightarrow +\infty} d(x, \gamma(t)) - d(\gamma(0), \gamma(t)) \\ &= \lim_{t \rightarrow +\infty} d(x, \gamma(t)) - td(\gamma(0), \gamma(1))\end{aligned}$$

Busemann Function

Let (X, d) be a geodesic metric space and $\gamma : [0, +\infty[\rightarrow X$ be a geodesic ray.

Busemann function associated to γ :

$$\begin{aligned}\forall x \in X, B^\gamma(x) &= \lim_{t \rightarrow +\infty} d(x, \gamma(t)) - d(\gamma(0), \gamma(t)) \\ &= \lim_{t \rightarrow +\infty} d(x, \gamma(t)) - td(\gamma(0), \gamma(1))\end{aligned}$$

For $X = \mathbb{R}^d$, $\gamma(t) = x_0 + tv$, $x_0, v \in \mathbb{R}^d$,

$$\begin{aligned}d(x, \gamma(t)) - d(\gamma(0), \gamma(t)) &= \|x - x_0 - tv\|_2 - \|x_0 - x_0 - tv\|_2 \\ &= t\|v\|_2 \sqrt{1 - \frac{2}{t\|v\|_2^2} \langle v, x - x_0 \rangle + o(t^{-1})} - t\|v\|_2 \\ &= t\|v\|_2 \left(1 - \frac{1}{t\|v\|_2^2} \langle v, x - x_0 \rangle + o(t^{-1})\right) - t\|v\|_2 \\ &= - \left\langle x - x_0, \frac{v}{\|v\|_2} \right\rangle + o(1)\end{aligned}$$

$$\rightarrow B^\gamma(x) = - \left\langle x - x_0, \frac{v}{\|v\|_2} \right\rangle$$

Busemann Function on \mathbb{R}^d

On \mathbb{R}^d : for $\gamma(t) = x_0 + tv$ with $x_0, v \in \mathbb{R}^d$,


$$\forall x \in \mathbb{R}^d, B^\gamma(x) = - \left\langle x - x_0, \frac{v}{\|v\|_2} \right\rangle$$

- $B^\gamma(x)$: coincides with the coordinate of the geodesic projection (up to a sign)

$$t = \underset{s \in \mathbb{R}}{\operatorname{argmin}} \|x - (x_0 + sv)\|_2^2 = \left\langle x - x_0, \frac{v}{\|v\|_2} \right\rangle$$

→ Projection: $\gamma(-B^\gamma(x))$

- Level sets of B^γ are hyperplanes orthogonal to v



Busemann Function on \mathbb{R}^d

On \mathbb{R}^d : for $\gamma(t) = x_0 + tv$ with $x_0, v \in \mathbb{R}^d$,

$$\forall x \in \mathbb{R}^d, B^\gamma(x) = -\left\langle x - x_0, \frac{v}{\|v\|_2} \right\rangle$$


- $B^\gamma(x)$: coincides with the coordinate of the geodesic projection (up to a sign)

$$t = \underset{s \in \mathbb{R}}{\operatorname{argmin}} \|x - (x_0 + sv)\|_2^2 = \left\langle x - x_0, \frac{v}{\|v\|_2} \right\rangle$$

→ Projection: $\gamma(-B^\gamma(x))$

- Level sets of B^γ are hyperplanes orthogonal to v

→ **What about on other spaces?**



Busemann Function on Hyperbolic Space


Let $x^0 = (1, 0, \dots, 0) \in \mathbb{R}^{d+1}$, $v \in T_{x^0} \mathbb{L}^d \cap S^d$,

$$\forall x \in \mathbb{L}^d, B^\gamma(x) = \log(-\langle x, x^0 + v \rangle_{\mathbb{L}})$$


→ different from geodesic projection (Bonet et al., 2023)

Horospheres: Levels sets of B^γ : $(B^\gamma)^{-1}(\{t\})$ for all $t \in \mathbb{R}$

→ Second generalization of hyperplanes




Projections along geodesics
submanifolds




Projections along
horospheres

Applications in Machine Learning

- Busemann used as a projection
 - HoroPCA ([Chami et al., 2021](#))
→ Project on a geodesic subspace along horospheres



(a) PGA (average distortion: 0.534)




(b) HOROPCA (average distortion: 0.078)

From ([Chami et al., 2021](#))


Applications in Machine Learning

- Busemann used as a projection
 - HoroPCA ([Chami et al., 2021](#))
→ Project on a geodesic subspace along horospheres
 - Sliced-Wasserstein on Cartan-Hadamard manifolds ([Bonet et al., 2023, 2025a](#))
→ Project on geodesics with the Busemann function




Applications in Machine Learning

- Busemann used as a projection
 - HoroPCA (Chami et al., 2021)
→ Project on a geodesic subspace along horospheres
 - Sliced-Wasserstein on Cartan-Hadamard manifolds (Bonet et al., 2023, 2025a)
→ Project on geodesics with the Busemann function
- Busemann used for alignments
→ Embedding with prototypes (Ghadimi Atigh et al., 2021)



Applications in Machine Learning

- Busemann used as a projection
 - HoroPCA ([Chami et al., 2021](#))
→ Project on a geodesic subspace along horospheres
 - Sliced-Wasserstein on Cartan-Hadamard manifolds ([Bonet et al., 2023, 2025a](#))
→ Project on geodesics with the Busemann function
- Busemann used for alignments
→ Embedding with prototypes ([Ghadimi Atigh et al., 2021](#))
- Busemann used for classification ([Fan et al., 2023; Doorenbos et al., 2024; Berg et al., 2024, 2025](#))
→ SVM, Random Forests, Logistic Regression...



From ([Berg et al., 2024](#))

Table of Contents

Busemann Function in Metric Spaces


Busemann Functions in Wasserstein Space

Comparison of Labeled Datasets

Datasets of Distributions

Examples

- Histograms (e.g. age distributions of countries, financial assets...)
- Documents: distributions of words (Kusner et al., 2015)
- Cells: distributions of genes (Bellazzi et al., 2021)
- Embedding of words in Gaussian distributions (Vilnis and McCallum, 2015)



Wasserstein Geometry (Ambrosio et al., 2008)

Definition (Wasserstein distance)

Let $\mu, \nu \in \mathcal{P}_2(\mathbb{R}^d)$ and denote by $\Pi(\mu, \nu)$ the set of coupling between μ, ν . Then, the Wasserstein distance is

$$W_2^2(\mu, \nu) = \inf_{\gamma \in \Pi(\mu, \nu)} \int \|x - y\|_2^2 d\gamma(x, y),$$

with $\Pi(\mu, \nu) = \{\gamma \in \mathcal{P}_2(\mathbb{R}^d \times \mathbb{R}^d), \pi_1^* \gamma = \mu, \pi_2^* \gamma = \nu\}$, $\pi^1 : (x, y) \mapsto x$, $\pi^2 : (x, y) \mapsto y$

Remainder: For $T : \mathbb{R}^d \rightarrow \mathbb{R}^p$ measurable, $X \sim \mu \implies T(X) \sim T_\# \mu$

Wasserstein Geometry (Ambrosio et al., 2008)

Definition (Wasserstein distance)

Let $\mu, \nu \in \mathcal{P}_2(\mathbb{R}^d)$ and denote by $\Pi(\mu, \nu)$ the set of coupling between μ, ν . Then, the Wasserstein distance is

$$W_2^2(\mu, \nu) = \inf_{\gamma \in \Pi(\mu, \nu)} \int \|x - y\|_2^2 d\gamma(x, y),$$

with $\Pi(\mu, \nu) = \{\gamma \in \mathcal{P}_2(\mathbb{R}^d \times \mathbb{R}^d), \pi_1^\# \gamma = \mu, \pi_2^\# \gamma = \nu\}$, $\pi^1 : (x, y) \mapsto x$, $\pi^2 : (x, y) \mapsto y$

Remainder: For $T : \mathbb{R}^d \rightarrow \mathbb{R}^p$ measurable, $X \sim \mu \implies T(X) \sim T_\# \mu$

Properties:

- W_2 distance, $(\mathcal{P}_2(\mathbb{R}^d), W_2)$: Wasserstein space
- Riemannian structure
- Geodesic metric space


Geodesics in Wasserstein Space

Let $\Pi_o(\mu, \nu) = \{\gamma, \gamma \in \operatorname{argmin}_{\gamma \in \Pi(\mu, \nu)} \int \|x - y\|_2^2 d\gamma(x, y)\}$

Geodesics

For $\mu_0, \mu_1 \in \mathcal{P}_2(\mathbb{R}^d)$, geodesic = displacement interpolation:

$$\forall t \in [0, 1], \mu_t = ((1-t)\pi^1 + t\pi^2)_\# \gamma \quad \text{for } \gamma \in \Pi_o(\mu_0, \mu_1)$$



Geodesics in Wasserstein Space

Let $\Pi_o(\mu, \nu) = \{\gamma, \gamma \in \operatorname{argmin}_{\gamma \in \Pi(\mu, \nu)} \int \|x - y\|_2^2 d\gamma(x, y)\}$

Geodesics

For $\mu_0, \mu_1 \in \mathcal{P}_2(\mathbb{R}^d)$, geodesic = displacement interpolation:

$$\forall t \in [0, 1], \mu_t = ((1-t)\pi^1 + t\pi^2)_\# \gamma \quad \text{for } \gamma \in \Pi_o(\mu_0, \mu_1)$$

For all $\nu \in \mathcal{P}_2(\mathbb{R}^d)$, $t \in [0, 1]$, (Ambrosio et al., 2008)

$$W_2^2(\mu_t, \nu) \geq (1-t)W_2^2(\mu_0, \nu) + tW_2^2(\mu_1, \nu) - t(1-t)W_2^2(\mu_0, \mu_1)$$

→ Positively curved space (PC Space)

Geodesics in Wasserstein Space

Let $\Pi_o(\mu, \nu) = \{\gamma, \gamma \in \operatorname{argmin}_{\gamma \in \Pi(\mu, \nu)} \int \|x - y\|_2^2 d\gamma(x, y)\}$

Geodesics

For $\mu_0, \mu_1 \in \mathcal{P}_2(\mathbb{R}^d)$, geodesic = displacement interpolation:

$$\forall t \in [0, 1], \mu_t = ((1-t)\pi^1 + t\pi^2)_\# \gamma \quad \text{for } \gamma \in \Pi_o(\mu_0, \mu_1)$$

For all $\nu \in \mathcal{P}_2(\mathbb{R}^d)$, $t \in [0, 1]$, (Ambrosio et al., 2008)

$$W_2^2(\mu_t, \nu) \geq (1-t)W_2^2(\mu_0, \nu) + tW_2^2(\mu_1, \nu) - t(1-t)W_2^2(\mu_0, \mu_1)$$

→ Positively curved space (PC Space)

By (Zhu et al., 2021): always at least one geodesic ray starting from μ_0

→ Conditions to extend $t \mapsto \mu_t$ to \mathbb{R}_+ ?

1D Wasserstein Space

Let $\mu, \nu \in \mathcal{P}_2(\mathbb{R})$,

- Cumulative distribution function:

$$\forall t \in \mathbb{R}, F_\mu(t) = \mu([-\infty, t]) = \int \mathbb{1}_{]-\infty, t]}(x) d\mu(x)$$

- Quantile function:

$$\forall u \in [0, 1], F_\mu^{-1}(u) = \inf \{x \in \mathbb{R}, F_\mu(x) \geq u\}$$

1D Wasserstein Distance

The optimal coupling is $\gamma^* = (F_\mu^{-1}, F_\nu^{-1})_{\#} \text{Unif}([0, 1])$, and

$$W_2^2(\mu, \nu) = \int_0^1 |F_\mu^{-1}(u) - F_\nu^{-1}(u)|^2 du = \|F_\mu^{-1} - F_\nu^{-1}\|_{L^2([0,1])}^2$$

1D Wasserstein Space

Let $\mu, \nu \in \mathcal{P}_2(\mathbb{R})$,

- Cumulative distribution function:

$$\forall t \in \mathbb{R}, F_\mu(t) = \mu([-\infty, t]) = \int \mathbb{1}_{]-\infty, t]}(x) d\mu(x)$$

- Quantile function:

$$\forall u \in [0, 1], F_\mu^{-1}(u) = \inf \{x \in \mathbb{R}, F_\mu(x) \geq u\}$$

1D Wasserstein Distance

The optimal coupling is $\gamma^* = (F_\mu^{-1}, F_\nu^{-1})_{\#} \text{Unif}([0, 1])$, and

$$W_2^2(\mu, \nu) = \int_0^1 |F_\mu^{-1}(u) - F_\nu^{-1}(u)|^2 du = \|F_\mu^{-1} - F_\nu^{-1}\|_{L^2([0,1])}^2$$

Geodesic between μ and ν : $\forall t \in [0, 1]$,

$$\mu_t = ((1-t)\pi^1 + t\pi^2)_{\#} \gamma^* = ((1-t)F_\mu^{-1} + tF_\nu^{-1})_{\#} \text{Unif}([0, 1])$$

$$\rightarrow F_t^{-1} = (1-t)F_\mu^{-1} + tF_\nu^{-1}$$

Geodesic Rays in 1D Wasserstein Space

For $\mu_0, \mu_1 \in \mathcal{P}_2(\mathbb{R})$, quantile F_t^{-1} of the geodesic $t \mapsto \mu_t$ characterized as

$$\forall t \in [0, 1], \quad F_t^{-1} = (1 - t)F_0^{-1} + tF_1^{-1} = F_0^{-1} + t(F_1^{-1} - F_0^{-1}).$$

- For all $t, s \in \mathbb{R}$,

$$\|F_t^{-1} - F_s^{-1}\|_{L^2([0,1])}^2 = (t - s)^2 \|F_0^{-1} - F_1^{-1}\|_{L^2([0,1])}^2$$

→ ok if F_t^{-1}, F_s^{-1} quantile functions, i.e. non-decreasing and left-continuous

Geodesic Rays in 1D Wasserstein Space

For $\mu_0, \mu_1 \in \mathcal{P}_2(\mathbb{R})$, quantile F_t^{-1} of the geodesic $t \mapsto \mu_t$ characterized as

$$\forall t \in [0, 1], \quad F_t^{-1} = (1-t)F_0^{-1} + tF_1^{-1} = F_0^{-1} + t(F_1^{-1} - F_0^{-1}).$$

- For all $t, s \in \mathbb{R}$,

$$\|F_t^{-1} - F_s^{-1}\|_{L^2([0,1])}^2 = (t-s)^2 \|F_0^{-1} - F_1^{-1}\|_{L^2([0,1])}^2$$

→ ok if F_t^{-1}, F_s^{-1} quantile functions, i.e. non-decreasing and left-continuous

- For all $t \geq 0, 0 < m < m' < 1$,

$$\begin{aligned} F_t^{-1}(m) - F_t^{-1}(m') &= F_0^{-1}(m) - F_0^{-1}(m') \\ &\quad + t(F_1^{-1}(m) - F_0^{-1}(m) - (F_1^{-1}(m') - F_0^{-1}(m'))) \leq 0 \end{aligned}$$

Proposition (Kloeckner, 2010)

$t \mapsto \mu_t$ is a geodesic ray if and only if $F_1^{-1} - F_0^{-1}$ is non-decreasing.

Illustrations - 1D Gaussian Case

Let $\mu_0 = \mathcal{N}(m_0, \sigma_0^2)$, $\mu_1 = \mathcal{N}(m_1, \sigma_1^2)$, ϕ^{-1} quantile function of $\mathcal{N}(0, 1)$.

$$\rightarrow F_0^{-1} = m_0 + \sigma_0 \phi^{-1}, F_1^{-1} = m_1 + \sigma_1 \phi^{-1}$$

$$\rightarrow \forall t \in [0, 1], \mu_t = \mathcal{N}((1-t)m_0 + tm_1, ((1-t)\sigma_0 + t\sigma_1)^2)$$

Illustrations - 1D Gaussian Case

Let $\mu_0 = \mathcal{N}(m_0, \sigma_0^2)$, $\mu_1 = \mathcal{N}(m_1, \sigma_1^2)$, ϕ^{-1} quantile function of $\mathcal{N}(0, 1)$.

$$\rightarrow F_0^{-1} = m_0 + \sigma_0 \phi^{-1}, F_1^{-1} = m_1 + \sigma_1 \phi^{-1}$$

$$\rightarrow \forall t \in [0, 1], \mu_t = \mathcal{N}\left((1-t)m_0 + tm_1, ((1-t)\sigma_0 + t\sigma_1)^2\right)$$

For all $0 < m < m' < 1$,

$$((F_1^{-1} - F_0^{-1})(m') - (F_1^{-1} - F_0^{-1})(m)) = (\sigma_1 - \sigma_0)(\phi^{-1}(m') - \phi^{-1}(m)) \geq 0$$

$$\iff \sigma_1 \geq \sigma_0$$

$t \mapsto \mu_t$ is a geodesic ray if and only if $\sigma_1 \geq \sigma_0$.

Illustrations - 1D Gaussian Case

Let $\mu_0 = \mathcal{N}(m_0, \sigma_0^2)$, $\mu_1 = \mathcal{N}(m_1, \sigma_1^2)$, ϕ^{-1} quantile function of $\mathcal{N}(0, 1)$.


$$\rightarrow F_0^{-1} = m_0 + \sigma_0 \phi^{-1}, F_1^{-1} = m_1 + \sigma_1 \phi^{-1}$$

$$\rightarrow \forall t \in [0, 1], \mu_t = \mathcal{N}((1-t)m_0 + tm_1, ((1-t)\sigma_0 + t\sigma_1)^2)$$

For all $0 < m < m' < 1$,

$$((F_1^{-1} - F_0^{-1})(m') - (F_1^{-1} - F_0^{-1})(m)) = (\sigma_1 - \sigma_0)(\phi^{-1}(m') - \phi^{-1}(m)) \geq 0 \\ \iff \sigma_1 \geq \sigma_0$$

$t \mapsto \mu_t$ is a geodesic ray if and only if $\sigma_1 \geq \sigma_0$.



Illustrations - 1D Gaussian Case

Let $\mu_0 = \mathcal{N}(m_0, \sigma_0^2)$, $\mu_1 = \mathcal{N}(m_1, \sigma_1^2)$, ϕ^{-1} quantile function of $\mathcal{N}(0, 1)$.


$$\rightarrow F_0^{-1} = m_0 + \sigma_0 \phi^{-1}, F_1^{-1} = m_1 + \sigma_1 \phi^{-1}$$

$$\rightarrow \forall t \in [0, 1], \mu_t = \mathcal{N}\left((1-t)m_0 + tm_1, ((1-t)\sigma_0 + t\sigma_1)^2\right)$$

For all $0 < m < m' < 1$,

$$\begin{aligned} ((F_1^{-1} - F_0^{-1})(m') - (F_1^{-1} - F_0^{-1})(m)) &= (\sigma_1 - \sigma_0)(\phi^{-1}(m') - \phi^{-1}(m)) \geq 0 \\ \iff \sigma_1 &\geq \sigma_0 \end{aligned}$$

$t \mapsto \mu_t$ is a geodesic ray if and only if $\sigma_1 \geq \sigma_0$.



Illustrations - Starting from a Dirac

Let $x_0 \in \mathbb{R}$, $\mu_0 = \delta_{x_0}$, $F_0^{-1}(p) = x_0$ for all $1 \geq p > 0$.

→ For all $\mu_1 \in \mathcal{P}_2(\mathbb{R})$, $F_1^{-1} - F_0^{-1}$ non-decreasing

→ $\gamma^* = \mu_0 \otimes \mu_1$, $\mu_t = ((1-t)x_0 + t\text{Id})_\# \mu_1$

For all $\mu_1 \in \mathcal{P}_2(\mathbb{R})$, $t \mapsto \mu_t$ is a geodesic ray.



Illustrations - Starting from a Dirac

Let $x_0 \in \mathbb{R}$, $\mu_0 = \delta_{x_0}$, $F_0^{-1}(p) = x_0$ for all $1 \geq p > 0$.

→ For all $\mu_1 \in \mathcal{P}_2(\mathbb{R})$, $F_1^{-1} - F_0^{-1}$ non-decreasing

→ $\gamma^* = \mu_0 \otimes \mu_1$, $\mu_t = ((1-t)x_0 + t\text{Id})_\# \mu_1$

For all $\mu_1 \in \mathcal{P}_2(\mathbb{R})$, $t \mapsto \mu_t$ is a geodesic ray.



Illustrations - Starting from a Dirac

Let $x_0 \in \mathbb{R}$, $\mu_0 = \delta_{x_0}$, $F_0^{-1}(p) = x_0$ for all $1 \geq p > 0$.

→ For all $\mu_1 \in \mathcal{P}_2(\mathbb{R})$, $F_1^{-1} - F_0^{-1}$ non-decreasing

→ $\gamma^* = \mu_0 \otimes \mu_1$, $\mu_t = ((1-t)x_0 + t\text{Id})_\# \mu_1$

For all $\mu_1 \in \mathcal{P}_2(\mathbb{R})$, $t \mapsto \mu_t$ is a geodesic ray.

→ extends to \mathbb{R}^d (Bertrand and Kloeckner, 2016, Lemma 2.1)




Illustration - Discrete Distributions

Let $x_1 < \dots < x_n$, $y_1 < \dots < y_n$, $\mu_0 = \frac{1}{n} \sum_{i=1}^n \delta_{x_i}$, $\mu_1 = \frac{1}{n} \sum_{i=1}^n \delta_{y_i}$.

$t \mapsto \mu_t$ is a geodesic ray if and only if for all $j > i$, $y_i - x_i \leq y_j - x_j$.

Illustration - Discrete Distributions

Let $x_1 < \dots < x_n$, $y_1 < \dots < y_n$, $\mu_0 = \frac{1}{n} \sum_{i=1}^n \delta_{x_i}$, $\mu_1 = \frac{1}{n} \sum_{i=1}^n \delta_{y_i}$.

$t \mapsto \mu_t$ is a geodesic ray if and only if for all $j > i$, $y_i - x_i \leq y_j - x_j$.

$$\mu_0 = \frac{1}{2}\delta_{-2} + \frac{1}{2}\delta_0, \quad \mu_1 = \frac{1}{2}\delta_{-1} + \frac{1}{2}\delta_{1.5}$$

$$\rightarrow y_1 - x_1 = 1 < y_2 - x_2 = 1.5$$




Illustration - Discrete Distributions

Let $x_1 < \dots < x_n$, $y_1 < \dots < y_n$, $\mu_0 = \frac{1}{n} \sum_{i=1}^n \delta_{x_i}$, $\mu_1 = \frac{1}{n} \sum_{i=1}^n \delta_{y_i}$.

$t \mapsto \mu_t$ is a geodesic ray if and only if for all $j > i$, $y_i - x_i \leq y_j - x_j$.

$$\mu_0 = \frac{1}{2}\delta_{-2} + \frac{1}{2}\delta_0, \quad \mu_1 = \frac{1}{2}\delta_{-1} + \frac{1}{2}\delta_{1.5}$$

$$\rightarrow y_1 - x_1 = 1 < y_2 - x_2 = 1.5$$




Illustration - Discrete Distributions

Let $x_1 < \dots < x_n$, $y_1 < \dots < y_n$, $\mu_0 = \frac{1}{n} \sum_{i=1}^n \delta_{x_i}$, $\mu_1 = \frac{1}{n} \sum_{i=1}^n \delta_{y_i}$.

$t \mapsto \mu_t$ is a geodesic ray if and only if for all $j > i$, $y_i - x_i \leq y_j - x_j$.

$$\mu_0 = \frac{1}{2}\delta_0 + \frac{1}{2}\delta_4, \quad \mu_1 = \frac{1}{2}\delta_{-1} + \frac{1}{2}\delta_{1.5}$$

$$\rightarrow y_1 - x_1 = -1 > y_2 - x_2 = -2.5$$

Not a ray (particles cross at $t > 0$)




Illustration - Discrete Distributions


Let $x_1 < \dots < x_n$, $y_1 < \dots < y_n$, $\mu_0 = \frac{1}{n} \sum_{i=1}^n \delta_{x_i}$, $\mu_1 = \frac{1}{n} \sum_{i=1}^n \delta_{y_i}$.

$t \mapsto \mu_t$ is a geodesic ray if and only if for all $j > i$, $y_i - x_i \leq y_j - x_j$.

$$\mu_0 = \frac{1}{2}\delta_0 + \frac{1}{2}\delta_4, \quad \mu_1 = \frac{1}{2}\delta_{-1} + \frac{1}{2}\delta_{1.5}$$

$$\rightarrow y_1 - x_1 = -1 > y_2 - x_2 = -2.5$$

Not a ray (particles cross at $t > 0$)



Geodesic Rays in Brenier's Setting

Proposition (Brenier's Theorem (Brenier, 1991))

$\mu_0 \ll \text{Leb} \implies \text{Optimal coupling } \gamma^* \text{ unique and } \gamma^* = (\text{Id}, \nabla \varphi)_\# \mu_0 \text{ with } \varphi \text{ convex}$

In this setting:

- Geodesic between $\mu_0 \in \mathcal{P}_{2,\text{ac}}(\mathbb{R}^d)$ and $\mu_1 \in \mathcal{P}_2(\mathbb{R}^d)$ unique
- For all $t \in [0, 1]$, $\mu_t = ((1-t)\text{Id} + tT)_\# \mu_0$ with $T_\# \mu_0 = \mu_1$, $T = \nabla \varphi$, φ convex

Geodesic Rays in Brenier's Setting

Proposition (Brenier's Theorem (Brenier, 1991))

$\mu_0 \ll \text{Leb} \implies \text{Optimal coupling } \gamma^* \text{ unique and } \gamma^* = (\text{Id}, \nabla \varphi)_\# \mu_0 \text{ with } \varphi \text{ convex}$

In this setting:

- Geodesic between $\mu_0 \in \mathcal{P}_{2,\text{ac}}(\mathbb{R}^d)$ and $\mu_1 \in \mathcal{P}_2(\mathbb{R}^d)$ unique
- For all $t \in [0, 1]$, $\mu_t = ((1-t)\text{Id} + tT)_\# \mu_0$ with $T_\# \mu_0 = \mu_1$, $T = \nabla \varphi$, φ convex
- If $t \mapsto \mu_t$ is a geodesic ray:

$$\begin{aligned}\forall s \geq 0, \quad W_2^2(\mu_s, \mu_0) &= s^2 W_2^2(\mu_1, \mu_0) \\ &= \int \|s(x - \nabla \varphi(x))\|_2^2 \, d\mu_0(x) \\ &= \int \|x - (1-s)x - s\nabla \varphi(x)\|_2^2 \, d\mu_0(x)\end{aligned}$$

$\rightarrow (1-s)x + s\nabla \varphi(x) = \nabla(x \mapsto (1-s)\frac{\|x\|_2^2}{2} + s\varphi(x))(x)$ must be the gradient of a convex function for all $s \geq 0$

\rightarrow true if and only if $\varphi - \frac{\|x\|_2^2}{2}$ convex

Geodesic Rays in Brenier's Setting

Let $\mu_0 \in \mathcal{P}_{2,\text{ac}}(\mathbb{R}^d)$, $\mu_1 \in \mathcal{P}_2(\mathbb{R}^d)$, $\mu_1 = (\nabla \varphi)_\# \mu_0$, φ convex.
 $\rightarrow t \mapsto \mu_t$ geodesic ray if and only if φ 1-strongly convex

Examples

- 1D Gaussian: Let $\mu_0 = \mathcal{N}(m_0, \sigma_0^2)$, $\mu_1 = \mathcal{N}(m_1, \sigma_1^2)$, $T(x) = \frac{\sigma_1}{\sigma_0}(x - m_0) + m_1$
 $T'(x) - 1 \geq 0 \iff \sigma_1 \geq \sigma_0$
- General Gaussian: Let $\mu_0 = \mathcal{N}(m_0, \Sigma_0)$, $\mu_1 = \mathcal{N}(m_1, \Sigma_1)$,
 $T(x) = A(x - m_0) + m_1$ with $A = \Sigma_0^{-\frac{1}{2}} (\Sigma_0^{\frac{1}{2}} \Sigma_1 \Sigma_0^{\frac{1}{2}})^{\frac{1}{2}} \Sigma_0^{-\frac{1}{2}}$,
 $\nabla T(x) - I_d \succeq 0 \iff (\Sigma_0^{\frac{1}{2}} \Sigma_1 \Sigma_0^{\frac{1}{2}})^{\frac{1}{2}} \succeq \Sigma_0$

Busemann Function in the Wasserstein Space

Let $\mu_0 \in \mathcal{P}_2(\mathbb{R}^d)$, $t \mapsto \mu_t$ a geodesic ray starting from μ_0

Busemann function in $(\mathcal{P}_2(\mathbb{R}^d), W_2)$:

$$\forall \nu \in \mathcal{P}_2(\mathbb{R}^d), B^\mu(\nu) = \lim_{t \rightarrow +\infty} W_2(\mu_t, \nu) - \kappa_\mu t,$$

with $\kappa_\mu = W_2(\mu_0, \mu_1)$.

→ Computation?

Computing the Busemann Function

Define $\Gamma(\mu_0, \mu_1, \nu) = \{\tilde{\gamma} \in \Pi(\mu_0, \mu_1, \nu), \pi_{\#}^{1,2}\tilde{\gamma} \in \Pi_o(\mu_0, \mu_1)\}$

- General case:

$$B^\mu(\nu) = \inf_{\tilde{\gamma} \in \Gamma(\mu_0, \mu_1, \nu)} -\kappa_\mu^{-1} \int \langle x_1 - x_0, y - x_0 \rangle \, d\tilde{\gamma}(x_0, x_1, y)$$

Computing the Busemann Function

Define $\Gamma(\mu_0, \mu_1, \nu) = \{\tilde{\gamma} \in \Pi(\mu_0, \mu_1, \nu), \pi_{\#}^{1,2}\tilde{\gamma} \in \Pi_o(\mu_0, \mu_1)\}$

- General case:

$$B^\mu(\nu) = \inf_{\tilde{\gamma} \in \Gamma(\mu_0, \mu_1, \nu)} -\kappa_\mu^{-1} \int \langle x_1 - x_0, y - x_0 \rangle \, d\tilde{\gamma}(x_0, x_1, y)$$

- If $\mu_0 \in \mathcal{P}_{2,\text{ac}}(\mathbb{R}^d)$, $\mu_1 = (\nabla \varphi)_\# \mu_0$, φ 1-strongly convex:

$$B^\mu(\nu) = \inf_{\gamma \in \Pi(\mu_0, \nu)} -\kappa_\mu^{-1} \int \langle \nabla \varphi(x_0) - x_0, y - x_0 \rangle \, d\gamma(x_0, y)$$

→ equivalent to OT problem with cost $c(x_0, y) = \|\nabla \varphi(x_0) - x_0 - y\|_2^2$

Computing the Busemann Function

Define $\Gamma(\mu_0, \mu_1, \nu) = \{\tilde{\gamma} \in \Pi(\mu_0, \mu_1, \nu), \pi_{\#}^{1,2}\tilde{\gamma} \in \Pi_o(\mu_0, \mu_1)\}$

- General case:

$$B^{\mu}(\nu) = \inf_{\tilde{\gamma} \in \Gamma(\mu_0, \mu_1, \nu)} -\kappa_{\mu}^{-1} \int \langle x_1 - x_0, y - x_0 \rangle \, d\tilde{\gamma}(x_0, x_1, y)$$

- If $\mu_0 \in \mathcal{P}_{2,\text{ac}}(\mathbb{R}^d)$, $\mu_1 = (\nabla \varphi)_{\#} \mu_0$, φ 1-strongly convex:

$$B^{\mu}(\nu) = \inf_{\gamma \in \Pi(\mu_0, \nu)} -\kappa_{\mu}^{-1} \int \langle \nabla \varphi(x_0) - x_0, y - x_0 \rangle \, d\gamma(x_0, y)$$

- If $\mu_0 = \delta_{x_0}$, $x_0 \in \mathbb{R}^d$, $\pi_{\#}^{1,2}\tilde{\gamma} = \mu_0 \otimes \mu_1$,

$$B^{\mu}(\nu) = \inf_{\gamma \in \Pi(\mu_1, \nu)} -\kappa_{\mu}^{-1} \int \langle x_1 - x_0, y - x_0 \rangle \, d\gamma(x_1, y)$$

→ Equivalent to $W_2^2(\mu_1, \nu)$

→ For $\mu_1 = \delta_{x_1}$, $\theta = x_1 - x_0 \in S^{d-1}$, $\gamma(t) = t\theta$, $B^{\mu}(\nu) = \int B^{\gamma}(y) d\nu(y)$

Computing the Busemann Function

Define $\Gamma(\mu_0, \mu_1, \nu) = \{\tilde{\gamma} \in \Pi(\mu_0, \mu_1, \nu), \pi_{\#}^{1,2}\tilde{\gamma} \in \Pi_o(\mu_0, \mu_1)\}$

- General case:

$$B^\mu(\nu) = \inf_{\tilde{\gamma} \in \Gamma(\mu_0, \mu_1, \nu)} -\kappa_\mu^{-1} \int \langle x_1 - x_0, y - x_0 \rangle \, d\tilde{\gamma}(x_0, x_1, y)$$

- If $\mu_0 \in \mathcal{P}_{2,\text{ac}}(\mathbb{R}^d)$, $\mu_1 = (\nabla \varphi)_\# \mu_0$, φ 1-strongly convex:

$$B^\mu(\nu) = \inf_{\gamma \in \Pi(\mu_0, \nu)} -\kappa_\mu^{-1} \int \langle \nabla \varphi(x_0) - x_0, y - x_0 \rangle \, d\gamma(x_0, y)$$

- If $\mu_0 = \delta_{x_0}$, $x_0 \in \mathbb{R}^d$, $\pi_{\#}^{1,2}\tilde{\gamma} = \mu_0 \otimes \mu_1$,

$$B^\mu(\nu) = \inf_{\gamma \in \Pi(\mu_1, \nu)} -\kappa_\mu^{-1} \int \langle x_1 - x_0, y - x_0 \rangle \, d\gamma(x_1, y)$$

→ Linear programs

Busemann Function in the 1D Wasserstein Space

Let $\mu_0, \mu_1 \in \mathcal{P}_2(\mathbb{R})$,

$$W_2^2(\mu_0, \mu_1) = \int_0^1 |F_0^{-1}(u) - F_1^{-1}(u)|^2 du = \|F_0^{-1} - F_1^{-1}\|_{L^2([0,1])}^2$$

→ Hilbert structure

Proposition (Closed-form for the Busemann function on $\mathcal{P}_2(\mathbb{R})$)

Let $(\mu_t)_{t \geq 0}$ be a unit-speed geodesic ray in $\mathcal{P}_2(\mathbb{R})$, then

$$\begin{aligned} \forall \nu \in \mathcal{P}_2(\mathbb{R}), \quad B^\mu(\nu) &= - \int_0^1 (F_1^{-1}(u) - F_0^{-1}(u))(F_\nu^{-1}(u) - F_0^{-1}(u)) du \\ &= -\langle F_1^{-1} - F_0^{-1}, F_\nu^{-1} - F_0^{-1} \rangle_{L^2([0,1])}. \end{aligned}$$

Example

Let $\mu_0 = \mathcal{N}(m_0, \sigma_0^2)$, $\mu_1 = \mathcal{N}(m_1, \sigma_1^2)$, $\nu = \mathcal{N}(m, \sigma^2)$,

$$B^\mu(\nu) = -(m_1 - m_0)(m - m_0) - (\sigma_1 - \sigma_0)(\sigma - \sigma_0)$$

Busemann Function on the Bures-Wasserstein Space

Let $\mu_0 = \mathcal{N}(m_0, \Sigma_0)$, $\mu_1 = \mathcal{N}(m_1, \Sigma_1)$,

$$W_2^2(\mu_0, \mu_1) = \|m_0 - m_1\|_2^2 + \text{tr} \left(\Sigma_0 + \Sigma_1 - 2(\Sigma_0^{\frac{1}{2}} \Sigma_1 \Sigma_0^{\frac{1}{2}})^{\frac{1}{2}} \right)$$

Proposition (Closed-form for the Busemann function on $BW(\mathbb{R}^d)$)

Let $(\mu_t)_{t \geq 0}$ be a unit-speed geodesic ray characterized by $\mu_0 = \mathcal{N}(m_0, \Sigma_0)$ and $\mu_1 = \mathcal{N}(m_1, \Sigma_1)$. Then, for any $\nu = \mathcal{N}(m, \Sigma)$,

$$\begin{aligned} B^\mu(\nu) &= -\langle m_1 - m_0, m - m_0 \rangle + \text{tr} (\Sigma_0(A - I_d)) \\ &\quad - \text{tr} \left((\Sigma^{\frac{1}{2}} (\Sigma_0 - \Sigma_0 A - A \Sigma_0 + \Sigma_1) \Sigma^{\frac{1}{2}})^{\frac{1}{2}} \right), \end{aligned}$$

where $A = \Sigma_0^{-\frac{1}{2}} (\Sigma_0^{\frac{1}{2}} \Sigma_1 \Sigma_0^{\frac{1}{2}})^{\frac{1}{2}} \Sigma_0^{-\frac{1}{2}}$.

Table of Contents

Busemann Function in Metric Spaces

Busemann Functions in Wasserstein Space

Comparison of Labeled Datasets

Labeled Datasets

$$\mathcal{D}_1 : \mu_1 = \frac{1}{m} \sum_{i=1}^m \delta_{(x_i^1, y_i^1)} \in \mathcal{P}(\mathbb{R}^d \times \{1, \dots, C\}),$$

$$\mathcal{D}_2 : \mu_2 = \frac{1}{m} \sum_{j=1}^m \delta_{(x_j^2, y_j^2)} \in \mathcal{P}(\mathbb{R}^d \times \{1, \dots, C\})$$

C : number of classes, n : number of sample in each class, $m = nC$


Question: how to compare datasets \mathcal{D}_1 and \mathcal{D}_2 ?



OTDD (Alvarez-Melis and Fusi, 2020)

Solution of Alvarez-Melis and Fusi (2020):

- Embed a label (a class) in $\mathcal{P}(\mathbb{R}^d)$ as $c \mapsto \nu_c^k = \frac{1}{n} \sum_{i=1}^n \delta_{x_i^k} \mathbb{1}_{\{y_i^k=c\}}$ for $k = 1, 2$




$$\rightarrow \mathcal{D}_k : \mu_k = \frac{1}{m} \sum_{i=1}^m \delta_{(x_i^k, \nu_{y_i^k}^k)} \in \mathcal{P}(\mathbb{R}^d \times \mathcal{P}(\mathbb{R}^d))$$

OTDD (Alvarez-Melis and Fusi, 2020)

Solution of Alvarez-Melis and Fusi (2020):

- Embed a label (a class) in $\mathcal{P}(\mathbb{R}^d)$ as $c \mapsto \nu_c^k = \frac{1}{n} \sum_{i=1}^n \delta_{x_i^k} \mathbb{1}_{\{y_i^k=c\}}$ for $k = 1, 2$



$$\rightarrow \mathcal{D}_k : \mu_k = \frac{1}{m} \sum_{i=1}^m \delta_{(x_i^k, \nu_{y_i^k}^k)} \in \mathcal{P}(\mathbb{R}^d \times \mathcal{P}(\mathbb{R}^d))$$


- Cost: $d((x, y), (x', y'))^2 = \|x - x'\|_2^2 + W_2^2(\nu_y, \nu_{y'})$
- **Optimal transport distance:** $O(C^2 n^3 \log n + n^3 C^3 \log(nC))$

$$\text{OTDD}(\mu_1, \mu_2) = \inf_{\gamma \in \Pi(\mu_1, \mu_2)} \int d((x, y), (x', y'))^2 \, d\gamma((x, y), (x', y')).$$

OTDD (Alvarez-Melis and Fusi, 2020)

Solution of Alvarez-Melis and Fusi (2020):

- Embed a label (a class) in $\mathbb{R}^p \times S_p^{++}(\mathbb{R})$ as $c \mapsto \nu_c^k \approx \mathcal{N}(m_c^k, \Sigma_c^k)$ for $k = 1, 2$





$$\rightarrow \mathcal{D}_k : \mu_k = \frac{1}{m} \sum_{i=1}^m \delta_{(x_i^k, m_{y_i^k}^k, \Sigma_{y_i^k}^k)} \in \mathcal{P}(\mathbb{R}^d \times \mathbb{R}^p \times S_p^{++}(\mathbb{R}))$$

- Cost: $d((x, y), (x', y'))^2 = \|x - x'\|_2^2 + \text{BW}_2^2(\nu_y, \nu_{y'})$
- **Optimal transport distance:** approximated in $O(C^2 d^3 + n^2 C^2 \log(nC)/\varepsilon^2)$

$$\text{OTDD}_\varepsilon(\mu_1, \mu_2) = \inf_{\gamma \in \Pi(\mu_1, \mu_2)} \int d((x, y), (x', y'))^2 \, d\gamma((x, y), (x', y')) + \varepsilon \mathcal{H}(\gamma).$$

Sliced-Wasserstein Distance



Definition (Sliced-Wasserstein (Rabin et al., 2011))

Let $\mu, \nu \in \mathcal{P}_2(\mathbb{R}^d)$,

$$\text{SW}_2^2(\mu, \nu) = \int_{S^{d-1}} \text{W}_2^2(P_\#^\theta \mu, P_\#^\theta \nu) \, d\lambda(\theta),$$

where $P^\theta(x) = \langle x, \theta \rangle$, λ uniform measure on S^{d-1} .

Sliced-Wasserstein Distance

Let $x_1, \dots, x_n, y_1, \dots, y_n \in \mathbb{R}^d$, $\alpha, \beta \in \Sigma_n$, $\mu = \sum_{i=1}^n \alpha_i \delta_{x_i}$, $\nu = \sum_{i=1}^n \beta_i \delta_{y_i}$.

Approximation via Monte-Carlo:

$$\widehat{\text{SW}}_{2,L}^2(\mu, \nu) = \frac{1}{L} \sum_{\ell=1}^L \text{W}_2^2(P_{\#}^{\theta_\ell} \mu, P_{\#}^{\theta_\ell} \nu),$$

$\theta_1, \dots, \theta_L \sim \lambda$.

→ Computational complexity: $O(L n (\log n + d))$

Goal: Define a SW distance on $\mathcal{P}_2(\mathbb{R}^d \times \mathcal{P}_2(\mathbb{R}^d))$

Previous Methods

- Sliced-Wasserstein on $\mathcal{P}_2(X \times Y)$ ([Nguyen and Ho, 2024](#)):

- Define 2 projections $P^\theta : X \rightarrow \mathbb{R}$, $Q^\phi : Y \rightarrow \mathbb{R}$
 - For $\alpha \in S^1$, define

$$\forall (x, y) \in X \times Y, P^{\alpha, \theta, \phi}(x, y) = \alpha_1 P^\theta(x) + \alpha_2 Q^\phi(y)$$

- For $\mu, \nu \in \mathcal{P}_2(X \times Y)$,

$$\text{SW}_2^2(\mu, \nu) = \int W_2^2(P_{\#}^{\alpha, \theta, \phi} \mu, P_{\#}^{\alpha, \theta, \phi} \nu) d\lambda(\alpha, \theta, \phi)$$

Previous Methods

- Sliced-Wasserstein on $\mathcal{P}_2(X \times Y)$ ([Nguyen and Ho, 2024](#)):

- Define 2 projections $P^\theta : X \rightarrow \mathbb{R}$, $Q^\phi : Y \rightarrow \mathbb{R}$
 - For $\alpha \in S^1$, define

$$\forall (x, y) \in X \times Y, P^{\alpha, \theta, \phi}(x, y) = \alpha_1 P^\theta(x) + \alpha_2 Q^\phi(y)$$

- For $\mu, \nu \in \mathcal{P}_2(X \times Y)$,

$$\text{SW}_2^2(\mu, \nu) = \int W_2^2(P_{\#}^{\alpha, \theta, \phi} \mu, P_{\#}^{\alpha, \theta, \phi} \nu) d\lambda(\alpha, \theta, \phi)$$

- Sliced-Wasserstein on $\mathcal{P}_2(\mathbb{R}^d \times \mathcal{P}_2(\mathbb{R}^d))$ ([SOTDD](#)) ([Nguyen et al., 2025](#))

- For a label $y \in \{1, \dots, C\}$, define $\varphi(y) = \frac{1}{n_y} \sum_{i=1}^n \delta_{x_i} \mathbb{1}_{\{y_i=y\}}$
 - Use for $\alpha \in S^k$,

$$P^{\alpha, \theta, \lambda}(x, y) = \alpha_1 P^\theta(x) + \sum_{i=1}^k \alpha_{i+1} \mathcal{M}^{\lambda_i} (P_{\#}^\theta \varphi(y)),$$

with $P^\theta : \mathbb{R}^d \rightarrow \mathbb{R}$, $\mathcal{M}^\lambda : \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R}$ the moment transform projection.

Previous Methods

- Sliced-Wasserstein on $\mathcal{P}_2(X \times Y)$ ([Nguyen and Ho, 2024](#)):

- Define 2 projections $P^\theta : X \rightarrow \mathbb{R}$, $Q^\phi : Y \rightarrow \mathbb{R}$
 - For $\alpha \in S^1$, define

$$\forall (x, y) \in X \times Y, P^{\alpha, \theta, \phi}(x, y) = \alpha_1 P^\theta(x) + \alpha_2 Q^\phi(y)$$

- For $\mu, \nu \in \mathcal{P}_2(X \times Y)$,

$$\text{SW}_2^2(\mu, \nu) = \int W_2^2(P_{\#}^{\alpha, \theta, \phi} \mu, P_{\#}^{\alpha, \theta, \phi} \nu) d\lambda(\alpha, \theta, \phi)$$

- Sliced-Wasserstein on $\mathcal{P}_2(\mathbb{R}^d \times \mathcal{P}_2(\mathbb{R}^d))$ ([SOTDD](#)) ([Nguyen et al., 2025](#))

- For a label $y \in \{1, \dots, C\}$, define $\varphi(y) = \frac{1}{n_y} \sum_{i=1}^n \delta_{x_i} \mathbb{1}_{\{y_i=y\}}$
 - Use for $\alpha \in S^k$,

$$P^{\alpha, \theta, \lambda}(x, y) = \alpha_1 P^\theta(x) + \sum_{i=1}^k \alpha_{i+1} \mathcal{M}^{\lambda_i} (P_{\#}^\theta \varphi(y)),$$

with $P^\theta : \mathbb{R}^d \rightarrow \mathbb{R}$, $\mathcal{M}^\lambda : \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R}$ the moment transform projection.

→ Use B^μ for projecting distributions on \mathbb{R}


Slicing Datasets with Busemann on Gaussian

With Gaussian approximation:

- Define $\Xi(\mu) = \mathcal{N}(m(\mu), \Sigma(\mu))$
- For all $y \in \{1, \dots, C\}$,

$$Q^\eta(y) = B^\eta(\Xi(\varphi(y))),$$

with η a geodesic ray on $BW(\mathbb{R}^d)$



Computational Complexity: $O(LCd^3 + Ln_y C(\log n_y C + d) + d^2 Cn_y)$


Slicing Datasets with Busemann in 1D

With 1D Projection:

- For all $y \in \{1, \dots, C\}$,

$$Q^{\eta, \theta}(y) = B^\eta(P_\#^\theta \varphi(y)),$$

with η a geodesic ray on $\mathcal{P}_2(\mathbb{R})$





Computation Complexity: $O(L n_y C (\log(n_y C + d)))$

Correlation vs OTDD


Goal: Measure correlation between sliced distances and OTDD

→ Compare randomly sampled subdatasets + Spearman and Pearson correlations



Flowing Labeled Datasets

- Model datasets as $\mathbb{P} = \frac{1}{C} \sum_{c=1}^C \delta_{\nu^c} \in \mathcal{P}_2(\mathcal{P}_2(\mathbb{R}^d))$ where $\nu^c = \frac{1}{n} \sum_{i=1}^n \delta_{x_i^c}$
- Flow a dataset \mathbb{P} towards \mathbb{Q} by minimizing a discrepancy D on $\mathcal{P}_2(\mathcal{P}_2(\mathbb{R}^d))$
→ minimization problem on $\mathcal{P}_2(\mathcal{P}_2(\mathbb{R}^d))$



Example

Let $\psi\left(\frac{1}{C} \sum_{c=1}^C \delta_{\mu^{n,c}}\right) = \left(\frac{1}{nC} \sum_{c=1}^C \sum_{i=1}^n \delta_{x_i^c}, \frac{1}{n} \sum_{i=1}^n \delta_{x_i^c}\right)$.
 $\rightarrow D(\mathbb{P}, \mathbb{Q}) = \text{SWB1DG}(\mathbb{P}, \mathbb{Q})$

Minimizing on $\mathcal{P}_2(\mathcal{P}_2(\mathbb{R}^d))$ (Bonet et al., 2025b)


Goal: minimize $\mathbb{F} : \mathcal{P}_2(\mathcal{P}_2(\mathbb{R}^d)) \rightarrow \mathbb{R}$

In practice: For $\mathbb{P}_k = \frac{1}{C} \sum_{c=1}^C \delta_{\mu_k^{c,n}}$ with $\mu_k^{c,n} = \frac{1}{n} \sum_{i=1}^n \delta_{x_{i,k}^c} \in \mathcal{P}_2(\mathbb{R}^d)$:

$\forall k \geq 0$, particle (image) i , class c , $x_{i,k+1}^c = x_{i,k}^c - \tau \nabla_{W_{W_2}} \mathbb{F}(\mathbb{P}_k)(\mu_k^{c,n})(x_{i,k}^c)$.

\mathbb{P}_k : inter-class interaction, $\mu_k^{c,n}$: intra-class interaction, $x_{i,k}^c$ image

$\nabla_{W_{W_2}} \mathbb{F}(\mathbb{P}_k)(\mu_k^{c,n})(x_{i,k}^c) = nC[\nabla F(\mathbf{x})]_{i,c}$ with $F(\mathbf{x}) = \mathbb{F}(\mathbb{P}_k)$, $\mathbf{x} = (x_{i,k}^c)_{i,c}$



Synthetic Data

Goal: minimize $\mathbb{F} : \mathcal{P}_2(\mathcal{P}_2(\mathbb{R}^d)) \rightarrow \mathbb{R}$

In practice: For $\mathbb{P}_k = \frac{1}{C} \sum_{c=1}^C \delta_{\mu_k^{c,n}}$ with $\mu_k^{c,n} = \frac{1}{n} \sum_{i=1}^n \delta_{x_{i,k}^c} \in \mathcal{P}_2(\mathbb{R}^d)$:

$\forall k \geq 0$, particle (image) i , class c , $x_{i,k+1}^c = x_{i,k}^c - \tau \nabla_{W_{W_2}} \mathbb{F}(\mathbb{P}_k)(\mu_k^{c,n})(x_{i,k}^c)$.

Let $\mathbb{Q} = \frac{1}{3} \sum_{c=1}^3 \delta_{\nu^c}$, ν^c ring

$$\min_{\mathbb{P}} \mathbb{F}(\mathbb{P}) = D(\mathbb{P}, \mathbb{Q})$$

SOTDD



SWB1DG



SWBG



Synthetic Data

Goal: minimize $\mathbb{F} : \mathcal{P}_2(\mathcal{P}_2(\mathbb{R}^d)) \rightarrow \mathbb{R}$

In practice: For $\mathbb{P}_k = \frac{1}{C} \sum_{c=1}^C \delta_{\mu_k^{c,n}}$ with $\mu_k^{c,n} = \frac{1}{n} \sum_{i=1}^n \delta_{x_{i,k}^c} \in \mathcal{P}_2(\mathbb{R}^d)$:

$\forall k \geq 0$, particle (image) i , class c , $x_{i,k+1}^c = x_{i,k}^c - \tau \nabla_{W_{W_2}} \mathbb{F}(\mathbb{P}_k)(\mu_k^{c,n})(x_{i,k}^c)$.

Let $\mathbb{Q} = \frac{1}{3} \sum_{c=1}^3 \delta_{\nu^c}$, ν^c ring

$$\min_{\mathbb{P}} \mathbb{F}(\mathbb{P}) = D(\mathbb{P}, \mathbb{Q})$$

SOTDD



SWB1DG



SWBG



Synthetic Data


Goal: minimize $\mathbb{F} : \mathcal{P}_2(\mathcal{P}_2(\mathbb{R}^d)) \rightarrow \mathbb{R}$

In practice: For $\mathbb{P}_k = \frac{1}{C} \sum_{c=1}^C \delta_{\mu_k^{c,n}}$ with $\mu_k^{c,n} = \frac{1}{n} \sum_{i=1}^n \delta_{x_{i,k}^c} \in \mathcal{P}_2(\mathbb{R}^d)$:

$\forall k \geq 0$, particle (image) i , class c , $x_{i,k+1}^c = x_{i,k}^c - \tau \nabla_{W_{W_2}} \mathbb{F}(\mathbb{P}_k)(\mu_k^{c,n})(x_{i,k}^c)$.

Let $\mathbb{Q} = \frac{1}{3} \sum_{c=1}^3 \delta_{\nu^c}$, ν^c ring

$$\min_{\mathbb{P}} \mathbb{F}(\mathbb{P}) = D(\mathbb{P}, \mathbb{Q})$$



Conclusion

Conclusion:

- Studied geometry of geodesics on the Wasserstein space
- Studied the existence and computation of the Busemann function on the Wasserstein space
- Defined new SW distances to compare labeled Datasets
- Also in the paper: SW distances to compare Gaussian mixtures
- Related to new sliced distances on Gaussian mixtures ([Baouan et al., 2025](#); [Piening and Beinert, 2025a](#)) and on $\mathcal{P}_2(\mathcal{P}_2(\mathbb{R}^d))$ ([Pieming and Beinert, 2025b](#))

Perspectives:

- Applications to PCA on the Wasserstein space
- Extension to $\mathcal{P}_2(\mathbb{L}^d)$
- Busemann along other curves ([Gallouët et al., 2025](#))

Conclusion

Thank you!

Paper: <https://arxiv.org/abs/2510.04579>



References |

- David Alvarez-Melis and Nicolo Fusi. Geometric Dataset Distances via Optimal Transport. *Advances in Neural Information Processing Systems*, 33: 21428–21439, 2020.
- Luigi Ambrosio, Nicola Gigli, and Giuseppe Savaré. *Gradient flows: in metric spaces and in the space of probability measures*. Springer, 2008.
- Ali Baouan, Mathieu Rosenbaum, and Sergio Pulido. An optimal transport based embedding to quantify the distance between playing styles in collective sports. *Journal of Quantitative Analysis in Sports*, 2025.
- Riccardo Bellazzi, Andrea Codegoni, Stefano Gualandi, Giovanna Nicora, and Eleonora Vercesi. The gene mover's distance: Single-cell similarity via optimal transport. *arXiv preprint arXiv:2102.01218*, 2021.
- Paul Berg, Bjoern Michele, Minh-Tan Pham, Laetitia Chapel, and Nicolas Courty. Horospherical Learning with Smart Prototypes. In *British Machine Vision Conference (BMVC)*, 2024.
- Paul Berg, Léo Buecher, Björn Michele, Minh-Tan Pham, Laetitia Chapel, and Nicolas Courty. Multi-Prototype Hyperbolic Learning Guided by Class Hierarchy. *International Journal of Computer Vision*, pages 1–16, 2025.

References II

- Jérôme Bertrand and Benoît Kloeckner. A geometric study of Wasserstein spaces: isometric rigidity in negative curvature. *International Mathematics Research Notices*, 2016(5):1368–1386, 2016.
- Clément Bonet, Laetitia Chapel, Lucas Drumetz, and Nicolas Courty. Hyperbolic Sliced-Wasserstein via Geodesic and Horospherical Projections. In *Proceedings of 2nd Annual Workshop on Topology, Algebra, and Geometry in Machine Learning (TAG-ML)*, pages 334–370. PMLR, 2023.
- Clément Bonet, Lucas Drumetz, and Nicolas Courty. Sliced-Wasserstein Distances and Flows on Cartan-Hadamard Manifolds. *Journal of Machine Learning Research*, 26(32):1–76, 2025a.
- Clément Bonet, Christophe Vauthier, and Anna Korba. Flowing Datasets with Wasserstein over Wasserstein Gradient Flows. In *International Conference on Machine Learning*. PMLR, 2025b.
- Yann Brenier. Polar factorization and monotone rearrangement of vector-valued functions. *Communications on pure and applied mathematics*, 44(4):375–417, 1991.
- Martin R Bridson and André Haefliger. *Metric Spaces of Non-Positive Curvature*, volume 319. Springer Science & Business Media, 2013.

References III

- Herbert Busemann. *The Geometry of Geodesics*. Academic Press, New York, 1955.
- Ines Chami, Albert Gu, Dat P Nguyen, and Christopher Ré. HoroPCA: Hyperbolic Dimensionality Reduction via Horospherical Projections. In *International Conference on Machine Learning*, pages 1419–1429. PMLR, 2021.
- Julie Delon and Agnes Desolneux. A Wasserstein-Type Distance in the space of Gaussian Mixture Models. *SIAM Journal on Imaging Sciences*, 13(2):936–970, 2020.
- Lars Doorenbos, Pablo Márquez Neila, Raphael Sznitman, and Pascal Mettes. Hyperbolic Random Forests. *Transactions on Machine Learning Research*, 2024. ISSN 2835-8856.
- Xiran Fan, Chun-Hao Yang, and Baba Vemuri. Horocycle Decision Boundaries for Large Margin Classification in Hyperbolic Space. *Advances in neural information processing systems*, 36:11194–11204, 2023.
- Thomas O Gallouët, Andrea Natale, and Gabriele Todeschi. Metric extrapolation in the wasserstein space. *Calculus of Variations and Partial Differential Equations*, 64(5):147, 2025.

References IV

- Jonathan Geuter, Clément Bonet, Anna Korba, and David Alvarez-Melis. DDEQs: Distributional Deep Equilibrium Models through Wasserstein Gradient Flows. In *The 28th International Conference on Artificial Intelligence and Statistics*, 2025.
- Mina Ghadimi Atigh, Martin Keller-Ressel, and Pascal Mettes. Hyperbolic Busemann Learning with Ideal Prototypes. *Advances in Neural Information Processing Systems*, 34:103–115, 2021.
- Benoit Kloeckner. A geometric study of Wasserstein spaces: Euclidean spaces. *Annali della Scuola Normale Superiore di Pisa-Classe di Scienze*, 9(2):297–323, 2010.
- Matt Kusner, Yu Sun, Nicholas Kolkin, and Kilian Weinberger. From word embeddings to document distances. In *International conference on machine learning*, pages 957–966. PMLR, 2015.
- Khai Nguyen and Nhat Ho. Hierarchical Hybrid Sliced Wasserstein: A Scalable Metric for Heterogeneous Joint Distributions. *Advances in Neural Information Processing Systems*, 37:108140–108166, 2024.
- Khai Nguyen, Hai Nguyen, Tuan Pham, and Nhat Ho. Lightspeed Geometric Dataset Distance via Sliced Optimal Transport. In *Forty-second International Conference on Machine Learning*, 2025.

References V

- Moritz Piening and Robert Beinert. Slicing the Gaussian Mixture Wasserstein Distance. *Transactions on Machine Learning Research*, 2025a. ISSN 2835-8856.
- Moritz Piening and Robert Beinert. Slicing Wasserstein Over Wasserstein Via Functional Optimal Transport. *arXiv preprint arXiv:2509.22138*, 2025b.
- Julien Rabin, Gabriel Peyré, Julie Delon, and Marc Bernot. Wasserstein barycenter and its application to texture mixing. In *International Conference on Scale Space and Variational Methods in Computer Vision*, pages 435–446. Springer, 2011.
- Yossi Rubner, Carlo Tomasi, and Leonidas J Guibas. The Earth Mover’s Distance as a Metric for Image Retrieval. *International journal of computer vision*, 40(2):99–121, 2000.
- Luke Vilnis and Andrew McCallum. Word representations via gaussian embedding. In Yoshua Bengio and Yann LeCun, editors, *3rd International Conference on Learning Representations, ICLR 2015, San Diego, CA, USA, May 7-9, 2015, Conference Track Proceedings*, 2015.
- Guomin Zhu, Wen-Long Li, and Xiaojun Cui. Busemann functions on the wasserstein space. *Calculus of Variations and Partial Differential Equations*, 60(3):97, 2021.