

Question

Given the following equations, find the values of x , y , and z

$$\begin{cases} 2 \cos x \sin y + 4 \sin y - 3 \cos x = -4 \\ 8 \sin y \sin z \cos z \cos 2z - 6 \sin 2z \cos 2z + 4 \sin y = -9 \\ 4 \cos x \cos z \cos 2z \sin z + 2 \cos x + 4 \sin 2z \cos 2z = 2 \end{cases}$$

Solution

Simplifying the Equations

Recall the double angle formula:

$$\sin 2x = 2 \sin x \cos x$$

The first equation has no terms in which we may apply the double angle formula to, therefore we leave it as is. Looking at the second equation:

$$8 \sin y \sin z \cos z \cos 2z - 6 \sin 2z \cos 2z + 4 \sin y = -9$$

We can simplify it:

$$\begin{aligned} 8 \sin y \sin z \cos z \cos 2z - 6 \sin 2z \cos 2z + 4 \sin y &= -9 \\ 4 \sin y \cos 2z (2 \sin z \cos z) - 3 (2 \sin 2z \cos 2z) + 4 \sin y &= -9 \\ 4 \sin y \cos 2z (\sin 2z) - 3 (\sin 2 (2z)) + 4 \sin y &= -9 \\ 4 \sin y \cos 2z \sin 2z - 3 \sin 4z + 4 \sin y &= -9 \\ 2 \sin y (2 \cos 2z \sin 2z) - 3 \sin 4z + 4 \sin y &= -9 \\ 2 \sin y (\sin 2 (2z)) - 3 \sin 4z + 4 \sin y &= -9 \\ 2 \sin y \sin 4z - 3 \sin 4z + 4 \sin y &= -9 \end{aligned} \tag{1}$$

The third equation can also be simplified:

$$\begin{aligned} 4 \cos x \cos z \cos 2z \sin z + 2 \cos x + 4 \sin 2z \cos 2z &= 2 \\ 2 \cos x \cos 2z (2 \sin z \cos z) + 2 \cos x + 2 (2 \sin 2z \cos 2z) &= 2 \\ 2 \cos x \cos 2z (\sin 2z) + 2 \cos x + 2 (\sin 2 (2z)) &= 2 \\ 2 \cos x \cos 2z \sin 2z + 2 \cos x + 2 \sin 4z &= 2 \\ \cos x (2 \cos 2z \sin 2z) + 2 \cos x + 2 \sin 4z &= 2 \\ \cos x (\sin 2 (2z)) + 2 \cos x + 2 \sin 4z &= 2 \\ \cos x \sin 4z + 2 \cos x + 2 \sin 4z &= 2 \end{aligned} \tag{2}$$

Factor by Grouping

Upon the completion of simplifying the given equations, it is now possible to further simplify with the process of factoring.

Here are the newly simplified equations obtained from the last section:

$$\begin{cases} 2 \cos x \sin y + 4 \sin y - 3 \cos x = -4 \\ 2 \sin y \sin 4z - 3 \sin 4z + 4 \sin y = -9 \\ \cos x \sin 4z + 2 \cos x + 2 \sin 4z = 2 \end{cases}$$

We can follow the steps for how to factor by grouping for the first equation:

$$\begin{aligned}
& 2 \cos x \sin y + 4 \sin y - 3 \cos x = -4 \\
& \text{the first two terms have a common factor of } 2 \sin y \\
& 2 \sin y(\cos x + 2) - 3 \cos x = -4 \\
& \text{we see the part inside the brackets is } \cos x + 2 \\
& 2 \sin y(\cos x + 2) - 3 \cos x - 6 + 6 = -4 \\
& \text{to obtain } \cos x + 2 \text{ from } -3 \cos x \text{ we need to factor out a } -3 \\
& \text{because } -3 \cos x \text{ has a coefficient of } -3 \text{ and} \\
& \text{we want } \cos x + 2 \text{ therefore we must subtract 6.} \tag{3} \\
& \text{if we subtracted 6, we also need to add 6 to keep the equation balanced} \\
& \text{now } -3 \text{ can be factored out of the last two terms on the left side} \\
& 2 \sin y(\cos x + 2) - 3(\cos x + 2) + 6 = -4 \\
& \text{we can move the } +6 \text{ to the other side of the equation} \\
& 2 \sin y(\cos x + 2) - 3(\cos x + 2) = -10 \\
& \text{factor the left side by grouping} \\
& (2 \sin y - 3)(\cos x + 2) = -10
\end{aligned}$$

The same concept can be applied for the second and third equation. Some algebraic rearrangements may be required. The factorization is left as an exercise for the reader. Solutions on next line:

$$\begin{aligned}
& 2 \sin y \sin 4z - 3 \sin 4z + 4 \sin y = -9 \\
& 2 \sin y \sin 4z + 4 \sin y - 3 \sin 4z = -9 \\
& 2 \sin y(\sin 4z + 2) - 3 \sin 4z = -9 \\
& 2 \sin y(\sin 4z + 2) - 3 \sin 4z - 6 + 6 = -9 \\
& 2 \sin y(\sin 4z + 2) - 3(\sin 4z + 2) + 6 = -9 \\
& 2 \sin y(\sin 4z + 2) - 3(\sin 4z + 2) = -15 \\
& (2 \sin y - 3)(\sin 4z + 2) = -15
\end{aligned} \tag{4}$$

$$\begin{aligned}
& \cos x \sin 4z + 2 \cos x + 2 \sin 4z = 2 \\
& \cos x(\sin 4z + 2) + 2 \sin 4z = 2 \\
& \cos x(\sin 4z + 2) + 2 \sin 4z + 4 - 4 = 2 \\
& \cos x(\sin 4z + 2) + 2(\sin 4z + 2) - 4 = 2 \\
& \cos x(\sin 4z + 2) + 2(\sin 4z + 2) = 6 \\
& (\cos x + 2)(\sin 4z + 2) = 6
\end{aligned} \tag{5}$$

Solving Systems of Equations

After finishing all factoring exercises which have been assigned to the reader, the reader should have a new system of equations:

$$\begin{cases} (2 \sin y - 3)(\cos x + 2) = -10 \\ (2 \sin y - 3)(\sin 4x + 2) = -15 \\ (\cos x + 2)(\sin 4x + 2) = 6 \end{cases}$$

Upon realizing the three equations share the same factors, we can substitute these factors as letters to visualize it better.

Substitute:

$$\begin{aligned}
& \text{Let } A = (\cos x + 2) \\
& \text{Let } B = (2 \sin y - 3) \\
& \text{Let } C = (\sin 4x + 2)
\end{aligned} \tag{6}$$

The original system of equations can now be written as

$$\begin{cases} BA = -10 \\ BC = -15 \\ AC = 6 \end{cases}$$

If all three equations in the system is multiplied together:

$$\begin{aligned} (AB) \cdot (BC) \cdot (AC) &= (-10)(-15)(6) \\ A^2 B^2 C^2 &= 900 \\ ABC &= \pm 30 \\ ABC &= 30 \text{ and } ABC = -30 \end{aligned} \tag{7}$$

If we divide ABC by the equations in our system of equations we should be able to obtain the values for the last variable.

$$\begin{aligned} \frac{ABC}{BA} &= \frac{\pm 30}{-10} \\ C &= \pm 3 \end{aligned}$$

$$\begin{aligned} \frac{ABC}{BC} &= \frac{\pm 30}{-15} \\ A &= \pm 2 \end{aligned} \tag{8}$$

$$\begin{aligned} \frac{ABC}{AC} &= \frac{\pm 30}{6} \\ B &= \pm 5 \end{aligned}$$

We can now substitute the factors back and write a new system of equations:

$$\begin{cases} (\cos x + 2) = \pm 2 \\ (2 \sin y - 3) = \pm 5 \\ (\sin 4x + 2) = \pm 3 \end{cases}$$

Solving Trig

For

$$\begin{cases} (\cos x + 2) = \pm 2 \\ (2 \sin y - 3) = \pm 5 \\ (\sin 4x + 2) = \pm 3 \end{cases}$$

not all values of the equation will provide real solutions therefore some values should be rejected.

Given $(\cos x + 2) = \pm 2$ we can come up with these two equations

$$\begin{cases} \cos x + 2 = 2 \\ \cos x + 2 = -2 \end{cases}$$

From $\cos x + 2 = 2$ we can get $\cos x = 0$

From $\cos x + 2 = -2$ we can get $\cos x = -4$

The range of $\cos x$ is $y \in \mathbb{R}, y \in [-1, 1]$

We can reject $\cos x = -4$ because it's outside the

range of $\cos x$ and won't be providing any real solutions

$\cos x = 0$ is $x = \pi n$ and repeats every $\frac{\pi}{2}$ units

therefore a general solution for $\cos x = 0$ is $x = \pi n + \frac{\pi}{2}, n \in \mathbb{Z}$

The same concept can be repeated for the second equation:

$$\begin{aligned}
 2 \sin y - 3 &= \pm 5 \\
 2 \sin y - 3 &= 5 \text{ and } 2 \sin y - 3 = -5 \\
 2 \sin y &= 8 \text{ and } 2 \sin y = -2 \\
 \sin y &= 4 \text{ and } \sin y = -1 \\
 \text{Reject } \sin y &= 4 \\
 \text{A solution of } \sin y &= -1 \text{ is } y = \frac{3\pi}{2} \\
 \text{The period of } f(y) = \sin y &\text{ is } 2\pi \\
 y &= 2\pi n + \frac{3\pi}{2}, n \in \mathbb{Z}
 \end{aligned} \tag{9}$$

For the last equation:

$$\begin{aligned}
 \sin 4z + 2 &= \pm 3 \\
 \sin 4z + 2 &= 3 \text{ and } \sin 4z + 2 = -3 \\
 \sin 4z &= 1 \text{ and } \sin 4z = -5 \\
 \text{Reject } \sin 4z &= -5 \\
 \text{Let } q &= 4z \\
 \text{When } \sin q &= 1, q = \frac{\pi}{2} \\
 \text{The period of } f(q) = \sin q &\text{ is } 2\pi \\
 q &= 2\pi n + \frac{\pi}{2}, n \in \mathbb{Z} \\
 4z &= 2\pi n + \frac{\pi}{2}, n \in \mathbb{Z} \\
 z &= \frac{2\pi n + \frac{\pi}{2}}{4}, n \in \mathbb{Z} \\
 z &= \frac{\pi n}{2} + \frac{\pi}{8}, n \in \mathbb{Z}
 \end{aligned} \tag{10}$$

$$\begin{aligned}
 \therefore x &= \pi n + \frac{\pi}{2}, n \in \mathbb{Z} \\
 y &= 2\pi n + \frac{3\pi}{2}, n \in \mathbb{Z} \\
 z &= \frac{\pi n}{2} + \frac{\pi}{8}, n \in \mathbb{Z}
 \end{aligned}$$