

Averaged Jacobian Regularity Can Misrank Few-Step Flow-Matching Schedules: A Certified Gaussian Counterexample

Clément Callaert
CentraleSupélec and Université Paris-Saclay
callaert.clement@gmail.com

Abstract

Few-step sampling in flow matching and stochastic interpolants requires choosing an interpolation schedule before endpoint error is observed. Averaged squared Lipschitzness of the drift, measured by the time-integrated squared 2-norm of its spatial Jacobian, is a plausible schedule-design criterion, but it is not a solver-specific discretization-error functional. Chen, Vanden-Eijnden, and Xu propose minimizing this quantity, denoted A_2 ; they do not claim or prove that A_2 universally ranks equal-NFE solver error. We study that narrower surrogate question. For independent $N(0, 1)$ and $N(0, 4)$, the exact regularity integrals are $5\pi/8 - 1$ and $\pi^2/16$, so regularity prefers trigonometric VP, whereas explicit Heun, a two-stage Runge–Kutta method, with a budget of eight function evaluations (NFE 8), prefers the linear path in Gaussian Wasserstein-2 distance. The inversion is certified by an exact rational Heun product and a nonnegative element of $\mathbb{Q}[\pi, \sqrt{2}]$. A complementary construction shows that, for every step count N and every admissible endpoint log-scale, the unique integrated-regularity minimizer can have strictly larger N -step Euler endpoint error than a higher-regularity competitor aligned with that solver grid. A specified finite Gaussian enumeration of four candidate paths on 36 tested blocks contains both agreement and pairwise disagreement. These results limit a universal surrogate interpretation; they do not evaluate learned velocity fields or estimate a population failure frequency.

1 Introduction

A few-step sampler commits to a probability path, a numerical solver, and a function-evaluation budget before any endpoint error is observed. Few-step sampling therefore makes the sampling schedule a design choice that must be made a priori. Classical one-step bounds involve a Lipschitz constant of the field [Hairer et al., 1993]. A scalar functional of the path that can be evaluated before discretization is an attractive surrogate.

In the stochastic interpolants and flow matching setting, Chen et al. [2025] propose minimizing averaged squared Lipschitzness of the drift (their Definition 3.2, denoted A_2) as a schedule-design objective. They prove KL equivalence of scalar schedules after a posteriori diffusion tuning, a transfer formula, and a logarithmic scaling law for the one-dimensional log-covariance minimizer. They minimize A_2

in specified admissible families. They do not claim, and do not prove, a universal pairwise implication of the form

$$R(q_1) < R(q_2) \implies \text{Err}_{\text{solver},N}(q_1) < \text{Err}_{\text{solver},N}(q_2). \quad (1)$$

The present paper studies that narrower ranking question on flow-matching marginal interpolant ODEs, not score-based probability-flow ODEs.

The object is an integrated unsigned Jacobian-energy functional. For a fixed finite-step solver, endpoint error depends on stagewise field values, temporal derivatives, quadrature structure, and signed cancellations that are not determined by that energy alone.

Already for independent $N(0, 1)$ and $N(0, 4)$, exact regularity integrals prefer trigonometric VP while explicit Heun at NFE 8 prefers the linear path in Gaussian W_2 (Proposition 2). The comparison uses exact W_2 between Gaussians and a fully certified scalar calculation.

For every step count N and every admissible endpoint log-scale $L > 0$, a solver-grid-aware Gaussian construction exhibits two smooth scalar paths with identical endpoints such that the unique minimizer of integrated squared field energy has strictly larger N -step Euler endpoint error than a higher-regularity competitor (Theorem 1). The competitor is constructed as a function of the solver resolution and exploits exact grid alignment. It is not offered as a model of naturally occurring or learned schedules.

A specified finite Gaussian enumeration then delimits where pairwise disagreement appears and where it does not (Section 6). Among four specified candidate paths, the per-mode log-covariance path has both the smallest $\mathcal{R}$ and the smallest W_2 in 36 of 36 tested blocks. Pairwise linear-versus-VP and VP-versus-scalar-log-covariance comparisons on those tested blocks contain inversions. These counts are a finite enumeration, not a population frequency and not a global optimality statement.

Scope. The analysis uses commuting affine Gaussian interpolants, three fixed-step Runge–Kutta methods, and small equal-NFE budgets. No learned neural velocity field is evaluated. The paper does not claim that average Jacobian regularity is generally useless, that it is empirically inferior in typical flow-matching training, that Chen et al.’s optimization results are invalid, or that ranking inversions are frequent in practice. Signed information is consistent with the certified inversion; it is not isolated as a unique causal source.

The contributions are as follows.

- A certified one-dimensional Gaussian ranking inversion: exact $\mathcal{R}$ prefers trigonometric VP while Heun at NFE 8 prefers the linear path in Gaussian W_2 (Proposition 2, Appendix A).
- Exact one-dimensional identities and solver-specific local truncation expansions for the implemented Euler and Heun factors (Section 4, Appendix B).
- A solver-grid-aware impossibility theorem: the unique integrated squared-energy minimizer need not minimize fixed-grid Euler endpoint error (Theorem 1).

- A finite Gaussian enumeration of four specified candidate paths on 36 tested blocks, containing both agreement and pairwise disagreement (Section 6).

2 Definitions and standing assumptions

Let $X_0 \sim N(\mu_0, \Sigma_0)$ and $X_1 \sim N(\mu_1, \Sigma_1)$ be independent in $\mathbb{R}^d$, with Σ_0, Σ_1 symmetric positive definite. A *shared schedule* is a pair of scalar functions used for every eigenmode, equivalently $Q(t) = \alpha(t)^2 \Sigma_0 + \sigma(t)^2 \Sigma_1$. We distinguish three regularity classes.

- *Class S* (shared C^1 interpolants): $\alpha, \sigma \in C^1([0, 1])$ with

$$\alpha(0) = 1, \quad \sigma(0) = 0, \quad \alpha(1) = 0, \quad \sigma(1) = 1, \quad (2)$$

and $(\alpha(t), \sigma(t)) \neq (0, 0)$ for every $t \in [0, 1]$. The linear path $(1 - t, t)$ and trigonometric VP $(\cos(\pi t/2), \sin(\pi t/2))$ belong to Class S, as does the grid-aware embedding of Appendix C.

- *Class Q* (regular covariance paths): a C^1 path of positive definite covariances $Q(t)$ with $Q(0) = \Sigma_0$, $Q(1) = \Sigma_1$, for which $A(t) = \frac{1}{2}Q'(t)Q(t)^{-1}$ extends continuously to $[0, 1]$. Every Class S interpolant induces a Class Q path, because the simultaneous-zero prohibition and $\Sigma_0, \Sigma_1 \succ 0$ give $Q(t) \succ 0$ on $[0, 1]$. The per-mode schedule $q_i(t) = \lambda_i^t$ is Class Q.
- *Class R* (regular squares): $\alpha^2, \sigma^2 \in C^1([0, 1])$, $\alpha^2 \geq 0$, $\sigma^2 \geq 0$, and

$$\alpha(0)^2 = 1, \quad \sigma(0)^2 = 0, \quad \alpha(1)^2 = 0, \quad \sigma(1)^2 = 1, \quad (3)$$

with $\alpha(t)^2 + \sigma(t)^2 > 0$ on $[0, 1]$. Square roots need not be differentiable at the endpoints. Chen et al.'s scalar log-covariance schedule $\sigma(t)^2 = (M^t - 1)/(M - 1)$, $\alpha(t)^2 = (M - M^t)/(M - 1)$ for $M > 0$, $M \neq 1$ is Class R: $\sigma(t) \asymp \sqrt{t}$ as $t \rightarrow 0$ and $\alpha(t) \asymp \sqrt{1 - t}$ as $t \rightarrow 1$, so it is not Class S. When $M = 1$ the formulas are read in the limit $\sigma(t)^2 = t$, $\alpha(t)^2 = 1 - t$. The covariance is C^∞ and A is continuous, so the path is Class Q.

Two paths are called shared-schedule Class Q paths when both lie in Class Q and both use a single scalar pair (α, σ) . We do not call a Class R path a Class S interpolant.

The interpolant of a Class S or Class R schedule is

$$X_t = \alpha(t)X_0 + \sigma(t)X_1, \quad (4)$$

as in stochastic interpolants and flow matching [Albergo et al., 2025, Lipman et al., 2023, Liu et al., 2023, Tong et al., 2024]. Throughout, $\alpha^2 + \sigma^2$ is not required to equal 1 except on the VP path and on Chen et al.'s scalar log-covariance schedule. The two classical paths compared below are the linear schedule $(\alpha, \sigma) = (1 - t, t)$ and the trigonometric VP schedule $(\cos(\pi t/2), \sin(\pi t/2))$.

The law of X_t is Gaussian with mean $m_t = \alpha\mu_0 + \sigma\mu_1$ and covariance $Q(t) = \alpha^2\Sigma_0 + \sigma^2\Sigma_1$. We assume $Q(t) \succ 0$ on $[0, 1]$. The marginal velocity of (4) is the affine field

$$b(t, x) = A(t)x + c(t), \quad A(t) = C_t Q(t)^{-1}, \quad c(t) = \dot{m}_t - A(t)m_t, \quad (5)$$

with $C_t = \frac{1}{2}(\alpha^2)' \Sigma_0 + \frac{1}{2}(\sigma^2)' \Sigma_1$, which extends continuously to $[0, 1]$ whenever $\alpha^2, \sigma^2 \in C^1([0, 1])$. On $(0, 1)$ this coincides with $C_t = \dot{\alpha} \alpha \Sigma_0 + \dot{\sigma} \sigma \Sigma_1$ wherever α and σ are differentiable and nonzero. This is a marginal interpolant ODE, not a score-based probability-flow ODE in the sense of Song et al. [2021]. If $\mu_0 = \mu_1 = 0$ and $\Sigma_0 = I$, then $c \equiv 0$ and $A = \frac{1}{2} Q' Q^{-1}$.

If $\Sigma_0 = I$, then $Q(t)$, $Q'(t)$, $A(t)$, and Σ_1 share the eigenbasis of Σ_1 . In that basis, mode i with eigenvalue $\lambda_i > 0$ obeys $x'_i = a_i(t)x_i + c_i(t)$ with $q_i(t) = \alpha(t)^2 + \lambda_i \sigma(t)^2$ and $a_i(t) = q'_i(t)/(2q_i(t))$. For a Class Q path that is not shared, the same identity holds modewise with no single pair (α, σ) . The exact one-step factor on $[t_n, t_{n+1}]$ is $\sqrt{q_i(t_{n+1})/q_i(t_n)}$. For centered modes the exact endpoint factor is $\sqrt{\lambda_i}$.

Euler uses $s = 1$ field evaluation per step, Heun $s = 2$, and classical RK4 $s = 4$. A budget B is admitted only if $B \equiv 0 \pmod{s}$. The step count is $S = B/s$ and the grid is the uniform partition of $[0, 1]$ with S steps. Equal NFE is not equal step count: at $B = 8$, Euler takes 8 steps, Heun 4, and RK4 2. A solver applied to (5) produces an affine endpoint map $x \mapsto Rx + d$. Error is the Gaussian W_2 of the endpoint laws [Gelbrich, 1990, Peyré and Cuturi, 2019],

$$W_2^2 = \|\hat{m} - \mu_1\|_2^2 + \text{tr}(\hat{C} + \Sigma_1 - 2(\Sigma_1^{1/2} \hat{C} \Sigma_1^{1/2})^{1/2}). \quad (6)$$

In the centered commuting case, $W_2^2 = \sum_i (|r_i| - \sqrt{\lambda_i})^2$.

Chen et al. [2025, Definition 3.2] introduce

$$A_2 = \int_0^1 \mathbb{E}[\|\nabla b_t(I_t)\|_2^2] dt. \quad (7)$$

For the affine field (5) the integrand is deterministic and

$$\mathcal{R}[b] = \int_0^1 \|A(t)\|_2^2 dt \quad (8)$$

coincides with A_2 . By definition $\mathcal{R}$ ignores $c(t)$. For commuting $A(t)$, $\|A(t)\|_2 = \max_i |a_i(t)|$. Headline comparisons use the continuous integral (8). The historical 24-node trapezoidal estimator is written $\hat{\mathcal{R}}_{24}$ and is used only as an implementation check.

Proposition 1. *For any Class Q one-dimensional variance path from 1 to $\lambda > 0$, $\int_0^1 a(t) dt = \frac{1}{2} \ln \lambda$. Consequently*

$$\mathcal{R} = \int_0^1 a(t)^2 dt = \left(\int_0^1 a(t) dt \right)^2 + \text{Var}_t(a) \geq \left(\frac{1}{2} \ln \lambda \right)^2, \quad (9)$$

with equality if and only if a is constant.

Proof. The integral identity is $\frac{1}{2} \log q(1) - \frac{1}{2} \log q(0)$. Expanding $\int a^2$ as $(\int a)^2 + \text{Var}_t(a)$ gives the bound. $\square$

Thus $\mathcal{R}$ measures only the temporal variance of the log-covariance rate. Its unconstrained Class Q minimizer is $a \equiv \frac{1}{2} \ln \lambda$, i.e. $q(t) = \lambda^t$, which is Chen et al. [2025, Example 3.3] in one dimension. In several dimensions the per-mode schedule $q_i(t) = \lambda_i^t$ attains $\mathcal{R} = (\frac{1}{2} \max_i |\ln \lambda_i|)^2$. If Σ_1 has at least two distinct eigenvalues,

that path is not a Class S shared interpolant: matching $\alpha^2 + \lambda_i \sigma^2 = \lambda_i^t$ on two distinct eigenvalues forces $\sigma(t) \asymp \sqrt{t}$ as $t \rightarrow 0$. If three or more distinct eigenvalues are present, no shared (α, σ) exists even in Class R, because $\alpha^2 + \lambda \sigma^2$ is affine in λ while λ^t is not. Isotropic spectra remain compatible with a shared schedule. We do not claim that $d \geq 2$ forbids a shared schedule.

3 Certified scalar Gaussian ranking inversion

Assumption 1. *The endpoints $X_0 \sim N(0, 1)$ and $X_1 \sim N(0, 4)$ are independent. The compared schedules are linear and trigonometric VP, both in Class S. The solver is explicit Heun with NFE $B = 8$, hence $S = 4$ steps of size $h = 1/4$. Error is Gaussian W_2 of the endpoint laws, which equals $|r - 2|$ because every Heun factor below is positive.*

Proposition 2. *Under Assumption 1,*

$$\mathcal{R}_{\text{lin}} = \frac{5\pi}{8} - 1, \quad \mathcal{R}_{\text{VP}} = \frac{\pi^2}{16}, \quad (10)$$

and the comparison $\mathcal{R}_{\text{lin}} > \mathcal{R}_{\text{VP}}$ is equivalent to $2 < \pi < 8$. Averaged regularity therefore prefers VP. The Heun endpoint factors satisfy

$$r_{\text{lin}} = \frac{6797469}{3559400}, \quad 0 < r_{\text{VP}} < \frac{187}{100}, \quad (11)$$

the second bound being a rational enclosure obtained by substituting $\sqrt{2} < 99/70$ and $\pi < 355/113$ into the exact element of $\mathbb{Q}[\pi, \sqrt{2}]$. Consequently

$$W_2^{\text{lin}} = \frac{321331}{3559400} < 0.091 < 0.130 < W_2^{\text{VP}}, \quad (12)$$

so equal-NFE Gaussian W_2 prefers the linear path. Pairwise ordering by integrated Jacobian regularity therefore need not agree with fixed-NFE Heun W_2 error.

Proof sketch. The linear drift is $a(t) = (5t - 1)/(1 - 2t + 5t^2)$. Completing the square and using $\arctan 2 + \arctan(1/2) = \pi/2$ yields $\mathcal{R}_{\text{lin}} = 5\pi/8 - 1$. The VP integral evaluates to $\pi^2/16$. Multiplying $5\pi/8 - 1 > \pi^2/16$ by 16 rearranges to $\pi^2 - 10\pi + 16 < 0$, i.e. $(\pi - 2)(\pi - 8) < 0$, which holds because $2 < \pi < 8$. The four linear Heun factors are the rationals $47/52$, $6/5$, $497/370$, and $97/74$, whose product is $6797469/3559400$. Then $321331/3559400 < 91/1000$ is an elementary comparison of integers. The VP grid drifts are exact elements of $\mathbb{Q}[\pi, \sqrt{2}]$; the four-step Heun product has only nonnegative coefficients, so substituting the rational upper bounds is valid and gives $r_{\text{VP}} < 187/100$. Details occupy Appendix A. $\square$

One certified inversion proves the logical non-implication. It does not by itself identify a unique causal mechanism, and it does not refute a universal pairwise theorem that Chen et al. did not state.




Figure 1: Certified scalar counterexample for independent $N(0, 1)$ and $N(0, 4)$. (a) Closed-form drifts; the shaded region is $a_{\text{lin}} < 0$; markers are Heun nodes at NFE 8. (b) Exact regularity integrals $5\pi/8 - 1$ and $\pi^2/16$. (c) Gaussian W_2 at Heun NFE 8: the linear value is the rational $321331/3559400$; VP is shown by its certified lower enclosure.

4 Solver-specific local error

The leading exact-minus-method coefficients of the one-step factor on $x' = a(t)x$ are $\frac{1}{2}(a' + a^2)h^2$ for explicit Euler and $(-a''/12 + a^3/6)h^3$ for the implemented Heun factor $1 + (h/2)(a(t) + a(t+h)(1 + ha(t)))$ (Appendix B). These involve different derivatives, signs, and stage locations. The scalar integral of $\|A\|_2^2$ does not determine these derivatives, signs, or stagewise values. The expansions establish that local solver error depends on temporal and stagewise information not determined by integrated Jacobian energy. They do not isolate sign loss as the unique source of the certified inversion.

The telescoping identity

$$\prod_{n=0}^{S-1} r_{i,n} - \prod_{n=0}^{S-1} e_{i,n} = \sum_{n=0}^{S-1} (r_{i,n} - e_{i,n}) \left(\prod_{j<n} r_{i,j} \right) \left(\prod_{j>n} e_{i,j} \right) \quad (13)$$

converts those local coefficients into endpoint factor error. The exact modal factor is $e_{i,n} = \sqrt{q_i(t_{n+1})/q_i(t_n)}$; $r_{i,n}$ is numerical. For any independent linear coupling, $a_{\text{lin}}(0) = -1 < 0 < a_{\text{lin}}(1) = +1$. The VP drift vanishes at the endpoints and does

not change sign on $(0, 1)$. This distinction is consistent with the Heun inversion; it is not a causal isolation.

5 Grid-aware impossibility

Proposition 2 compares two fixed Class S paths. The following theorem addresses a different question: even the global minimizer of integrated squared field energy need not minimize fixed-grid Euler error. The competitor is chosen after the grid.

Theorem 1. *Fix $L > 0$ and an integer $N \geq 1$. Let $h = 1/N$ and let forward Euler use the left-endpoint grid $t_n = nh$ on $[0, 1]$. Set $\varepsilon_N = N(e^{L/N} - 1) - L > 0$ and define $a_0(t) \equiv L$ and $a_{1,N}(t) = L + \varepsilon_N \cos(2\pi Nt)$. Consider the scalar ODE $x' = a(t)x$ with $x(0) = 1$, and the Class S Gaussian interpolants of Appendix C with endpoints $N(0, 1)$ and $N(0, e^{2L})$. Then: both exact endpoints equal e^L ; a_0 uniquely minimizes $\int_0^1 a^2$ among scalar fields with integral L ; $\int_0^1 a_{1,N}^2 = L^2 + \varepsilon_N^2/2 > \int_0^1 a_0^2$; left-endpoint Euler is exact for $a_{1,N}$; and Euler has strictly positive endpoint error for a_0 . Endpoint error may equivalently be read as $|r - e^L|$ or as Gaussian W_2 between $N(0, r^2)$ and $N(0, e^{2L})$.*

Proof. The solution of $x' = a(t)x$, $x(0) = 1$, is $x(t) = \exp(\int_0^t a(s) ds)$. The cosine perturbation has integer period, so $\int_0^1 \cos(2\pi Nt) dt = 0$ and both exact endpoints equal e^L . Expanding the square and using the same orthogonality gives $\int_0^1 a_{1,N}^2 = L^2 + \varepsilon_N^2/2$. The inequality $\varepsilon_N > 0$ is $e^z > 1 + z$ at $z = L/N > 0$. Uniqueness of the minimizer is Proposition 1. On the left-endpoint Euler grid, $\cos(2\pi Nt_n) = \cos(2\pi n) = 1$, so $1 + ha_{1,N}(t_n) = e^{L/N}$ and the product is e^L . For the constant field the product is $(1 + L/N)^N < e^L$ because $\log(1 + z) < z$ for $z = L/N > 0$. The Class S embedding is Appendix C. $\square$

The field $a_{1,N}$ is chosen after the grid. Shifted nodes, phase offsets, and frequencies other than N destroy exact Euler aliasing (Table 8 in Appendix G). Heun is not exact on the aligned oscillation. The construction is a solver-grid-dependent impossibility result, not evidence that naturally occurring schedules typically behave this way.

6 Finite Gaussian enumeration

Source $N(0, I)$. Four centered targets, with seed 271828 for the headline factor matrices: anisotropic Gaussians with condition number 4 in dimensions 2 and 8 (eigenvalues a geometric sequence from $1/2$ to 2); and factor-plus-noise covariances $\Sigma = FF^\top + 0.05I$ with $F \in \mathbb{R}^{d \times 2}$ having i.i.d. standard-normal entries. When $d = 2$ the factor has rank equal to the ambient dimension; when $d = 8$ it is rank 2 plus a noise floor. The dimension-2 eigenvalues are 0.291 and 2.97; the dimension-8 eigenvalues are 0.05 (six times), 4.65, and 10.7. Paths, solvers, budgets, and status occupy Table 1. Precision is float64, audited against an 80-digit Gelbrich reference. Phase 4 is confirmatory on a grid frozen before the rerun, not a protocol recorded before discovery.

Given a geometry, a path, a solver, and a budget, $\mathcal{R}$ and W_2 are deterministic numerical quantities. The 36-block grid is a finite enumeration of 4 geometries $\times$ 3

Table 1: Experimental grid. Headline $\mathcal{R}$ is the continuous integral. “Confirmatory” means reproduced on a frozen grid; “post-hoc” means added after the Phase 3 observation. The Gaussian drift and W_2 formula are closed form; multimode $\mathcal{R}$ is adaptive quadrature.

experiment	grid	paths	status
pairwise comparator	4 geom. $\times$ 3 solvers $\times$ {8, 16, 32}	linear, VP	confirmatory
shared-schedule pairwise	same 36 tested blocks	VP, Ex. 3.3 scalar	post-hoc
four-path enumeration	same 36 tested blocks	$+q_i = \lambda_i^t$	post-hoc
inversion map	$\lambda \in [0.05, 100] \times 3 \times 3$	linear, VP	post-hoc
geometry draws	$N = 50$ i.i.d. $F, d \in \{2, 8\}$	linear, VP	post-hoc
robustness $\pm 10\%$, NFE 64, 128	66 factor-plus-noise blocks	linear, VP	post-hoc
non-centered stress test	$2 \times 3 \times 3$	linear, VP	confirmatory

solvers $\times$ 3 budgets. Because $\mathcal{R}$ is independent of solver and budget, and because the two anisotropic geometries share one $\mathcal{R}$ pair, the enumeration contains 3 distinct linear-versus-VP $\mathcal{R}$ comparisons and nests into 12 geometry $\times$ solver cells. Each inversion margin exceeds the 80-digit floor by a factor $\geq 10^3$. No population-level inference is intended.

Linear versus VP invert in 5 of 12 geometry $\times$ solver cells; 4 of those cells invert at every NFE in {8, 16, 32}. On this enumeration the paired concordance score $(n_{\text{agree}} - n_{\text{invert}})/n$ equals 2/9 over 36 tested blocks. Stratified by solver ($n = 12$ blocks per stratum) the score is +1/2 under Euler and RK4 and $-1/3$ under Heun. The strongest block is rank-2 factor-plus-noise $d = 8$, Euler, NFE 8: $\mathcal{R}$ prefers linear (2.944 versus 4.731) while W_2 prefers VP (0.811 versus 0.456).

As a post-hoc pairwise comparison of two shared-schedule Class Q paths, trigonometric VP versus Chen Example 3.3 with $M = \lambda_{\max}$ inverts in 9 of 36 tested blocks and 4 of 12 cells. This is not a test of Chen et al. [2025, Proposition 3.9].

Among the four specified candidate paths on the same 36 tested blocks, the per-mode log-covariance path has the smallest continuous $\mathcal{R}$ and the smallest Gaussian W_2 in 36 of 36 tested blocks. When Σ_1 has at least two distinct eigenvalues, that path is not a Class S shared interpolant. Theorem 1 blocks any reading of this count as a global minimizer theorem.

A log-uniform sweep $\lambda \in [0.05, 100]$ locates linear-versus-VP inversions (Figure 4). At $\lambda = 4$ the inversion is present under Heun at every tested budget NFE $\in \{4, 8, 12, 16, 24, 32, 64, 128, 256\}$, absent under Euler, and mixed under RK4. Euler does invert at more extreme λ . The scalar example is solver-dependent, not an artefact of Heun at budget 8 alone.

Redrawing F from `torch.Generator` seeds 10000 through 10049 at $d \in \{2, 8\}$, every one of the 50 draws has at least one linear-versus-VP inversion among the nine solver-budget conditions. This descriptive fraction is reported only for the finite sampled geometries; no population-level inference is intended. The event is a post-hoc “at least one inversion” summary, not a preregistered geometry-level law.

Mixture targets are excluded. A confirmatory non-centered family (Appendix E) shows that $\mathcal{R}$, which ignores $c(t)$, is blind to a mean contribution in W_2^2 .




Figure 2: Three comparisons on commuting Gaussian interpolants. Continuous $\mathcal{R}$ (left) and equal-NFE Gaussian W_2 (right). (a) Pairwise linear versus VP on rank-2 factor-plus-noise $d = 8$, Euler, NFE 8. (b) VP versus Chen Ex. 3.3 scalar on anisotropic $d = 8$, Euler, NFE 8. (c) Four specified candidate paths on rank-2 factor-plus-noise $d = 8$, Euler, NFE 8. Among those four paths, the per-mode path has the smallest $\mathcal{R}$ and the smallest W_2 in 36 of 36 tested blocks.

7 Related work

Flow matching and stochastic interpolants supply the Gaussian path family [Lipman et al., 2023, Albergo et al., 2025, Liu et al., 2023, Tong et al., 2024]. Lipschitz-guided schedule design proposes the selection criterion tested here [Chen et al., 2025]: minimize A_2 over scalar schedules, with a transfer formula that applies a designed schedule at inference without retraining. The present negative result is a limitation of a universal pairwise surrogate reading, not a refutation of that optimization programme.

Tao and Choi [2026] give a broader dynamical error analysis, decomposing $\nabla v = S + \Omega$ and controlling Euler error by the logarithmic norm $\mu_2 = \lambda_{\max}(S)$, together with material-derivative structure and learned-model experiments. In one dimension $\mu_2(a) = a$ and $\|A\|_2 = |a|$. Their Theorem 5 concerns McCann displacement interpolation, which is not the independent-coupling linear or VP path studied here. The present contribution is narrower: an exact, fully certified scalar Gaussian ranking inversion, a closed-form W_2 comparison, a solver-grid-aware impossibility construction, and a transparent finite Gaussian enumeration. It does not provide a general numerical-error theory.




Figure 3: Four specified candidate paths on the four centered geometries. Left: continuous $\mathcal{R}$. Right: Euler Gaussian W_2 at NFE 8.




Figure 4: Linear versus VP ranking inversion versus target variance λ . Dark cells: $\mathcal{R}$ and W_2 disagree. The dashed line is $\lambda = 4$ (Proposition 2).

Stéphanovitch [2026] proves sharp Lipschitz regularity of canonical flow-matching fields and Euler-type W_2 rates of order $\sqrt{d}/N$ up to logs. That is a bound, not a pairwise ranking of two interpolants at equal NFE. Solver-specific schedules and exact linear parts are documented in diffusion practice [Sabour et al., 2024, Karras et al., 2022, Lu et al., 2022, Zhang and Chen, 2023]. Benton et al. [2024] bound W_2 of a learned flow. Those objects are distinct from the ranking question tested here.

Complementary solver-aware approaches use temporal information rather than an integrated spatial-Jacobian energy. Gupta et al. [2026] construct SharpEuler, calibrating Euler grids from trajectory acceleration estimated on pretrained flow-matching models. Khan [2026] regularizes material acceleration during training. Other work modifies the coupling or prior design to produce straighter, OT-like flows [Malnick et al., 2026]. The present contribution is neither a sampler nor a training regularizer: it gives a certified ranking non-implication and a solver-grid-

aware impossibility result for the stated classes.

8 Limitations

The systems are commuting affine Gaussian interpolants. No learned neural field, image, video, or world-model quantity is reported. The solvers are three fixed-step Runge–Kutta methods at small equal-NFE budgets. Exponential integrators solve affine scalar modes exactly, so the comparisons concern generic RK discretizations of those modes rather than the best integrator for an affine field. Theorem 1 depends on the grid: the competitor is chosen after the grid and exploits exact aliasing. Census counts are hierarchical enumerations of specified candidates, not population frequencies. No counterexample to the signed logarithmic norm μ_2 is provided. Sign loss is consistent with the Heun inversion and is not isolated as its unique cause.

9 Conclusion

An integrated unsigned Jacobian-energy functional does not contain enough solver-specific temporal information to universally determine few-step endpoint error. A certified scalar Gaussian example shows that pairwise ordering by integrated Jacobian regularity can disagree with fixed-NFE Heun W_2 error. Local solver error depends on temporal and stagewise information not determined by that energy. A solver-grid-aware Gaussian construction shows that the global integrated-regularity minimizer need not minimize fixed-grid Euler error. A specified finite Gaussian enumeration contains both agreement and disagreement. The results establish a limitation of a universal surrogate interpretation.

References

- Michael S. Albergo, Nicholas M. Boffi, and Eric Vanden-Eijnden. Stochastic interpolants: A unifying framework for flows and diffusions. *Journal of Machine Learning Research*, 26 (209):1–80, 2025. arXiv:2303.08797.
- Joe Benton, George Deligiannidis, and Arnaud Doucet. Error bounds for flow matching methods. *Transactions on Machine Learning Research*, 2024. arXiv:2305.16860.
- Yifan Chen, Eric Vanden-Eijnden, and Jiawei Xu. Lipschitz-guided design of interpolation schedules in generative models. *arXiv preprint arXiv:2509.01629*, 2025. version 3, 16 May 2026.
- Matthias Gelbrich. On a formula for the L^2 Wasserstein metric between measures on Euclidean and Hilbert spaces. *Mathematische Nachrichten*, 147:185–203, 1990. doi: 10.1002/mana.19901470121.
- Aditi Gupta, Soon Hoe Lim, Annan Yu, and N. Benjamin Erichson. Sharpen your flow: Sharpness-aware sampling for flow matching. *arXiv preprint arXiv:2605.11547*, 2026.
- Ernst Hairer, Syvert P. Nørsett, and Gerhard Wanner. *Solving Ordinary Differential Equations I: Nonstiff Problems*. Springer, 2nd edition, 1993.
- Tero Karras, Miika Aittala, Timo Aila, and Samuli Laine. Elucidating the design space of diffusion-based generative models. In *Advances in Neural Information Processing Systems*, 2022. arXiv:2206.00364.

- Tauhid Khan. Isokinetic flow matching for pathwise straightening of generative flows. *arXiv preprint arXiv:2604.04491*, 2026.
- Yaron Lipman, Ricky T. Q. Chen, Heli Ben-Hamu, Maximilian Nickel, and Matthew Le. Flow matching for generative modeling. In *International Conference on Learning Representations (ICLR)*, 2023. arXiv:2210.02747.
- Xingchao Liu, Chengyue Gong, and Qiang Liu. Flow straight and fast: Learning to generate and transfer data with rectified flow. In *International Conference on Learning Representations (ICLR)*, 2023. arXiv:2209.03003.
- Cheng Lu, Yuhao Zhou, Fan Bao, Jianfei Chen, Chongxuan Li, and Jun Zhu. DPM-solver: A fast ODE solver for diffusion probabilistic model sampling in around 10 steps. In *Advances in Neural Information Processing Systems*, 2022. arXiv:2206.00927.
- Shimon Malnick, Matan Rusanovsky, Ohad Fried, and Shai Avidan. Optimal transport flow matching by design. *arXiv preprint arXiv:2606.04092*, 2026.
- Gabriel Peyré and Marco Cuturi. *Computational Optimal Transport*. Now Publishers, 2019. arXiv:1803.00567.
- Amirmojtaba Sabour, Sanja Fidler, and Karsten Kreis. Align your steps: Optimizing sampling schedules in diffusion models. In *International Conference on Machine Learning (ICML)*, 2024. arXiv:2404.14507.
- Yang Song, Jascha Sohl-Dickstein, Diederik P. Kingma, Abhishek Kumar, Stefano Ermon, and Ben Poole. Score-based generative modeling through stochastic differential equations. In *International Conference on Learning Representations (ICLR)*, 2021. arXiv:2011.13456.
- Arthur Stéphanovitch. Lipschitz regularity in flow matching and diffusion models: Sharp sampling rates and functional inequalities. *arXiv preprint arXiv:2604.06065*, 2026.
- Chenxi Tao and Seung-Kyum Choi. On the role of strain and vorticity in numerical integration error for flow matching. *arXiv preprint arXiv:2605.06680*, 2026.
- Alexander Tong, Kilian Fatras, Nikolay Malkin, Guillaume Hugué, Yanlei Zhang, Jarrod Rector-Brooks, Guy Wolf, and Yoshua Bengio. Improving and generalizing flow-based generative models with minibatch optimal transport. *Transactions on Machine Learning Research*, 2024. arXiv:2302.00482.
- Qinsheng Zhang and Yongxin Chen. Fast sampling of diffusion models with exponential integrator. In *International Conference on Learning Representations (ICLR)*, 2023. arXiv:2204.13902.

A Proof of Proposition 2

Linear drift and integral. $q_{\text{lin}}(t) = (1-t)^2 + 4t^2 = 1 - 2t + 5t^2$ and $q' = -2 + 10t$, so $a_{\text{lin}}(t) = (5t-1)/q_{\text{lin}}(t)$. Completing the square yields the two equivalent identities

$$q_{\text{lin}}(t) = 5 \left[\left(t - \frac{1}{5} \right)^2 + \left(\frac{2}{5} \right)^2 \right], \quad (14)$$

$$q_{\text{lin}}(t) = 5 \left(t - \frac{1}{5} \right)^2 + \frac{4}{5}. \quad (15)$$

Thus $a_{\text{lin}}(t) = (t - 1/5) / ((t - 1/5)^2 + (2/5)^2)$. The substitution $u = t - 1/5 = (2/5) \tan \theta$ gives $du = (2/5) \sec^2 \theta d\theta$ and converts $\int a_{\text{lin}}^2$ into $(5/2) \int \sin^2 \theta d\theta$. The definite limits are $\theta(0) = \arctan(-1/2)$ and $\theta(1) = \arctan 2$. The antiderivative $(5/4)(\theta - \sin \theta \cos \theta)$ therefore contributes $(5/4)(\arctan 2 + \arctan(1/2)) - (5/4)(4/5)$. Both angles lie in $(0, \pi/2)$ and $\tan(\arctan 2 + \arctan(1/2))$ is undefined, so the sum is $\pi/2$. Hence $\mathcal{R}_{\text{lin}} = 5\pi/8 - 1$. (The subtracted 1 is the boundary evaluation of $\sin \theta \cos \theta = \tan \theta / (1 + \tan^2 \theta)$.)

VP drift and integral. $q_{\text{VP}}(t) = \cos^2(\pi t/2) + 4 \sin^2(\pi t/2) = (5 - 3 \cos(\pi t))/2$ and $q' = (3\pi/2) \sin(\pi t)$, hence

$$a_{\text{VP}}(t) = \frac{(3\pi/2) \sin(\pi t)}{5 - 3 \cos(\pi t)} = \frac{(3\pi/4) \sin(\pi t)}{q_{\text{VP}}(t)}. \quad (16)$$

The change of variables $u = \pi t$ yields $\mathcal{R}_{\text{VP}} = (9\pi/4) \int_0^\pi \sin^2 u / (5 - 3 \cos u)^2 du$. The standard Weierstrass evaluation of the integrand gives $\int_0^\pi \sin^2 u / (5 - 3 \cos u)^2 du = \pi/36$, hence $\mathcal{R}_{\text{VP}} = \pi^2/16$. The comparison $\mathcal{R}_{\text{lin}} > \mathcal{R}_{\text{VP}}$ rearranges to $\pi^2 - 10\pi + 16 < 0$, i.e. $(\pi - 2)(\pi - 8) < 0$, which holds because $2 < \pi < 8$. Both closed forms were checked by adaptive quadrature (absolute discrepancy below 10^{-12} ; quadrature error estimates are not used as certificates).

Heun recurrence. On $x' = a(t)x$ the implemented Heun factor is $r = 1 + (h/2)(a(t_n) + a(t_{n+1})(1 + ha(t_n)))$. With $h = 1/4$ the four linear factors are the rationals $47/52$, $6/5$, $497/370$, and $97/74$, all positive, with product $6797469/3559400$. Then $2 - r = 321331/3559400$, and $321331 \cdot 1000 < 91 \cdot 3559400$ is an integer comparison, so $W_2^{\text{lin}} < 0.091$.

Exact VP grid and rational certificate. At the Heun nodes the VP drift takes the values

$$\begin{array}{cccccc} t & 0 & 1/4 & 1/2 & 3/4 & 1 \\ a_{\text{VP}}(t) & 0 & \frac{(9+15\sqrt{2})\pi}{82} & \frac{3\pi}{10} & \frac{(-9+15\sqrt{2})\pi}{82} & 0 \end{array} \quad (17)$$

Positivity of $a(3/4)$ follows from $15\sqrt{2} > 9$, or $15^2 \cdot 2 - 9^2 = 369 > 0$ after squaring. Each Heun factor is therefore strictly positive, so $0 < r_{\text{VP}}$ and $W_2^{\text{VP}} = 2 - r_{\text{VP}}$. The four-step product is the following element $P(\pi, \sqrt{2})$ of $\mathbb{Q}[\pi, \sqrt{2}]$, obtained by expanding the four Heun factors and reducing $\sqrt{2}^2 = 2$:

$$\begin{aligned} P(\pi, \sqrt{2}) = & 1 + \frac{3}{40}\pi + \frac{15}{164}\pi\sqrt{2} + \frac{78579}{10758400}\pi^2 + \frac{45}{5248}\pi^2\sqrt{2} \\ & + \frac{11421}{17213440}\pi^3 + \frac{4671}{17213440}\pi^3\sqrt{2} \\ & + \frac{5103}{275415040}\pi^4 + \frac{567}{55083008}\pi^4\sqrt{2} \\ & + \frac{243}{2203320320}\pi^5 + \frac{729}{2203320320}\pi^5\sqrt{2} + \frac{729}{176265625600}\pi^6. \end{aligned} \quad (18)$$

Each of the twelve coefficients is a ratio of nonnegative integers, hence nonnegative, and the constant term is 1. Substituting any pair of upper bounds for π and $\sqrt{2}$ is therefore valid. The bound $\sqrt{2} < 99/70$ is the integer comparison $99^2 - 2 \cdot 70^2 = 1$. The bound $\pi < 355/113$ follows from Machin's formula $\pi = 16 \arctan(1/5) - 4 \arctan(1/239)$ together with Leibniz remainders: seven Taylor terms upper-bound $\arctan(1/5)$ and four terms lower-bound $\arctan(1/239)$. The resulting rational is strictly less than $355/113$ by an integer comparison. Substituting these enclosures yields

$$P\left(\frac{355}{113}, \frac{99}{70}\right) = \frac{192113353671412470139303}{102753427879939708813312} < \frac{187}{100}, \quad (19)$$

equivalently the integer comparison

$$19211335367141247013930300 < 19214891013548725548089344. \quad (20)$$

Hence $r_{\text{VP}} < 187/100$ and $W_2^{\text{VP}} > 13/100$. An independent interval enclosure at 40 decimal digits and an 80-digit non-interval product agree with this bound and with the float64 implementation to 10^{-14} . They are cross-checks, not the proof.

B Solver formulas and local truncation expansions

On $[t, t+h]$ for $x' = a(t)x$,

$$r_{\text{Euler}} = 1 + ha(t), \quad (21)$$

$$r_{\text{Heun}} = 1 + \frac{h}{2}(a(t) + a(t+h)(1 + ha(t))), \quad (22)$$

and classical RK4 with stages at t , $t+h/2$ (twice), and $t+h$. Equal NFE requires $B \equiv 0 \pmod{s}$ and sets $S = B/s$.

Assume a is C^3 . The exact factor admits $1 + ha + (h^2/2)(a' + a^2) + (h^3/6)(a'' + 3aa' + a^3) + O(h^4)$. Subtracting Euler leaves $\frac{1}{2}(a' + a^2)h^2 + O(h^3)$. Substituting $a(t+h)$ into the implemented Heun factor leaves $(-a''/12 + a^3/6)h^3 + O(h^4)$. The aa' terms cancel. The expansion was confirmed with independent symbolic algebra on the implemented expression.

C Class S embedding of the grid-aware fields

The ODE comparison of Theorem 1 is proved in the main text. This appendix records the Class S Gaussian embedding and the positivity hypotheses. The field $a_{1,N}$ is chosen after the grid. The construction is not the mechanism of Proposition 2.

The same fields are Class S Gaussian interpolants. Set $\lambda = e^{2L}$ and

$$\begin{aligned} q_k(t) &= \exp\left(2 \int_0^t a_k(s) ds\right), \\ \alpha_k(t) &= \sqrt{q_k(t)} \cos(\pi t/2), \\ \sigma_k(t) &= \sqrt{q_k(t)/\lambda} \sin(\pi t/2). \end{aligned} \quad (23)$$

Then $q_k(0) = 1$, $\int_0^1 a_k = L$ so $q_k(1) = \lambda$, and the endpoint conditions $\alpha_k(0) = \sigma_k(1) = 1$, $\alpha_k(1) = \sigma_k(0) = 0$ hold. Positivity $q_k > 0$ and C^∞ regularity of a_k give $\alpha_k, \sigma_k \in C^\infty([0, 1])$, hence the embedding belongs to Class S. The identity $\alpha_k^2 + \lambda \sigma_k^2 = q_k$ is trigonometric, and $q'_k/(2q_k) = a_k$ by construction. Among scalar fields with fixed integral L , a_0 uniquely minimizes $\int_0^1 a^2$ (Proposition 1). Euler has strictly positive endpoint error for a_0 , while

$a_{1,N}$ has strictly larger regularity and left-endpoint Euler is exact for $a_{1,N}$. Thus a_0 is an admissible Class S $\mathcal{R}$ -minimizer among one-dimensional variance paths from 1 to λ , yet left-endpoint Euler has strictly larger endpoint error for a_0 than for the higher- $\mathcal{R}$ path $a_{1,N}$. This construction blocks any inference from a 36 of 36 count among four specified candidate paths on 36 tested blocks to the statement that a global $\mathcal{R}$ -minimizer minimizes fixed-NFE error.

D Full block tables

Table 2 lists every inverted linear-versus-VP solver-budget row under continuous $\mathcal{R}$. These rows are a hierarchical listing of the 5 geometry $\times$ solver cells, not a population rate. Table 3 records the strongest centered block. Table 4 reports the four-path enumeration: among four specified candidate paths, the per-mode schedule has the smallest $\mathcal{R}$ and the smallest W_2 in 36 of 36 tested blocks. This finite enumeration does not imply that a global $\mathcal{R}$ -minimizer minimizes fixed-NFE error. When Σ_1 has at least two distinct eigenvalues, that path is not a Class S shared interpolant. Table 5 lists the nine shared-schedule pairwise inversions. Figure 5 illustrates identity (13) on the strongest block; Figure 6 shows the factor-plus-noise solver-path split.

Table 2: All 14 inverted solver-budget rows among 36 centered blocks, grouped into 5 of 12 geometry $\times$ solver cells. Census listing, not a sample.

family	d	solver	NFE	R prefers	W_2 prefers	W_2 margin
anisotropic	2	heun	8	VP	linear	1.905310×10^{-3}
anisotropic	2	heun	16	VP	linear	1.939806×10^{-3}
anisotropic	2	heun	32	VP	linear	6.634439×10^{-4}
anisotropic	8	heun	16	VP	linear	1.758942×10^{-3}
anisotropic	8	heun	32	VP	linear	7.628173×10^{-4}
factor+noise	2	heun	8	VP	linear	1.836251×10^{-2}
factor+noise	2	heun	16	VP	linear	6.039925×10^{-3}
factor+noise	2	heun	32	VP	linear	1.698035×10^{-3}
factor+noise	2	rk4	8	VP	linear	1.516722×10^{-3}
factor+noise	2	rk4	16	VP	linear	1.289981×10^{-4}
factor+noise	2	rk4	32	VP	linear	1.118892×10^{-5}
rank-2 factor+noise	8	euler	8	linear	VP	3.543761×10^{-1}
rank-2 factor+noise	8	euler	16	linear	VP	2.046946×10^{-1}
rank-2 factor+noise	8	euler	32	linear	VP	1.096576×10^{-1}

Table 3: Strongest centered inversion: rank-2 factor-plus-noise, $d = 8$, Euler, NFE 8. $\mathcal{R}$ is the continuous integral; $\widehat{\mathcal{R}}_{24}$ is shown only as a check. Eigenvalues: 0.05 (six times), 4.64827725629505, 10.675638226546441. Deterministic calculation: $\mathcal{R}$ is evaluated by adaptive quadrature, while W_2 uses exact Gaussian moments propagated through the modal Runge–Kutta updates.

path	$\mathcal{R}$	$\widehat{\mathcal{R}}_{24}$	Gaussian W_2
linear	2.9441044083	2.9476523251	0.8108540111
VP	4.7305438136	4.7295206355	0.4564779075

E Non-centered structural stress test

The quantity $\mathcal{R}$ depends only on $A(t)$. If $c(t) \neq 0$, the mean still contributes $\|\hat{m} - \mu_1\|_2^2$ to W_2^2 . Dimensions $d \in \{2, 8\}$, coordinates indexed by $i = 0, \dots, d - 1$. Source $N(\mu_0, I)$ with

Table 4: Four specified candidate paths on the centered grid. Among these four paths, the per-mode schedule has the smallest $\mathcal{R}$ and the smallest W_2 in 36 of 36 tested blocks. This finite enumeration does not imply that a global $\mathcal{R}$ -minimizer minimizes fixed-NFE error. When Σ_1 has at least two distinct eigenvalues, that path is not a Class S shared interpolant.

family	d	solver	NFE	R min	W_2 min	$W_{2,\log}$	$W_{2,\text{lin}}$
anisotropic	2	euler	8	log-cov	log-cov	0.01164	0.2441
anisotropic	2	euler	16	log-cov	log-cov	0.005874	0.1266
anisotropic	2	euler	32	log-cov	log-cov	0.002952	0.06454
anisotropic	2	heun	8	log-cov	log-cov	0.0006613	0.03243
anisotropic	2	heun	16	log-cov	log-cov	0.0001682	0.006327
anisotropic	2	heun	32	log-cov	log-cov	4.244e-05	0.001359
anisotropic	2	rk4	8	log-cov	log-cov	3.834e-06	0.01119
anisotropic	2	rk4	16	log-cov	log-cov	2.473e-07	0.000213
anisotropic	2	rk4	32	log-cov	log-cov	1.575e-08	9.781e-06
anisotropic	8	euler	8	log-cov	log-cov	0.0131	0.4509
anisotropic	8	euler	16	log-cov	log-cov	0.006608	0.2337
anisotropic	8	euler	32	log-cov	log-cov	0.003319	0.1191
anisotropic	8	heun	8	log-cov	log-cov	0.0007027	0.04645
anisotropic	8	heun	16	log-cov	log-cov	0.0001786	0.008457
anisotropic	8	heun	32	log-cov	log-cov	4.503e-05	0.001746
anisotropic	8	rk4	8	log-cov	log-cov	3.897e-06	0.01812
anisotropic	8	rk4	16	log-cov	log-cov	2.513e-07	0.0003324
anisotropic	8	rk4	32	log-cov	log-cov	1.6e-08	1.37e-05
factor+noise	2	euler	8	log-cov	log-cov	0.03303	0.2933
factor+noise	2	euler	16	log-cov	log-cov	0.01683	0.1523
factor+noise	2	euler	32	log-cov	log-cov	0.008497	0.07773
factor+noise	2	heun	8	log-cov	log-cov	0.002998	0.0585
factor+noise	2	heun	16	log-cov	log-cov	0.0007696	0.0122
factor+noise	2	heun	32	log-cov	log-cov	0.0001953	0.002707
factor+noise	2	rk4	8	log-cov	log-cov	4.7e-05	0.004578
factor+noise	2	rk4	16	log-cov	log-cov	2.974e-06	0.000432
factor+noise	2	rk4	32	log-cov	log-cov	1.89e-07	1.732e-05
rank-2 factor+noise	8	euler	8	log-cov	log-cov	0.2736	0.8109
rank-2 factor+noise	8	euler	16	log-cov	log-cov	0.1445	0.424
rank-2 factor+noise	8	euler	32	log-cov	log-cov	0.07439	0.2175
rank-2 factor+noise	8	heun	8	log-cov	log-cov	0.05274	0.4094
rank-2 factor+noise	8	heun	16	log-cov	log-cov	0.01399	0.08936
rank-2 factor+noise	8	heun	32	log-cov	log-cov	0.00363	0.02047
rank-2 factor+noise	8	rk4	8	log-cov	log-cov	0.004721	0.06962
rank-2 factor+noise	8	rk4	16	log-cov	log-cov	0.0002675	0.00746
rank-2 factor+noise	8	rk4	32	log-cov	log-cov	1.687e-05	0.0001322

Table 5: Shared-schedule pairwise inversions: trigonometric VP versus Chen Example 3.3 with $M = \lambda_{\max}$. Nine of 36 blocks; four of 12 cells. Post-hoc pairwise census, not a global family optimum.

family	d	solver	NFE	R prefers	W_2 prefers
anisotropic	2	euler	16	scalar	VP
anisotropic	2	euler	32	scalar	VP
anisotropic	8	euler	8	scalar	VP
anisotropic	8	euler	16	scalar	VP
anisotropic	8	euler	32	scalar	VP
factor+noise	2	euler	8	VP	scalar
factor+noise	2	heun	8	VP	scalar
factor+noise	2	heun	16	VP	scalar
factor+noise	2	heun	32	VP	scalar




Figure 5: Modal contributions to squared Wasserstein error and transported signed defects for the strongest centered inversion. The bars reconstruct the telescoping identity (13). Exact modal identity on one block; illustration, not an independent causal test.




Figure 6: Signed factor-plus-noise W_2 margin at equal NFE. Positive values prefer VP; negative values prefer linear. Vertical axis: symmetric-log with linear threshold 10^{-4} . Exact-moment modal Runge–Kutta.

$\mu_{0,i} = 0.75(-1)^i$. Target $N(\mu_1, D)$ with $\mu_{1,i} = 1 + 0.25i$ and $D = \text{diag}(\lambda_i)$, where the λ_i form a geometric sequence from $6^{-1/2}$ to $6^{1/2}$ (condition number 6). Then $c(t) \neq 0$. Paths are linear and trigonometric VP. Solvers are Euler, Heun, and RK4 at NFE $B \in \{8, 16, 32\}$. Write $M = \|\hat{m} - \mu_1\|_2^2$ and $C = W_2^2 - M$, and

$$\Delta_{\text{mean}} = M_{\text{lin}} - M_{\text{VP}}, \quad \Delta_{\text{cov}} = C_{\text{lin}} - C_{\text{VP}}, \quad \Delta_{\text{total}} = \Delta_{\text{mean}} + \Delta_{\text{cov}}. \quad (24)$$

A block is mean-driven if $\Delta_{\text{mean}}\Delta_{\text{total}} > 0$ and $|\Delta_{\text{mean}}| > |\Delta_{\text{cov}}|$, and covariance-driven if $\Delta_{\text{cov}}\Delta_{\text{total}} > 0$ and $|\Delta_{\text{cov}}| > |\Delta_{\text{mean}}|$. Otherwise it is mixed. The same paths, solvers, and budgets give 18 blocks, of which 11 invert under continuous $\mathcal{R}$. Table 6 reports the driver of each path preference. Several inversions, including all dimension-8 Euler and Heun inversions, are mean-driven: $\mathcal{R}$ cannot see them. The family was specified before execution as a stress test of this structural blind spot, not as an independent replication of the centered inversions.




Figure 7: Signed path differences on all 18 non-centered blocks. $\Delta_{\text{mean}} = M_{\text{lin}} - M_{\text{VP}}$ and $\Delta_{\text{cov}} = C_{\text{lin}} - C_{\text{VP}}$. A positive bar means the linear path has the larger contribution; a negative bar means VP has the larger contribution. The axis is symmetric-log with linear threshold 10^{-6} . Stars mark ranking inversions. Exact numerical values appear in Table 7.

F Quadrature, precision, and interval procedures

Table 9 compares the continuous integral with the historical 24-node trapezoidal estimator on every headline geometry. Path orderings agree. Trapezoidal values at 24, 48, 96, and 192 nodes converge to the same sign. Adaptive quadrature is used for multi-mode $\mathcal{R}$; the scalar integrals of Appendix A are closed form. Scipy error estimates are not interval certificates.

Because $\mathcal{R}$ depends on neither solver nor budget, the 36 linear-versus-VP rows nest into 12 geometry $\times$ solver cells. The two anisotropic geometries share one $\mathcal{R}$ pair, so the census contains three distinct $\mathcal{R}$ comparisons:

geometry	$(\mathcal{R}_{\text{lin}}, \mathcal{R}_{\text{VP}})$	distinct $\mathcal{R}$ pair
anisotropic $d = 2$ and $d = 8$	(0.972, 0.192)	1 (shared)
factor+noise $d = 2$	(1.246, 0.625)	1
rank-2 factor+noise $d = 8$	(2.944, 4.731)	1

Frozen Gaussian W_2 on the centered census was checked against an 80-digit Gelbrich reference. The maximum observed gap is the numerical floor 9.7×10^{-10} . The paired concordance score $(n_{\text{agree}} - n_{\text{invert}})/n$ between the $\mathcal{R}$ order and the W_2 order is a descriptive census statistic, not a test (Table 12). On the 36 linear-versus-VP blocks, the score is 2/9. Stratified by solver ($n = 12$ blocks per stratum) the score is +1/2 under Euler and RK4 and $-1/3$ under Heun. The two anisotropic geometries share one $\mathcal{R}$ pair, a structural tie in the value of $\mathcal{R}$, not in the denominator. Stratified by geometry ($n = 9$ blocks per stratum) the scores are 1/3, 5/9, $-1/3$, and 1/3.

Scalar linear Heun factors are computed in exact rationals. The VP Heun product is certified by substituting $\pi < 355/113$ and $\sqrt{2} < 99/70$ into a nonnegative element of $\mathbb{Q}[\pi, \sqrt{2}]$; the ranking inequality is the integer comparison in Appendix A. An independent interval enclosure at 40 decimal digits (mpmath 1.3.0) and an 80-digit (non-interval) product are recorded as cross-checks against the float64 Heun implementation. Default numerical precision for the benchmark runs is float64.

Table 11 reports the log-Lebesgue measure of the linear-versus-VP inversion set on $[0.05, 100]$. The declared resolution is 1281 log-uniform nodes; doubling to 2561 nodes

Table 6: Non-centered family: all 18 blocks, with the signed-difference driver of the W_2^2 path preference ($\Delta_{\text{mean}}, \Delta_{\text{cov}}$).

d	solver	NFE	inversion	W_2 prefers	W_2^2 driver
2	euler	8	no	VP	covariance
2	euler	16	no	VP	covariance
2	euler	32	no	VP	covariance
2	heun	8	yes	linear	mean
2	heun	16	yes	linear	mean
2	heun	32	yes	linear	covariance
2	rk4	8	no	VP	covariance
2	rk4	16	no	VP	covariance
2	rk4	32	yes	linear	covariance
8	euler	8	yes	linear	mean
8	euler	16	yes	linear	mean
8	euler	32	yes	linear	mean
8	heun	8	yes	linear	mean
8	heun	16	yes	linear	mean
8	heun	32	yes	linear	mean
8	rk4	8	no	VP	covariance
8	rk4	16	no	VP	covariance
8	rk4	32	yes	linear	mean

Table 7: Signed path differences for the non-centered family. $\Delta_{\text{mean}} = M_{\text{lin}} - M_{\text{VP}}$, $\Delta_{\text{cov}} = C_{\text{lin}} - C_{\text{VP}}$, $\Delta_{\text{total}} = \Delta_{\text{mean}} + \Delta_{\text{cov}}$.

d	solver	NFE	inv.	Δ_{mean}	Δ_{cov}	Δ_{total}
2	euler	8	no	-1.969780×10^{-2}	7.008429×10^{-2}	5.038649×10^{-2}
2	euler	16	no	-5.010761×10^{-3}	1.895811×10^{-2}	1.394734×10^{-2}
2	euler	32	no	-1.266182×10^{-3}	4.941067×10^{-3}	3.674885×10^{-3}
2	heun	8	yes	-1.401380×10^{-3}	-8.296535×10^{-4}	-2.231034×10^{-3}
2	heun	16	yes	-8.132945×10^{-5}	-7.740260×10^{-5}	-1.587321×10^{-4}
2	heun	32	yes	-4.800285×10^{-6}	-5.428459×10^{-6}	-1.022874×10^{-5}
2	rk4	8	no	-1.931000×10^{-6}	7.169013×10^{-5}	6.975913×10^{-5}
2	rk4	16	no	$-5.235990 \times 10^{-10}$	1.106023×10^{-8}	1.053664×10^{-8}
2	rk4	32	yes	$-7.527744 \times 10^{-12}$	$-5.405709 \times 10^{-11}$	$-6.158484 \times 10^{-11}$
8	euler	8	yes	-3.252232×10^{-1}	2.207633×10^{-1}	-1.044599×10^{-1}
8	euler	16	yes	-8.388778×10^{-2}	5.947257×10^{-2}	-2.441521×10^{-2}
8	euler	32	yes	-2.129903×10^{-2}	1.546831×10^{-2}	-5.830720×10^{-3}
8	heun	8	yes	-5.927520×10^{-3}	-4.969512×10^{-4}	-6.424471×10^{-3}
8	heun	16	yes	-3.450747×10^{-4}	-9.787629×10^{-5}	-4.429510×10^{-4}
8	heun	32	yes	-2.074391×10^{-5}	-7.576731×10^{-6}	-2.832065×10^{-5}
8	rk4	8	no	-3.909251×10^{-6}	3.088406×10^{-4}	3.049314×10^{-4}
8	rk4	16	no	-1.137695×10^{-8}	7.433020×10^{-8}	6.295325×10^{-8}
8	rk4	32	yes	$-7.287617 \times 10^{-11}$	2.093892×10^{-11}	$-5.193726 \times 10^{-11}$

changes every cell by less than 1%.

G Robustness grids

Table 8 records aligned and perturbed evaluations of Theorem 1 at $L = \log 2$. Ranking inversion is recorded only when exact endpoints still match and the constant minimizer has strictly larger solver error. Phase offsets and interior nodes preserve endpoints but destroy Euler exactness. Frequencies outside $(\frac{1}{2}\mathbb{Z})$ break the common-endpoint constraint; the tested half-integer frequency ($f = 4.5$) preserves the endpoints while destroying Euler

exactness. Heun is not exact on the aligned oscillation. These checks show how quickly the aliasing phenomenon disappears when grid alignment is perturbed; they are not a population frequency.

Table 8: Grid-aware robustness at $L = \log 2$. “Ends” records whether the two exact endpoints still match. “Invert” records whether the constant $\mathcal{R}$ -minimizer has strictly larger solver error.

N	solver	ϕ	f	θ	err a_0	err $a_{1,N}$	ends	invert
4	euler	0	4	0	0.105	$< 10^{-12}$	yes	yes
8	euler	0	8	0	0.056	$< 10^{-12}$	yes	yes
16	euler	0	16	0	0.029	$< 10^{-12}$	yes	yes
4	euler	$\pi/2$	4	0	0.105	0.105	yes	no
4	euler	π	4	0	0.105	0.206	yes	no
8	euler	$\pi/4$	8	0	0.056	0.017	yes	yes
4	euler	0	4.5	0	0.105	0.079	yes	yes
8	euler	0	7	0	0.056	0.056	yes	no
8	euler	0	9	0	0.056	0.056	yes	no
8	euler	0	8	0.5	0.056	0.111	yes	no
4	heun	0	4	0	6.09e-03	0.123	yes	no
8	heun	0	8	0	1.63e-03	0.061	yes	no

The post-hoc factor-plus-noise grid has 66 preference blocks: 2 dimensions, 3 solvers, and, for each pair, the three primary NFE budgets $\{8, 16, 32\}$ at noise-floor perturbations $\delta \in \{-0.1, 0, 0.1\}$, plus NFE $\{64, 128\}$ at $\delta = 0$ only. The perturbation multiplies the noise variance 0.05 by $(1 + \delta)$; the factor F is regenerated from seed 271828 and is not redrawn. Extra budgets are not run at $\delta \neq 0$. Table 10 summarizes the 66 blocks. All 66 follow the same solver-path split as the primary factor-plus-noise census: VP under Euler, linear under Heun and RK4.

The geometry-draw experiment of Section 6 is the one place F is redrawn. The paired concordance score on the centered enumeration is discussed in Appendix F and recorded in Table 12.

H Reproducibility, artifacts, and provenance

Code, compact artifacts, frozen inputs, and reproducibility instructions are available at <https://github.com/clement-callaert/fewstep-field-regularity>. Default numerical precision is float64. The scalar calculation uses exact rational and algebraic expressions together with validated bounds; the precision audit uses 80-digit arithmetic. Compact artifacts live under `paper/arxiv/artifacts/` and are committed files. Frozen Phase 4 Hydra tables used by the eigenmode and non-centered figures live under `paper/arxiv/frozen_runs/`. Eigenvalues, exact-moment modal Runge–Kutta, and those compact files suffice for Table 3 and Proposition 2. The raw Hydra directory `outputs/` is not required to read the PDF. The 80-digit Gelbrich audit is the frozen precision table in `paper/arxiv/frozen_runs/`.

Table 9: Continuous $\mathcal{R}$ versus $\widehat{\mathcal{R}}_{24}$ on headline geometries, including the non-centered anisotropic family. Quadrature, not an exact integral except on the scalar pair.

geometry	$\mathcal{R}_{\text{lin}}$	$\widehat{\mathcal{R}}_{24}^{\text{lin}}$	$\mathcal{R}_{\text{VP}}$	$\widehat{\mathcal{R}}_{24}^{\text{VP}}$
scalar $\lambda = 4$	0.963495	0.964206	0.616850	0.616850
anisotropic $d = 2, 8$	0.972296	0.972637	0.192270	0.192361
factor+noise $d = 2$	1.245531	1.244898	0.625461	0.625454
rank-2 factor+noise $d = 8$	2.944104	2.947652	4.730544	4.729521
non-centered anisotropic $d = 2, 8$	1.093537	1.093851	0.342354	0.342536

Table 10: Post-hoc factor-plus-noise robustness: all 66 blocks follow the same solver-path split as the primary grid. The factor F is not redrawn.

solver	$d = 2$ blocks	$d = 8$ blocks	preferred path
euler	11	11	VP
heun	11	11	linear
rk4	11	11	linear

Table 11: Lebesgue measure in $\log \lambda$ of the linear-versus-VP inversion region on $[0.05, 100]$. Resolution 1281 nodes; doubling changes every entry by less than 1%. Not a count of inverted grid points.

solver	NFE	meas $_{\log \lambda}$	fraction of $[\ln 0.05, \ln 100]$
euler	8	2.444	0.321
euler	16	2.556	0.336
euler	32	2.669	0.351
heun	8	3.337	0.439
heun	16	3.723	0.490
heun	32	3.985	0.524
rk4	8	2.212	0.291
rk4	16	3.082	0.405
rk4	32	3.432	0.452

Table 12: Paired concordance score $(n_{\text{agree}} - n_{\text{invert}})/n$ between the $\mathcal{R}$ order and the W_2 order on the linear-versus-VP census. Geometry strata use $n = 9$ blocks (3 solvers $\times$ 3 budgets); solver strata use $n = 12$ blocks (4 geometries $\times$ 3 budgets). Descriptive statistic on a complete enumeration, not Kendall's τ and not an inferential test.

stratum	paired concordance
tested 36-block grid	2/9
anisotropic $d = 2$	1/3
anisotropic $d = 8$	5/9
factor+noise $d = 2$	-1/3
rank-2 factor+noise $d = 8$	1/3
Euler	1/2
Heun	-1/3
RK4	1/2

Source: <https://github.com/clement-callaert/fewstep-field-regularity>.