

When Averaged Field Regularity Fails to Rank Few-Step Generative Paths

Clément Callaert
CentraleSupélec and Université Paris-Saclay
callaert.clement@gmail.com

Abstract

Choosing a probability path for few-step generative sampling is often guided by an averaged regularity scalar of the velocity field: a path with smaller averaged squared Jacobian norm is expected to discretize better at a fixed budget of function evaluations (NFE). This paper studies that ranking question in commuting Gaussian probability-flow ODEs, where endpoint laws and Gaussian 2-Wasserstein (W_2) errors are available in closed form, the regularity integrand is analytically known, and sampling, training, and estimation noise are absent. On a registered grid of 72 equal-NFE configurations (36 two-path comparisons) the averaged-regularity ordering of linear versus variance-preserving paths does not determine the fixed-NFE W_2 ordering: 14 of 36 blocks invert. The same phenomenon appears in a separately specified non-centered commuting family (11 of 18 blocks). All reported W_2 values and inversion orderings survive an 80-digit reference calculation. An exact modal decomposition writes endpoint factor error as a transported sum of signed, solver-stage-dependent local defects, information that an unsigned time average of Jacobian norms cannot retain. An explicit scalar construction proves the corresponding non-implication for fixed-grid left-endpoint Euler, for every $L > 0$ and every integer grid size $N \geq 1$. The construction is grid-aware and is not offered as the mechanism of the Gaussian inversions. The study is a controlled limitation: it does not refute regularity-guided schedule design, and no learned model is evaluated. The counts 14 of 36 and 11 of 18 are descriptive benchmark frequencies, not estimates of a population probability.

1 Introduction

Few-step sampling of a continuous-time generative model commits to a probability path, a numerical solver, and a small NFE budget before any endpoint error is observed. Classical one-step error bounds involve Lipschitz and stability constants of the vector field [Hairer et al., 1993]. In the stochastic interpolant framework, Chen et al. [2025] propose minimizing averaged squared drift Lipschitzness as a schedule criterion (their Definition 3.2). Bounding an error, however, is weaker than determining which of two paths errs less at an equal small budget.

This paper isolates that ranking question where it can be answered without estimators. The systems are commuting Gaussian interpolants. The marginal velocity is affine. A fixed-step Runge–Kutta method induces an affine endpoint map, reconstructed from $d + 1$ deterministic probes. Endpoint error is the closed-form Gaussian W_2 [Gelbrich, 1990, Peyré and Cuturi, 2019]. The regularity scalar is a trapezoidal quadrature of an analytically known spectral-norm integrand. Two paths are compared: the linear interpolant and the trigonometric variance-preserving (VP) interpolant.

Contributions.

- A controlled closed-form W_2 benchmark on commuting Gaussian flows: 14 inversions among 36 equal-NFE comparisons in a registered grid of 72 configurations, all 80-digit-verified, and a separately specified non-centered commuting family with 11 inversions among 18 comparisons.




Figure 1: Conceptual schematic, not experimental data. (a) Averaging collapses two time-dependent fields to two scalars, while a solver samples only stage locations (markers). (b) Local defects are signed: same-sign defects accumulate, alternating defects cancel. (c) After transport to the endpoint, the error ranking can oppose the averaged ranking. The panel illustrates the accounting of Section 5, not the measured mechanism of any specific Gaussian inversion.

- An exact eigenmode analysis: the endpoint factor error is a transported sum of signed, solver-stage-dependent local defects.
- An explicit grid-aware smooth scalar construction proving that, for every $L > 0$ and every integer $N \geq 1$, strictly larger averaged squared Jacobian is compatible with zero fixed-grid Euler endpoint error.

Solver dependence of good schedules is already documented in diffusion practice [Sabour et al., 2024, Karras et al., 2022]. The contribution here is the exact, controlled non-implication and its modal accounting: what the tested averaged-regularity scalar certifies, and what it discards, about fixed-budget *rankings* in these systems. Contemporaneous analyses of strain versus vorticity and of material-derivative regularization [Tao and Choi, 2026, Khan, 2026] address related numerical structure. They do not study the linear-versus-VP ranking by averaged spectral Jacobian on independent-coupling interpolants.

2 Background and problem formulation

Definition (interpolant). Let $X_0 \sim N(\mu_0, \Sigma_0)$ and $X_1 \sim N(\mu_1, \Sigma_1)$ be independent in $\mathbb{R}^d$, with Σ_0, Σ_1 symmetric positive definite. Let $\alpha, \sigma : [0, 1] \rightarrow \mathbb{R}$ be C^3 scalar schedules. The interpolant is

$$X_t = \alpha(t)X_0 + \sigma(t)X_1, \quad (1)$$

as in stochastic interpolants and flow matching [Albergo et al., 2025, Lipman et al., 2023, Liu et al., 2023, Tong et al., 2024]. The two schedules used as *paths* in the benchmark are the linear path $(\alpha, \sigma) = (1 - t, t)$ and the trigonometric VP path $(\cos(\pi t/2), \sin(\pi t/2))$, the interpolant analogue of VP diffusion schedules [Song et al., 2021].

Problem. Fix a solver from {Euler, Heun, RK4}, an NFE budget B divisible by the solver’s stage count, and two paths b and b' . Let $\mathcal{R}$ be the averaged regularity scalar of Section 4 and let E be the Gaussian W_2 endpoint error of Section 4. Does $\mathcal{R}[b] < \mathcal{R}[b']$ imply $E[b] < E[b']$?

The paper answers this in the negative for the tested commuting Gaussian systems, as a logical and empirical non-implication, not as a claim about learned neural fields.

3 Gaussian probability paths and marginal velocity fields

Assumption (independent Gaussian endpoints and positive definite marginals). The endpoints are independent Gaussians as above, and $Q(t) := \text{Cov}(X_t) = \alpha(t)^2 \Sigma_0 + \sigma(t)^2 \Sigma_1$ is positive definite for every $t \in [0, 1]$.

Under this assumption the marginal law of X_t is Gaussian with mean $m_t = \alpha\mu_0 + \sigma\mu_1$. Differentiating the interpolant and taking the Gaussian conditional expectation yields the affine marginal velocity

$$b(t, x) = A(t)x + c(t), \quad A(t) = C_t Q(t)^{-1}, \quad c(t) = \dot{m}_t - A(t)m_t, \quad (2)$$

with $C_t = \dot{\alpha}\alpha \Sigma_0 + \dot{\sigma}\sigma \Sigma_1$. This is the standard Gaussian conditional formula, written in the notation of Chen et al. [2025, Example 3.3] and extended to non-centered non-isotropic endpoints. Invertibility of $Q(t)$ is required at every solver stage.

Centered specialization. If $\mu_0 = \mu_1 = 0$ and $\Sigma_0 = I$, then $c \equiv 0$ and $A(t) = \frac{1}{2}Q'(t)Q(t)^{-1}$.

Assumption (commuting covariance flow). Throughout the benchmark, $\Sigma_0 = I$ and Σ_1 is simultaneously diagonalized with I . Then $Q(t)$, $Q'(t)$, and $A(t)$ commute. In the eigenbasis of Σ_1 , mode i with eigenvalue $\lambda_i > 0$ obeys $x'_i = a_i(t)x_i + c_i(t)$ with $q_i(t) = \alpha(t)^2 + \lambda_i\sigma(t)^2$ and $a_i(t) = q'_i(t)/(2q_i(t))$ when $\Sigma_0 = I$. The exact one-step factor on $[t_n, t_{n+1}]$ is $\sqrt{q_i(t_{n+1})/q_i(t_n)}$. For centered modes, $q_i(0) = 1$ and $q_i(1) = \lambda_i$, so the exact endpoint factor is $\sqrt{\lambda_i}$.

This commuting hypothesis is essential. Noncommuting covariance flows are outside the theorems and the tables.

4 Averaged field regularity and fixed-budget error

Definition (averaged regularity scalar). Following Chen et al. [2025, Definition 3.2],

$$\mathcal{R}[b] = \int_0^1 \|A(t)\|_2^2 dt \quad (3)$$

for the affine field (2), where $\|\cdot\|_2$ is the matrix spectral norm. For state-independent $A(t)$ this coincides with the path expectation in their Eq. (3.4). By definition $\mathcal{R}$ ignores $c(t)$.

Numerical evaluation (not an exact integral). In every reported number, $\mathcal{R}$ is a trapezoidal quadrature of the analytically known integrand on 24 uniform nodes in $[0, 1]$. The implementation records this quantity as non-exact. Rankings in the registered grid are nevertheless strict: no comparison block has a tied $\mathcal{R}$ value.

Definition (equal-NFE accounting). Euler uses 1 field evaluation per step, Heun 2, and classical RK4 4. A requested budget B is admitted only if B is divisible by the stage count s . The step count is $S = B/s$. The time grid is the uniform partition of $[0, 1]$ with S steps, including both endpoints. Jacobian evaluations used for metrics are not counted as NFE. Equal NFE is not equal step count: at $B = 8$, Euler takes 8 steps, Heun 4, and RK4 2.

Definition (endpoint error). A solver applied to (2) produces an affine endpoint map $x \mapsto Rx + d$, recovered from the images of 0 and the canonical basis. Source moments are pushed through this map. Error is the Gaussian W_2 [Gelbrich, 1990, Peyré and Cuturi, 2019],

$$W_2^2 = \|m - \mu_1\|_2^2 + \text{tr}(C + \Sigma_1 - 2(\Sigma_1^{1/2}C\Sigma_1^{1/2})^{1/2}), \quad (4)$$

with PSD square roots. In the centered commuting case, if r_i is the numerical endpoint factor of mode i ,

$$W_2^2 = \sum_i (|r_i| - \sqrt{\lambda_i})^2. \quad (5)$$

The absolute value appears because the PSD square root of r_i^2 is $|r_i|$. This is endpoint *distribution* error, not trajectory error.

Reported float64 values are compared with an 80-digit `mpmath` propagation of the same modal maps. The comparison is a precision audit, not a second experiment.

5 Exact solver-stage and modal error analysis

Let $e_{i,n} = \sqrt{q_i(t_{n+1})/q_i(t_n)}$ be the exact factor of mode i and let $r_{i,n}$ be the numerical one-step factor (Appendix A). For nonzero factors the algebraic identity

$$\prod_{n=0}^{S-1} r_{i,n} - \prod_{n=0}^{S-1} e_{i,n} = \sum_{n=0}^{S-1} (r_{i,n} - e_{i,n}) \left(\prod_{j<n} r_{i,j} \right) \left(\prod_{j>n} e_{i,j} \right) \quad (6)$$

holds with no smallness or positivity assumption. Prefixes are numerical; suffixes are exact. This is the transported signed-defect decomposition.

For smooth a , the leading exact-minus-method coefficients are $\frac{1}{2}(a' + a^2)h^2$ for Euler and $(-\frac{1}{12}a'' + \frac{1}{6}a^3)h^3$ for the implemented Heun method (Appendix B). They involve different derivatives, signs, and stage locations. Classical RK4 cancels all lower-order terms; no closed RK4 coefficient is asserted.

Identity (6) identifies what $\mathcal{R}$ discards: which times the stages sample, the derivatives of A , the signs of local defects, their cancellation, and their transport to the endpoint. An unsigned average of $\|A\|_2^2$ therefore need not determine the fixed-NFE ordering. This is an accounting statement. It is not, by itself, a proof that any particular Gaussian pair inverts.

6 Controlled benchmark

Registered grid. Source $N(0, I)$. Targets: anisotropic Gaussians with condition number 4, and seeded low-rank Gaussians $FF^\top + 0.05I$ with rank 2 and geometry seed 271828. Dimensions $d \in \{2, 8\}$. Paths: linear and VP. Solvers: Euler, Heun, RK4. NFE $\in \{8, 16, 32\}$ with equal-NFE accounting. This is 72 float64 CPU configurations in 36 two-path blocks. Mixture targets are excluded: dimension-8 mixture endpoint estimators failed a post-result calibration diagnostic, and Phase 4 reproduction accepts Gaussian targets only.

Inversion. A block is an inversion when the strict $\mathcal{R}$ ordering disagrees with the strict W_2 ordering. There are no ties in this grid.

Chronology. Inversions were first identified in a registered broader gate benchmark, then the Gaussian grid above was frozen and rerun from clean code. The decomposition, local-error proxies, extra NFE budgets, and perturbations are post-hoc relative to that gate. A non-centered family was specified in writing before its result commit (Section 7).

7 Main empirical results

All 72 rows reproduce the registered precursor within absolute tolerance 10^{-12} . Independent grouping of the frozen `results.json` yields 14 inversions among 36 blocks (Figure 2; full list in Table 2 of the appendix). These are descriptive counts on the tested grid.

Table 1: Strongest inversion in the registered grid: low-rank Gaussian, $d = 8$, Euler, NFE 8. Values are generated from the frozen artifact; $\mathcal{R}$ is trapezoidal with 24 nodes. Lower $\mathcal{R}$ is the regularity preference; lower W_2 is the error preference. The table supports a ranking disagreement on this block. It does not support a universal preference for VP.

path	$\mathcal{R}[b]$	Gaussian W_2
linear	2.9476523251	0.8108540111
variance-preserving	4.7295206355	0.4564779075




Figure 2: Left and middle: strongest inversion in both quantities; hatching marks VP. Right: W_2 margins of all 14 inversion blocks on a logarithmic axis (which compresses, never exaggerates, magnitudes). Marker shape and color give the W_2 -preferred path. Source: frozen Phase 4 reproduction, 36 blocks.

In the strongest inversion (low-rank, $d = 8$, Euler, NFE 8; Table 1), $\mathcal{R}$ prefers linear while W_2 prefers VP, with margin 0.3543761036 . All 72 float64 W_2 values agree with the 80-digit reference to at most $9.7050430488 \times 10^{-10}$. The smallest inversion margin, $1.1188920612 \times 10^{-5}$ (low-rank, $d = 2$, RK4, NFE 32), is 11528 times that global precision gap. Every inverted block keeps the same W_2 ordering at 80 digits.

Solver–path interaction (low-rank family only). On all 18 low-rank primary blocks, VP has smaller W_2 under Euler and linear has smaller W_2 under Heun and RK4, at both dimensions and every primary budget (Figure 3). The pattern persists on a post-hoc robustness grid with NFE 64 and 128 and with $\pm 10\%$ target perturbations (66 of 66 low-rank robustness blocks). Solver-dependent schedule preference is known [Sabour et al., 2024]. Here the tested averaged-regularity scalar does not capture the interaction. The split is not claimed for the anisotropic family, whose primary inversions are Heun-only.

Non-centered supporting family. Before execution, one additional family was frozen: source $N(\mu_0, I)$ with $\mu_{0,i} = 0.75(-1)^i$; target $N(\mu_1, D)$ with $\mu_{1,i} = 1 + 0.25i$ and D diagonal, geometric eigenvalues, condition number 6. Then $c(t) \neq 0$ in (2). The same paths, solvers, and budgets give 18 blocks. A documented amendment between the freeze and the result commit changed only the endpoint sanity check (moment gap instead of a W_2 threshold that float64 identical Gaussians cannot meet). The family itself was not changed.

11 of 18 blocks invert. $\mathcal{R}$, which ignores $c(t)$, prefers VP with margin 0.7513; W_2 prefers




Figure 3: Signed low-rank W_2 margin at equal NFE. Positive values prefer VP; negative values prefer linear. The axis is symmetric-log (linear threshold 10^{-4}) because margins span several orders of magnitude and change sign. The figure supports a solver-dependent preference on this family and these budgets. It does not establish a schedule rule outside the tested systems.




Figure 4: Exact modal W_2^2 contributions and transported signed local defects on the dominant mode for the strongest comparison. The decomposition is an identity on the recorded factors. Aggregate solver proxies are not shown as predictive.

linear on the inverted blocks. All inverted blocks pass the 80-digit audit (maximum float64 deviation 2.07×10^{-11} ; smallest inverted margin 1.42×10^{-6}). The family remains Gaussian, diagonal, and commuting. It is one supporting replication, not out-of-distribution evidence.

Post-hoc local-error proxy. On the same 36 centered blocks, $\mathcal{R}$ agrees with the observed W_2 preference in 22 blocks. A solver-specific leading local-error proxy agrees in 29 blocks. That comparison is post-hoc and in-sample. It is not a predictive claim. Exact transported defects account for endpoint factor error by (6).

8 Fixed-grid Euler non-implication

Proposition 1. Fix $L > 0$ and an integer $N \geq 1$. Let $h = 1/N$ and let forward Euler use the left-endpoint grid $t_n = nh$ on $[0, 1]$. Set $\varepsilon_N = N(e^{L/N} - 1) - L > 0$ and define $f_k(t, x) = a_k(t)x$ with $a_0(t) = L$ and $a_{1,N}(t) = L + \varepsilon_N \cos(2\pi Nt)$. For $x(0) = 1$: (i) both exact endpoints equal e^L ; (ii) $\int_0^1 a_{1,N}^2 = L^2 + \varepsilon_N^2/2 > \int_0^1 a_0^2$; (iii) Euler is exact for $a_{1,N}$; (iv) Euler has strictly positive

endpoint error for a_0 . Hence larger averaged squared Jacobian does not imply larger fixed-grid Euler endpoint error.

Proof. (i) $\int_0^1 \cos(2\pi Nt) dt = 0$ for integer N , so both coefficients integrate to L and $x(1) = e^L$.
(ii) Orthogonality gives the displayed integrals, and $\varepsilon_N > 0$ because $e^z > 1 + z$ for $z = L/N > 0$.
(iii) $\cos(2\pi Nt_n) = 1$ at every left node, so each Euler factor equals $e^{L/N}$ and the product is e^L .
(iv) $\log(1 + z) < z$ for $z > 0$ yields $(1 + L/N)^N < e^L$. $\square$

The quantifiers matter. The field $a_{1,N}$ is chosen after the grid. Its amplitude vanishes as $N \rightarrow \infty$. Shifted grids and multi-stage solvers sample different phases. The construction shows non-implication for fixed-grid left-endpoint Euler only. It is not the mechanism behind every Gaussian inversion, and it is not presented as a new numerical-analysis phenomenon. Edge cases, including $N = 1$ and the exclusion of $L = 0$, are recorded in Appendix C. An 80-digit check confirms that the oscillatory Euler product equals e^L .

9 Relation to existing schedule and solver design

Flow matching and stochastic interpolants supply the Gaussian path family and its fields [Lipman et al., 2023, Albergo et al., 2025, Liu et al., 2023, Tong et al., 2024]. Lipschitz-guided schedule design proposes the criterion tested here [Chen et al., 2025]. Fast diffusion solvers exploit exactly integrated linear parts and solver-specific expansions [Lu et al., 2022a,b, Zhang and Chen, 2023]. EDM separates schedule from discretization [Karras et al., 2022], and Align Your Steps tunes schedules per solver, model, and dataset [Sabour et al., 2024]. Runge–Kutta and backward error analysis supply the local-error machinery [Butcher, 1963, Hairer et al., 1993, 2006].

Tao and Choi [2026] decompose Jacobian strain and vorticity in flow-matching integration error and prove that McCann displacement interpolation has vanishing material derivative, so Euler is exact on those Lagrangian trajectories. The paths in the present benchmark are independent-coupling interpolants, not McCann maps, so that exactness result does not apply to the linear and VP fields used here. Khan [2026] regularize the material derivative of learned fields. Neither paper produces the commuting inversion table or Proposition 1.

A literature search on 2026-08-13 (arXiv and the sources listed in the accompanying audit) did not locate an identical commuting Gaussian ranking counterexample. Search absence is not a proof that none exists.

10 Limitations and scope

The systems are closed-form commuting affine Gaussian flows with two paths, three fixed-step solvers, $d \in \{2, 8\}$, small equal-NFE budgets, and one non-centered commuting replication. No learned neural field is evaluated. No image, video, or world-model quantity is reported. Mixture targets were excluded after estimator calibration failure; they do not support any conclusion here. Proposition 1 is grid-aware and Euler-only. Exponential and exact-linear integrators solve affine scalar modes exactly, so the numerical comparisons concern generic fixed-stage Runge–Kutta discretization. Robustness NFE 64 and 128 and $\pm 10\%$ perturbations are post-hoc. The leading local-error proxy is post-hoc and in-sample.

The result does not refute Lipschitz-guided schedules. In the tested systems the averaged-regularity *ordering* does not determine the fixed-NFE W_2 ordering, because the tested scalar does not retain solver-stage sampling, signs, cancellation, or transport.

11 Conclusion

On a registered commuting Gaussian grid, ordering two interpolants by trapezoidal averaged squared Jacobian does not determine their equal-NFE Gaussian W_2 ordering. The mismatch

is visible in 14 of 36 blocks and in 11 of 18 blocks of a specified non-centered family, and it is compatible with an exact transported-defect identity and with a grid-aware Euler non-implication. The statement is as narrow as the evidence.

12 Reproducibility statement

Default analytical precision is float64. Figures and numerical macros in this PDF are generated by `scripts/make_arxiv_figures.py` and `scripts/make_arxiv_tables.py` from checksum-pinned JSON artifacts. The primary artifact `phase4_gaussian_reproduction_2026-07-24-v1:results` has SHA-256 `b8930142cba5655ee553aae5ff400cd884c1137e77547d9a5fa94bd4e354973f`. Validate a run directory with `pythonscripts/validate_artifacts.py` outputs/`<run_id>`. Tests: `pytest`. Release-ready runs refuse a dirty worktree. Code commit for this manuscript audit: `e48c9390e62b38f206342e6aeb7f160122ccc79c`. Repository: <https://github.com/clement-callaert/fewstep-field-regularity>.

References

- Michael S. Albergo, Nicholas M. Boffi, and Eric Vanden-Eijnden. Stochastic interpolants: A unifying framework for flows and diffusions. *Journal of Machine Learning Research*, 26(209): 1–80, 2025. arXiv:2303.08797.
- John C. Butcher. Coefficients for the study of Runge-Kutta integration processes. *Journal of the Australian Mathematical Society*, 3(2):185–201, 1963.
- Yifan Chen, Eric Vanden-Eijnden, and Jiawei Xu. Lipschitz-guided design of interpolation schedules in generative models. *arXiv preprint arXiv:2509.01629*, 2025. version 3, 16 May 2026.
- Matthias Gelbrich. On a formula for the L^2 Wasserstein metric between measures on Euclidean and Hilbert spaces. *Mathematische Nachrichten*, 147:185–203, 1990. doi: 10.1002/mana.19901470121.
- Ernst Hairer, Syvert P. Nørsett, and Gerhard Wanner. *Solving Ordinary Differential Equations I: Nonstiff Problems*. Springer, 2nd edition, 1993.
- Ernst Hairer, Christian Lubich, and Gerhard Wanner. *Geometric Numerical Integration: Structure-Preserving Algorithms for Ordinary Differential Equations*. Springer, 2nd edition, 2006.
- Tero Karras, Miika Aittala, Timo Aila, and Samuli Laine. Elucidating the design space of diffusion-based generative models. In *Advances in Neural Information Processing Systems*, 2022. arXiv:2206.00364.
- Tauhid Khan. Isokinetic flow matching for pathwise straightening of generative flows. *arXiv preprint arXiv:2604.04491*, 2026.
- Yaron Lipman, Ricky T. Q. Chen, Heli Ben-Hamu, Maximilian Nickel, and Matthew Le. Flow matching for generative modeling. In *International Conference on Learning Representations (ICLR)*, 2023. arXiv:2210.02747.
- Xingchao Liu, Chengyue Gong, and Qiang Liu. Flow straight and fast: Learning to generate and transfer data with rectified flow. In *International Conference on Learning Representations (ICLR)*, 2023. arXiv:2209.03003.

- Cheng Lu, Yuhao Zhou, Fan Bao, Jianfei Chen, Chongxuan Li, and Jun Zhu. DPM-solver: A fast ODE solver for diffusion probabilistic model sampling in around 10 steps. In *Advances in Neural Information Processing Systems*, 2022a. arXiv:2206.00927.
- Cheng Lu, Yuhao Zhou, Fan Bao, Jianfei Chen, Chongxuan Li, and Jun Zhu. DPM-solver++: Fast solver for guided sampling of diffusion probabilistic models. *arXiv preprint arXiv:2211.01095*, 2022b.
- Gabriel Peyré and Marco Cuturi. *Computational Optimal Transport*. Now Publishers, 2019. arXiv:1803.00567.
- Amirmojtaba Sabour, Sanja Fidler, and Karsten Kreis. Align your steps: Optimizing sampling schedules in diffusion models. In *International Conference on Machine Learning (ICML)*, 2024. arXiv:2404.14507.
- Yang Song, Jascha Sohl-Dickstein, Diederik P. Kingma, Abhishek Kumar, Stefano Ermon, and Ben Poole. Score-based generative modeling through stochastic differential equations. In *International Conference on Learning Representations (ICLR)*, 2021. arXiv:2011.13456.
- Chenxi Tao and Seung-Kyum Choi. On the role of strain and vorticity in numerical integration error for flow matching. *arXiv preprint arXiv:2605.06680*, 2026.
- Alexander Tong, Kilian Fatras, Nikolay Malkin, Guillaume Hugué, Yanlei Zhang, Jarrod Rector-Brooks, Guy Wolf, and Yoshua Bengio. Improving and generalizing flow-based generative models with minibatch optimal transport. *Transactions on Machine Learning Research*, 2024. arXiv:2302.00482.
- Qinsheng Zhang and Yongxin Chen. Fast sampling of diffusion models with exponential integrator. In *International Conference on Learning Representations (ICLR)*, 2023. arXiv:2204.13902.

A Solver formulas and equal-NFE accounting

On a step $[t, t + h]$ for $x' = a(t)x$, the implemented factors are

$$r_{\text{Euler}} = 1 + ha(t), \quad (7)$$

$$r_{\text{Heun}} = 1 + \frac{h}{2}(a(t) + a(t+h)(1 + ha(t))), \quad (8)$$

and classical RK4 with stages at t , $t + h/2$ (twice), and $t + h$. These match the vector solvers applied to affine fields. Equal-NFE requires $B \equiv 0 \pmod{s}$ and sets $S = B/s$.

B Local truncation expansions

Assume a is C^3 . The exact factor admits the Taylor expansion $1 + ha + (h^2/2)(a' + a^2) + (h^3/6)(a'' + 3aa' + a^3) + O(h^4)$. Subtracting the Euler factor leaves $\frac{1}{2}(a' + a^2)h^2 + O(h^3)$. Substituting the Taylor expansion of $a(t+h)$ into the implemented Heun factor and subtracting from the exact series leaves $(-a''/12 + a^3/6)h^3 + O(h^4)$. The aa' terms cancel. This expansion was confirmed with independent symbolic algebra on the implemented expression. RK4 is compared numerically to the exact factor; no memorized non-autonomous RK4 coefficient is used.

C Proof details for Proposition 1

The body proof is complete for the stated quantifiers. Additional remarks: $N = 1$ holds, with $\varepsilon_1 = e^L - 1 - L$ and a single Euler factor e^L . The case $L = 0$ is excluded because the metric ordering is not strict. As $N \rightarrow \infty$ with L fixed, $\varepsilon_N = L^2/(2N) + O(N^{-2})$, so the oscillatory amplitude vanishes. Every Euler factor in the construction is positive. An 80-digit evaluation of $(1 + h(L + \varepsilon_N))^N$ equals e^L to working precision for representative pairs including $(L, N) = (1, 8)$. The construction does not apply to Heun, RK4, right-endpoint Euler, or a fixed oscillatory ODE under grid refinement.

D Full inversion tables

Table 2 lists every inverted block in the registered centered grid. Table 3 lists every comparison block in the non-centered family. Both tables are generated from pinned artifacts.

Table 2: All 14 inversions among 36 centered blocks. Margins are absolute Gaussian W_2 differences. Descriptive grid counts only.

family	d	solver	NFE	R prefers	W_2 prefers	W_2 margin
anisotropic	2	heun	8	VP	linear	1.905310×10^{-3}
anisotropic	2	heun	16	VP	linear	1.939806×10^{-3}
anisotropic	2	heun	32	VP	linear	6.634439×10^{-4}
anisotropic	8	heun	16	VP	linear	1.758942×10^{-3}
anisotropic	8	heun	32	VP	linear	7.628173×10^{-4}
low-rank	2	heun	8	VP	linear	1.836251×10^{-2}
low-rank	2	heun	16	VP	linear	6.039925×10^{-3}
low-rank	2	heun	32	VP	linear	1.698035×10^{-3}
low-rank	2	rk4	8	VP	linear	1.516722×10^{-3}
low-rank	2	rk4	16	VP	linear	1.289981×10^{-4}
low-rank	2	rk4	32	VP	linear	1.118892×10^{-5}
low-rank	8	euler	8	linear	VP	3.543761×10^{-1}
low-rank	8	euler	16	linear	VP	2.046946×10^{-1}
low-rank	8	euler	32	linear	VP	1.096576×10^{-1}

E Precision-audit method

Each modal factor is recomputed in `mpmath` at 80 decimal digits with the same stage formulas. Gaussian W_2 is formed from those factors. The recorded maximum absolute float64-versus-reference difference on the centered grid is $9.7050430488 \times 10^{-10}$, below the pre-set tolerance 2×10^{-9} . A distinct residual, $9.7050436884 \times 10^{-10}$, measures eigenmode reconstruction of float64 W_2 and is not the 80-digit gap.

F Excluded mixture experiments

Phase 2 and Phase 3 contain Gaussian-mixture runners and sliced-Wasserstein estimators. Dimension-8 mixture blocks failed a post-result calibration diagnostic and were excluded from decision evidence. Dimension-2 two-mode blocks were used only in a gate condition and are not part of the Phase 4 conclusion. Phase 4 reproduction refuses non-Gaussian targets.

Table 3: Non-centered family: all 18 blocks. Positive $R_{\text{lin}} - R_{\text{VP}}$ means $\mathcal{R}$ prefers VP. Negative W_2 difference means W_2 prefers linear.

d	solver	NFE	inversion	$R_{\text{lin}} - R_{\text{VP}}$	$W_2^{\text{lin}} - W_2^{\text{VP}}$
2	euler	8	no	0.7513	1.225004×10^{-1}
2	euler	16	no	0.7513	6.613705×10^{-2}
2	euler	32	no	0.7513	3.436720×10^{-2}
2	heun	8	yes	0.7513	-2.054454×10^{-2}
2	heun	16	yes	0.7513	-6.514141×10^{-3}
2	heun	32	yes	0.7513	-1.793866×10^{-3}
2	rk4	8	no	0.7513	5.088161×10^{-3}
2	rk4	16	no	0.7513	1.869678×10^{-5}
2	rk4	32	yes	0.7513	-2.150047×10^{-6}
8	euler	8	yes	0.7513	-1.001525×10^{-1}
8	euler	16	yes	0.7513	-4.567154×10^{-2}
8	euler	32	yes	0.7513	-2.153309×10^{-2}
8	heun	8	yes	0.7513	-4.058830×10^{-2}
8	heun	16	yes	0.7513	-1.255211×10^{-2}
8	heun	32	yes	0.7513	-3.420523×10^{-3}
8	rk4	8	no	0.7513	1.303688×10^{-2}
8	rk4	16	no	0.7513	8.699785×10^{-5}
8	rk4	32	yes	0.7513	-1.423382×10^{-6}

G Artifact map and reproduction commands

alias	artifact
A1	phase4_gaussian_reproduction_2026-07-24-v1:results
A2	phase4_precision_2026-07-24-v1:table
A3	phase4_diagnostics_2026-07-24-v1:table
A4	phase4_robustness_2026-07-24-v1:table
A5–A7	external-validation results, inversions, precision

Install with Python 3.11+ and then:

```
python -m pip install -e ".[dev]"
pytest
python scripts/validate_artifacts.py \
    outputs/phase4_gaussian_reproduction_2026-07-24-v1
python scripts/make_arxiv_tables.py
python scripts/make_arxiv_figures.py
```

Release-ready experiment reruns require a clean worktree and `overwrite: false`.