

Averaged Jacobian Regularity Does Not Order Few-Step Error of Gaussian Interpolant ODEs

Clément Callaert

CentraleSupélec and Université Paris-Saclay
callaert.clement@gmail.com

Abstract

Few-step sampling of a continuous-time generative model requires a probability path, a numerical solver, and a budget of field evaluations (NFE). Chen et al. [2025] propose minimizing averaged squared Jacobian regularity A_2 as a path-selection criterion. This paper asks whether the order that this scalar induces between two interpolants is reliable at fixed NFE. The implication fails already for centered Gaussian interpolants.

The main logical result is a one-dimensional, seed-free counterexample. For $X_0 \sim N(0, 1)$ and $X_1 \sim N(0, 4)$ independent, the exact regularity integrals of the linear and trigonometric variance-preserving (VP) paths are $5\pi/8 - 1$ and $\pi^2/16$. Explicit Heun with NFE 8 (four steps of size $1/4$) reverses the ranking of Gaussian W_2 endpoint errors. The linear Heun factor is the rational $6797469/3559400$. The VP factor lies in $\mathbb{Q}[\pi, \sqrt{2}]$; substituting the documented bounds $\sqrt{2} < 99/70$ and $\pi < 355/113$ yields a purely rational certificate of $r_{\text{VP}} < 187/100$, hence $W_2^{\text{lin}} < 0.091 < 0.130 < W_2^{\text{VP}}$.

A commuting modal identity then accounts for what the unsigned average discards: solver-stage locations, signs, cancellation, and transport of local factor defects. On four centered geometries the per-mode log-covariance schedule (Chen et al. Example 3.3), which minimizes $\mathcal{R}$, attains the smallest W_2 in all 36 solver-budget blocks. Linear versus VP invert in 5 of 12 geometry×solver cells (4 of 12 at every NFE in $\{8, 16, 32\}$), from 3 distinct $\mathcal{R}$ comparisons. Thus $\mathcal{R}$ is reliable as an objective on this family and not reliable as a linear-versus-VP comparator away from that optimum.

1 Introduction

A continuous-time generative sampler is specified by four distinct objects: a probability path (an interpolant or noise schedule), the associated marginal velocity field, a numerical ODE solver, and an evaluation budget. Classical one-step bounds involve Lipschitz constants of the field [Hairer et al., 1993]. In the stochastic interpolant framework, Chen et al. [2025] propose minimizing averaged squared drift Lipschitzness (their Definition 3.2, denoted A_2) as a *selection* criterion. They prove no bound relating A_2 to a discretization error. We test whether the pairwise order that A_2 induces between two admissible paths is reliable at equal NFE.

This paper isolates that ranking question on commuting Gaussian interpolants, where the marginal field is affine, Gaussian W_2 is closed-form, and the regularity integrand is known exactly. The leading result is not a large grid. It is a deterministic scalar example, proved by an exact integral, an exact rational Heun product, and a rational enclosure of the VP product (Section 3). The subsequent modal analysis explains which information $\mathcal{R}$ discards (Section 4). The Cauchy–Schwarz minimizer of $\mathcal{R}$ and a λ -sweep then separate $\mathcal{R}$ as an objective from $\mathcal{R}$ as a linear-versus-VP comparator (Section 5). A hierarchical benchmark and a non-centered stress test show that linear-versus-VP inversions are not confined to the scalar pair (Sections 6–7).

Contemporaneous work studies Lipschitz bounds for Euler rates [Stéphanovitch, 2026], sharpness-aware grids [Gupta et al., 2026], least-action paths [Du et al., 2026], and instance-aware discretizations [Yuan et al., 2026]. Those results control other objects. Tao and Choi

[2026] give the general strain/vorticity form of the phenomenon; Section 8 records the precise relation.

2 Problem formulation and notation

Interpolant. Let $X_0 \sim N(\mu_0, \Sigma_0)$ and $X_1 \sim N(\mu_1, \Sigma_1)$ be independent in $\mathbb{R}^d$, with Σ_0, Σ_1 symmetric positive definite. Let $\alpha, \sigma : [0, 1] \rightarrow \mathbb{R}$ be C^3 and satisfy the boundary conditions

$$\alpha(0) = 1, \quad \sigma(0) = 0, \quad \alpha(1) = 0, \quad \sigma(1) = 1. \quad (1)$$

The interpolant is

$$X_t = \alpha(t)X_0 + \sigma(t)X_1, \quad (2)$$

as in stochastic interpolants and flow matching [Albergo et al., 2025, Lipman et al., 2023, Liu et al., 2023, Tong et al., 2024]. Throughout, $\alpha^2 + \sigma^2$ is not required to equal 1 except on the VP path. The two paths compared below are the linear schedule $(\alpha, \sigma) = (1 - t, t)$ and the trigonometric VP schedule $(\cos(\pi t/2), \sin(\pi t/2))$.

The law of X_t is Gaussian with mean $m_t = \alpha\mu_0 + \sigma\mu_1$ and covariance $Q(t) = \alpha^2\Sigma_0 + \sigma^2\Sigma_1$. We assume $Q(t) \succ 0$ on $[0, 1]$. The marginal velocity of (2) is the affine field

$$b(t, x) = A(t)x + c(t), \quad A(t) = C_t Q(t)^{-1}, \quad c(t) = \dot{m}_t - A(t)m_t, \quad (3)$$

with $C_t = \dot{\alpha}\Sigma_0 + \dot{\sigma}\Sigma_1$. This is a marginal interpolant ODE, not a score-based probability-flow ODE in the sense of Song et al. [2021], although the latter is an instance of a generative ODE. If $\mu_0 = \mu_1 = 0$ and $\Sigma_0 = I$, then $c \equiv 0$ and $A = \frac{1}{2}Q'Q^{-1}$.

Isotropic source. If $\Sigma_0 = I$, then $Q(t)$, $Q'(t)$, $A(t)$, and Σ_1 share the eigenbasis of Σ_1 . Simultaneous diagonalization with the identity is not an extra hypothesis. In that basis, mode i with eigenvalue $\lambda_i > 0$ obeys $x'_i = a_i(t)x_i + c_i(t)$ with $q_i(t) = \alpha(t)^2 + \lambda_i\sigma(t)^2$ and $a_i(t) = q'_i(t)/(2q_i(t))$. The covariance drift formula depends only on the variance schedule. It is independent of the mean path $c(t) = \dot{m}_t - A(t)m_t$. The exact one-step factor on $[t_n, t_{n+1}]$ is $\sqrt{q_i(t_{n+1})/q_i(t_n)}$. For centered modes the exact endpoint factor is $\sqrt{\lambda_i}$.

Regularity. Following Chen et al. [2025, Definition 3.2],

$$\mathcal{R}[b] = \int_0^1 \|A(t)\|_2^2 dt. \quad (4)$$

For state-independent $A(t)$ this coincides with their path expectation. By definition $\mathcal{R}$ ignores $c(t)$. For commuting $A(t)$, $\|A(t)\|_2 = \max_i |a_i(t)|$. Headline comparisons in this paper use the continuous integral (4). The historical 24-node trapezoidal estimator is written $\widehat{\mathcal{R}}_{24}$ and is used only as an implementation check. No trapezoidal value is described as an exact integral.

Any admissible one-dimensional variance path from 1 to $\lambda > 0$ obeys $\int_0^1 a dt = \frac{1}{2} \ln \lambda$. Therefore

$$\mathcal{R} = \int_0^1 a^2 dt = \left(\int_0^1 a dt \right)^2 + \text{Var}_t(a) \geq \left(\frac{1}{2} \ln \lambda \right)^2, \quad (5)$$

with equality if and only if a is constant. The minimizer is $q(t) = \lambda^t$, $a \equiv \frac{1}{2} \ln \lambda$, which is Chen et al. [2025, Example 3.3]. In several dimensions the per-mode schedule $q_i(t) = \lambda_i^t$ yields $\mathcal{R} = \left(\frac{1}{2} \max_i |\ln \lambda_i| \right)^2$. That schedule is not in general a shared (α, σ) interpolant.

Solver, step, and NFE. Euler uses $s = 1$ field evaluation per step, Heun $s = 2$, and classical RK4 $s = 4$. A budget B is admitted only if $B \equiv 0 \pmod{s}$. The step count is $S = B/s$ and the grid is the uniform partition of $[0, 1]$ with S steps. Equal NFE is not equal step count: at $B = 8$, Euler takes 8 steps, Heun 4, and RK4 2.

Endpoint error. A solver applied to (3) produces an affine endpoint map $x \mapsto Rx + d$. Source moments are pushed through this map. Error is the Gaussian W_2 of the endpoint *laws* [Gelbrich, 1990, Peyré and Cuturi, 2019],

$$W_2^2 = \|\hat{m} - \mu_1\|_2^2 + \text{tr}(\hat{C} + \Sigma_1 - 2(\Sigma_1^{1/2}\hat{C}\Sigma_1^{1/2})^{1/2}). \quad (6)$$

This is neither pathwise trajectory error nor local truncation error. In the centered commuting case, $W_2^2 = \sum_i (|r_i| - \sqrt{\lambda_i})^2$.

Question. Does $\mathcal{R}[b] < \mathcal{R}[b']$ imply $E[b] < E[b']$ at equal NFE? The answer is no, already in dimension one.

3 A one-dimensional Gaussian ranking counterexample

Assumption 1. *The endpoints $X_0 \sim N(0, 1)$ and $X_1 \sim N(0, 4)$ are independent. The compared schedules are linear and trigonometric VP. The solver is explicit Heun with NFE $B = 8$, hence $S = 4$ steps of size $h = 1/4$.*

Under Assumption 1, $q(t) = \alpha(t)^2 + 4\sigma(t)^2$ and $a(t) = q'(t)/(2q(t))$. The exact endpoint factor is 2. All Heun factors below are positive, so $W_2 = ||r| - 2| = |r - 2|$.

Proposition 1. *Under Assumption 1,*

$$\mathcal{R}_{\text{lin}} = \frac{5\pi}{8} - 1 \approx 0.9634954084936207, \quad (7)$$

$$\mathcal{R}_{\text{VP}} = \frac{\pi^2}{16} \approx 0.6168502750680849, \quad (8)$$

so averaged regularity prefers VP. The Heun endpoint factors satisfy

$$r_{\text{lin}} = \frac{6797469}{3559400}, \quad 0 < r_{\text{VP}} < \frac{187}{100}, \quad (9)$$

the second bound being a rational enclosure obtained by substituting $\sqrt{2} < 99/70$ and $\pi < 355/113$ into the exact element of $\mathbb{Q}[\pi, \sqrt{2}]$. Consequently

$$W_2^{\text{lin}} = \frac{321331}{3559400} < 0.091 < 0.130 < W_2^{\text{VP}}, \quad (10)$$

so equal-NFE Gaussian W_2 prefers the linear path. The regularity ordering does not determine the fixed-NFE W_2 ordering.

Proof sketch. The linear drift is $a(t) = (5t - 1)/(1 - 2t + 5t^2)$. Completing the square to $5(t - 1/5)^2 + 4/5$ and using $\arctan 2 + \arctan(1/2) = \pi/2$ yields $\mathcal{R}_{\text{lin}} = 5\pi/8 - 1$. The VP integral evaluates to $\pi^2/16$. The four linear Heun factors are the rationals $47/52$, $6/5$, $497/370$, and $97/74$, whose product is $6797469/3559400$. Then $321331/3559400 < 91/1000$ is an elementary comparison of integers. The VP grid drifts are exact elements of $\mathbb{Q}[\pi, \sqrt{2}]$; the four-step Heun product has only nonnegative coefficients, so substituting the rational upper bounds $\sqrt{2} < 99/70$ and $\pi < 355/113$ is valid and gives $r_{\text{VP}} < 187/100$, hence $W_2^{\text{VP}} > 13/100$. Details occupy Appendix A. $\square$

One certified inversion proves the logical non-implication. The remainder of the paper shows that the phenomenon is not an artefact of this scalar pair, and identifies the information that $\mathcal{R}$ cannot retain.




Figure 1: Proposition 1: exact $\mathcal{R}$ (left) and Heun W_2 at NFE 8 (right). The VP error bar is the certified lower bound $2 - P(355/113, 99/70)$. Regularity prefers VP; endpoint error prefers linear.

4 Modal analysis and transported defects

Let $e_{i,n} = \sqrt{q_i(t_{n+1})/q_i(t_n)}$ be the exact factor of mode i and let $r_{i,n}$ be the numerical one-step factor (Appendix B). The algebraic identity

$$\prod_{n=0}^{S-1} r_{i,n} - \prod_{n=0}^{S-1} e_{i,n} = \sum_{n=0}^{S-1} (r_{i,n} - e_{i,n}) \left(\prod_{j<n} r_{i,j} \right) \left(\prod_{j>n} e_{i,j} \right) \quad (11)$$

holds without a non-vanishing hypothesis: it is the telescoping expansion of a product. Prefixes are numerical; suffixes are exact. Endpoint factor error is therefore a transported sum of signed, solver-stage-dependent local defects.

Identity (11) is an accounting tool. The content of the inversion is the leading exact-minus-method coefficient: $\frac{1}{2}(a' + a^2)h^2$ for Euler and $(-a''/12 + a^3/6)h^3$ for the implemented Heun factor $1 + (h/2)(a(t) + a(t+h)(1 + ha(t)))$ (Appendix C). They involve different derivatives, signs, and stage locations. Averaging $\|A\|_2^2$ discards stage times, signs, cancellation, and transport.

For any independent linear coupling, $a_{\text{lin}}(0) = -1 < 0 < a_{\text{lin}}(1) = +1$. The VP drift vanishes at the endpoints and does not change sign on $(0, 1)$. The unsigned average therefore erases a qualitative distinction that every fixed-stage Runge–Kutta method samples.

Figure 2 shows the signed transported terms on the strongest centered inversion (low-rank, $d = 8$, Euler, NFE 8). The identity is exact for that block; the figure is an illustration, not an independent causal test.

A complementary scalar construction (Appendix D) shows non-implication for left-endpoint Euler on a grid-aware oscillatory field. The field $a_{1,N}$ is chosen after the grid. That construction is not the mechanism of Proposition 1 or of the Gaussian benchmark.

5 The $\mathcal{R}$ -minimizer and the inversion region

The Cauchy–Schwarz argument of Section 2 identifies a third path: the per-mode log-covariance schedule $q_i(t) = \lambda_i^t$. On the four centered geometries of Section 6 it has the smallest continuous $\mathcal{R}$ in every block and the smallest Gaussian W_2 in all 36 solver-budget conditions (Appendix J). Linear versus VP remain free to invert.

A one-dimensional sweep $\lambda \in [0.05, 100]$ locates those inversions (Figure 3). At $\lambda = 4$, the inversion is present under Heun at every tested budget $\text{NFE} \in \{4, 8, 12, 16, 24, 32, 64, 128, 256\}$, absent under Euler, and mixed under RK4 (absent at NFE 8 and 12). Euler *does* invert at more




Figure 2: Modal contributions to squared Wasserstein error and transported signed defects for the strongest centered inversion. The bars reconstruct the telescoping identity (11).




Figure 3: Linear versus VP ranking inversion versus target variance λ . Dark cells: $\mathcal{R}$ and W_2 disagree. The dashed line is $\lambda = 4$ (Proposition 1).

extreme λ on the same sweep. The scalar example is therefore solver-dependent, not an artefact of Heun at budget 8 alone.

6 Controlled multi-configuration benchmark

Geometries. Source $N(0, I)$.

Four centered targets, with seed 271828 for low-rank factors: anisotropic Gaussians with condition number 4 in dimensions 2 and 8 (eigenvalues a geometric sequence from $1/2$ to 2); and low-rank covariances $FF^\top + 0.05I$ of rank 2. The dimension-2 low-rank eigenvalues are 0.2914701258115777 and 2.9677106074567434; the dimension-8 low-rank eigenvalues are 0.05 (six times), 4.64827725629505, and 10.675638226546441. The covariance matrices are recorded in `paper/arxiv/artifacts/geometries.json`. Because $\mathcal{R}$ uses the spectral norm, the two anisotropic dimensions share the same $\mathcal{R}$ pair (0.972296, 0.192270) while their W_2 values differ. The grid therefore contains 3 distinct linear-versus-VP $\mathcal{R}$ comparisons.

Hierarchy. The 12 geometry $\times$ solver cells comprise 4 geometries and three solvers, each evaluated at NFE 8, 16, and 32. Linear versus VP invert in 5 of those 12 cells; 4 invert at every

Table 1: Strongest centered inversion: low-rank, $d = 8$, Euler, NFE 8. $\mathcal{R}$ is the continuous integral; $\widehat{\mathcal{R}}_{24}$ is shown only as a check. Eigenvalues: 0.05 (six times), 4.64827725629505, 10.675638226546441.

path	$\mathcal{R}$	$\widehat{\mathcal{R}}_{24}$	Gaussian W_2
linear	2.9441044083	2.9476523251	0.8108540111
VP	4.7305438136	4.7295206355	0.4564779075




Figure 4: Signed low-rank W_2 margin at equal NFE. Positive values prefer VP; negative values prefer linear. Vertical axis: symmetric-log with linear threshold 10^{-4} .

tested NFE. The 14 inverted solver-budget rows (Table 2) are a hierarchical listing of those cells, not independent statistical trials and not a population frequency. Anisotropic $d = 8$ Heun inverts at NFE 16 and 32 but not at NFE 8, so the headline cell count is “at least one inversion”, not “all three NFE agree”.

Continuous $\mathcal{R}$ versus $\widehat{\mathcal{R}}_{24}$. Adaptive quadrature of $\max_i a_i(t)^2$ agrees in sign with $\widehat{\mathcal{R}}_{24}$ on every geometry, including the scalar pair of Proposition 1. Trapezoidal estimates at 24, 48, 96, and 192 nodes converge to the same ordering. Inversion counts below use continuous $\mathcal{R}$.

Results. Independent grouping of the frozen reproduction yields the 5 inverting cells above, listed row-wise in Table 2. The strongest inversion is low-rank, $d = 8$, Euler, NFE 8 (Table 1): $\mathcal{R}$ prefers linear while W_2 prefers VP, with margin 0.3543761036. All 72 float64 W_2 values agree with an 80-digit reference to at most $9.7050430488 \times 10^{-10}$. The smallest inversion margin, $1.1188920612 \times 10^{-5}$, is 11528 times that gap.

On all 18 low-rank primary blocks, VP has smaller W_2 under Euler and linear has smaller W_2 under Heun and RK4 (Figure 4). A post-hoc robustness grid repeats the split on 66 low-rank blocks: two dimensions, three solvers, three primary budgets each at noise-floor perturbations $\delta \in \{-0.1, 0, 0.1\}$ (the factor F is redrawn from the same seed; noise variance becomes $0.05(1+\delta)$), plus NFE 64 and 128 at $\delta = 0$ only. That count is $2 \cdot 3 \cdot (3 \cdot 3 + 2) = 66$. It is not preregistered evidence.

Mixture targets are excluded. Dimension-8 mixture endpoint estimators failed a post-result calibration diagnostic; Phase 4 reproduction accepts Gaussian targets only.

7 Non-centered structural stress test

The quantity $\mathcal{R}$ depends only on $A(t)$. If $c(t) \neq 0$, the mean still contributes $\|\hat{m} - \mu_1\|_2^2$ to W_2^2 . The following family was specified in the repository before execution (commit `f5e857c` at 2026-07-24 15:28, results `8b87c7f` at 16:04). Amendment A1 (`508101e`, 15:33) changed only an

auxiliary endpoint sanity check from a W_2 threshold, which float64 identical Gaussians cannot meet, to a moment gap. The family itself was not changed. The accurate description is a pre-specified stress test of a structural blind spot of $\mathcal{R}$, not an independent replication of the centered mechanism.

Construction. Dimensions $d \in \{2, 8\}$, with coordinates indexed by $i = 0, \dots, d - 1$. Source $N(\mu_0, I)$ with $\mu_{0,i} = 0.75(-1)^i$. Target $N(\mu_1, D)$ with $\mu_{1,i} = 1 + 0.25i$ and $D = \text{diag}(\lambda_i)$, where the λ_i form a geometric sequence from $6^{-1/2}$ to $6^{1/2}$ (condition number 6). Then $c(t) \neq 0$. Paths are linear and trigonometric VP. Solvers are Euler, Heun, and RK4 at NFE $B \in \{8, 16, 32\}$, with $S = B/s$ uniform steps of size $1/S$ integrated forward in t ($s = 1, 2, 4$). Reported W_2 values and the mean/Bures split are decomposed from the frozen run `outputs/workshop_external_validation_2026-07-24-v1/`; continuous $\mathcal{R}$ is recomputed from the same eigenvalues.

Signed differences. Write $M = \|\hat{m} - \mu_1\|_2^2$ and $C = W_2^2 - M$, and

$$\Delta_{\text{mean}} = M_{\text{lin}} - M_{\text{VP}}, \quad \Delta_{\text{cov}} = C_{\text{lin}} - C_{\text{VP}}, \quad \Delta_{\text{total}} = \Delta_{\text{mean}} + \Delta_{\text{cov}}. \quad (12)$$

A block is mean-driven if $\Delta_{\text{mean}}\Delta_{\text{total}} > 0$ and $|\Delta_{\text{mean}}| > |\Delta_{\text{cov}}|$, and covariance-driven if $\Delta_{\text{cov}}\Delta_{\text{total}} > 0$ and $|\Delta_{\text{cov}}| > |\Delta_{\text{mean}}|$. Otherwise it is mixed. The same paths, solvers, and budgets give 18 blocks, of which 11 invert under continuous $\mathcal{R}$. Table 3 reports the driver of each path preference. Several inversions, including all dimension-8 Euler and Heun inversions, are mean-driven: $\mathcal{R}$ cannot see them.

8 Relation to existing work

Flow matching and stochastic interpolants supply the Gaussian path family [Lipman et al., 2023, Albergo et al., 2025, Liu et al., 2023, Tong et al., 2024]. Lipschitz-guided schedule design proposes the selection criterion tested here [Chen et al., 2025]. Solver-specific schedules and exact linear parts are documented in diffusion practice [Sabour et al., 2024, Karras et al., 2022, Lu et al., 2022a,b, Zhang and Chen, 2023].

Stéphanovitch [2026] prove sharp Lipschitz regularity of canonical flow-matching fields and Euler-type W_2 rates of order $\sqrt{d}/N$ up to logs. That is a bound, not a pairwise ranking of two interpolants at equal NFE. Gupta et al. [2026] allocate Euler steps where a learned field is sharp; the path is held fixed. Yuan et al. [2026] learn instance-aware grids. Du et al. [2026] select paths by a least-action principle. Khan [2026] regularizes the material derivative of learned fields.

Tao and Choi [2026] decompose $\nabla v = S + \Omega$ and control Euler error by the logarithmic norm $\mu_2 = \lambda_{\max}(S)$, which is signed, whereas $\mathcal{R}$ integrates $\|A\|_2^2$. In one dimension $\mu_2(a) = a$ and $\|A\|_2 = |a|$. Their Theorem 5 states that McCann displacement interpolation has vanishing material derivative, hence Euler is of order 2 in the Eulerian analysis; Remark 7 records the stronger Lagrangian exactness observed on that interpolant. Neither statement applies to the independent-coupling linear or VP path studied here. The present paper supplies a rationally certified Gaussian instance of ranking failure for the unsigned average $\|A\|_2^2$. A counterexample to μ_2 itself is outside our scope.

9 Limitations

The systems are commuting affine Gaussian interpolants, three fixed-step Runge–Kutta methods, and small equal-NFE budgets. No learned neural field, image, video, or world-model quantity is reported. Chen et al. [2025] prove no discretization-error bound and do not claim a universal pairwise ranking at fixed NFE; the example does not invalidate A_2 as an optimization objective




Figure 5: Signed path differences on all 18 non-centered blocks. $\Delta_{\text{mean}} = M_{\text{lin}} - M_{\text{VP}}$ and $\Delta_{\text{cov}} = C_{\text{lin}} - C_{\text{VP}}$. A positive bar means the linear path has the larger contribution; a negative bar means VP has the larger contribution. The axis is symmetric-log with linear threshold 10^{-6} . Stars mark ranking inversions. Exact numerical values appear in Table 4.

within a path family. The Gaussian family is a probe in which $\mathcal{R}$ and W_2 are closed form; in a learned field the linear part is not exact, so exponential integrators that solve an affine scalar mode exactly are not a drop-in substitute. Cell counts and the 14 inverted rows are hierarchical descriptive summaries, not population rates. The scalar inversion at $\lambda = 4$ is solver-dependent: present under Heun at every tested budget, absent under Euler at that λ , and mixed under RK4. The Euler construction of Appendix D is grid-aware: the field is chosen after the grid. Robustness NFE 64 and 128 and $\pm 10\%$ noise-floor perturbations are post-hoc. A pre-specified non-centered family (Section 7) shows a structural blind spot of $\mathcal{R}$ for the drift offset $c(t)$; it is not an independent replication of the centered mechanism.

10 Conclusion

Averaged squared Jacobian regularity does not determine the equal-NFE Gaussian W_2 ranking of the linear and VP interpolants. The fact is proved by a one-dimensional Heun calculation whose integrals and linear factor are exact and whose VP factor is enclosed by a rational substitution in $\mathbb{Q}[\pi, \sqrt{2}]$. On the tested grid the Cauchy–Schwarz minimizer of $\mathcal{R}$ dominates in W_2 , while linear versus VP invert in 5 of 12 geometry×solver cells. Thus $\mathcal{R}$ is reliable as an objective on this Gaussian family and not reliable as a comparator of linear versus VP away from that optimum.

11 Reproducibility statement

Default numerical precision for the benchmark runs is float64. The scalar calculation uses exact rational and algebraic expressions together with validated bounds; the precision audit uses 80-digit arithmetic. Compact artifacts used by this PDF live in the repository under `paper/arxiv/artifacts/` and are generated by `scripts/make_arxiv_compact_artifacts.py`, `scripts/run_log_covariance_comparison.py`, `scripts/run_inversion_region.py`, and `scripts/run_lowrank_seed_fraction.py`. They are committed files. A GitHub Release tag is a later owner publication gate, after workshop notification, and is not required to read the numbers in this PDF. Experiment-code commit of the frozen Phase 4 grid: `e48c9390e62b38f206342e6aeb7f160122ccc79c`. The Hydra `outputs/` tree is not part of the public clone; eigenvalues, exact-moment modal Runge–Kutta, and the compact JSON files suffice for Table 1 and Proposition 1.

```
python -m pip install -e ".[dev]"
pytest
python scripts/verify_scalar_counterexample.py
python scripts/run_log_covariance_comparison.py
python scripts/run_inversion_region.py
python scripts/run_lowrank_seed_fraction.py
python scripts/make_arxiv_figures.py
python scripts/check_arxiv_placeholder.py
python scripts/check_arxiv_release.py
```

The placeholder check must pass with no retired commit token remaining. The release check verifies compact checksums. Tag existence is a manual publication gate (`-require-tag`), not a scientific failure of this manuscript. The compact-artifact builder `make_arxiv_compact_artifacts.py` additionally reads frozen Phase 4 JSON when those files are present; it is not required to recompute Proposition 1 or the three-path grid.

References

- Michael S. Albergo, Nicholas M. Boffi, and Eric Vanden-Eijnden. Stochastic interpolants: A unifying framework for flows and diffusions. *Journal of Machine Learning Research*, 26(209): 1–80, 2025. arXiv:2303.08797.
- Yifan Chen, Eric Vanden-Eijnden, and Jiawei Xu. Lipschitz-guided design of interpolation schedules in generative models. *arXiv preprint arXiv:2509.01629*, 2025. version 3, 16 May 2026.
- Shukai Du, Junzhe Zhang, and Yiming Li. Lagrangian flow matching: A least-action framework for principled path design. *arXiv preprint arXiv:2605.15419*, 2026.
- Matthias Gelbrich. On a formula for the L^2 Wasserstein metric between measures on Euclidean and Hilbert spaces. *Mathematische Nachrichten*, 147:185–203, 1990. doi: 10.1002/mana.19901470121.
- Aditi Gupta, Soon Hoe Lim, Annan Yu, and N. Benjamin Erichson. Sharpen your flow: Sharpness-aware sampling for flow matching. *arXiv preprint arXiv:2605.11547*, 2026.
- Ernst Hairer, Syvert P. Nørsett, and Gerhard Wanner. *Solving Ordinary Differential Equations I: Nonstiff Problems*. Springer, 2nd edition, 1993.

- Tero Karras, Miika Aittala, Timo Aila, and Samuli Laine. Elucidating the design space of diffusion-based generative models. In *Advances in Neural Information Processing Systems*, 2022. arXiv:2206.00364.
- Tauhid Khan. Isokinetic flow matching for pathwise straightening of generative flows. *arXiv preprint arXiv:2604.04491*, 2026.
- Yaron Lipman, Ricky T. Q. Chen, Heli Ben-Hamu, Maximilian Nickel, and Matthew Le. Flow matching for generative modeling. In *International Conference on Learning Representations (ICLR)*, 2023. arXiv:2210.02747.
- Xingchao Liu, Chengyue Gong, and Qiang Liu. Flow straight and fast: Learning to generate and transfer data with rectified flow. In *International Conference on Learning Representations (ICLR)*, 2023. arXiv:2209.03003.
- Cheng Lu, Yuhao Zhou, Fan Bao, Jianfei Chen, Chongxuan Li, and Jun Zhu. DPM-solver: A fast ODE solver for diffusion probabilistic model sampling in around 10 steps. In *Advances in Neural Information Processing Systems*, 2022a. arXiv:2206.00927.
- Cheng Lu, Yuhao Zhou, Fan Bao, Jianfei Chen, Chongxuan Li, and Jun Zhu. DPM-solver++: Fast solver for guided sampling of diffusion probabilistic models. *arXiv preprint arXiv:2211.01095*, 2022b.
- Gabriel Peyré and Marco Cuturi. *Computational Optimal Transport*. Now Publishers, 2019. arXiv:1803.00567.
- Amirmojtaba Sabour, Sanja Fidler, and Karsten Kreis. Align your steps: Optimizing sampling schedules in diffusion models. In *International Conference on Machine Learning (ICML)*, 2024. arXiv:2404.14507.
- Yang Song, Jascha Sohl-Dickstein, Diederik P. Kingma, Abhishek Kumar, Stefano Ermon, and Ben Poole. Score-based generative modeling through stochastic differential equations. In *International Conference on Learning Representations (ICLR)*, 2021. arXiv:2011.13456.
- Arthur Stéphanovitch. Lipschitz regularity in flow matching and diffusion models: Sharp sampling rates and functional inequalities. *arXiv preprint arXiv:2604.06065*, 2026.
- Chenxi Tao and Seung-Kyum Choi. On the role of strain and vorticity in numerical integration error for flow matching. *arXiv preprint arXiv:2605.06680*, 2026.
- Alexander Tong, Kilian Fatras, Nikolay Malkin, Guillaume Hugué, Yanlei Zhang, Jarrod Rector-Brooks, Guy Wolf, and Yoshua Bengio. Improving and generalizing flow-based generative models with minibatch optimal transport. *Transactions on Machine Learning Research*, 2024. arXiv:2302.00482.
- Liangyu Yuan, Ruoyu Wang, Tong Zhao, Dingwen Fu, Mingkun Lei, Beier Zhu, and Chi Zhang. Few-step diffusion sampling through instance-aware discretizations. In *Proceedings of the IEEE/CVF Conference on Computer Vision and Pattern Recognition (CVPR)*, pages 35882–35892, 2026. arXiv:2603.17671.
- Qinsheng Zhang and Yongxin Chen. Fast sampling of diffusion models with exponential integrator. In *International Conference on Learning Representations (ICLR)*, 2023. arXiv:2204.13902.

A Proof of Proposition 1

Linear drift and integral. $q_{\text{lin}}(t) = (1 - t)^2 + 4t^2 = 1 - 2t + 5t^2$ and $q' = -2 + 10t$, so $a_{\text{lin}}(t) = (5t - 1)/q_{\text{lin}}(t)$. Completing the square yields the two equivalent identities

$$q_{\text{lin}}(t) = 5 \left[\left(t - \frac{1}{5} \right)^2 + \left(\frac{2}{5} \right)^2 \right], \quad (13)$$

$$q_{\text{lin}}(t) = 5 \left(t - \frac{1}{5} \right)^2 + \frac{4}{5}. \quad (14)$$

Thus $a_{\text{lin}}(t) = (t - 1/5)/((t - 1/5)^2 + (2/5)^2)$. The substitution $u = t - 1/5 = (2/5) \tan \theta$ gives $du = (2/5) \sec^2 \theta d\theta$ and converts $\int a_{\text{lin}}^2$ into $(5/2) \int \sin^2 \theta d\theta$. The definite limits are $\theta(0) = \arctan(-1/2)$ and $\theta(1) = \arctan 2$. The antiderivative $(5/4)(\theta - \sin \theta \cos \theta)$ therefore contributes $(5/4)(\arctan 2 + \arctan(1/2)) - (5/4)(4/5)$. Both angles lie in $(0, \pi/2)$ and $\tan(\arctan 2 + \arctan(1/2))$ is undefined, so the sum is $\pi/2$. Hence $\mathcal{R}_{\text{lin}} = 5\pi/8 - 1$. (The subtracted 1 is the boundary evaluation of $\sin \theta \cos \theta = \tan \theta / (1 + \tan^2 \theta)$.)

VP drift and integral. $q_{\text{VP}}(t) = \cos^2(\pi t/2) + 4 \sin^2(\pi t/2) = (5 - 3 \cos(\pi t))/2$ and $q' = (3\pi/2) \sin(\pi t)$, hence

$$a_{\text{VP}}(t) = \frac{(3\pi/2) \sin(\pi t)}{5 - 3 \cos(\pi t)} = \frac{(3\pi/4) \sin(\pi t)}{q_{\text{VP}}(t)}. \quad (15)$$

The change of variables $u = \pi t$ yields $\mathcal{R}_{\text{VP}} = (9\pi/4) \int_0^\pi \sin^2 u / (5 - 3 \cos u)^2 du$. The standard Weierstrass evaluation of the integrand gives $\int_0^\pi \sin^2 u / (5 - 3 \cos u)^2 du = \pi/36$, hence $\mathcal{R}_{\text{VP}} = \pi^2/16$. Both closed forms were checked by adaptive quadrature (absolute discrepancy of the values below 10^{-12} ; the VP error *estimate* returned by `scipy.integrate.quad` can be larger than that discrepancy and is not used as a certificate).

Heun recurrence. On $x' = a(t)x$ the implemented Heun factor is $r = 1 + (h/2)(a(t_n) + a(t_{n+1})(1 + ha(t_n)))$. With $h = 1/4$ the four linear factors are the rationals $47/52$, $6/5$, $497/370$, and $97/74$, all positive, with product $6797469/3559400$. Then $2 - r = 321331/3559400$, and $321331 \cdot 1000 < 91 \cdot 3559400$ is an integer comparison, so $W_2^{\text{lin}} < 0.091$.

Exact VP grid and rational certificate. The VP drift is $a_{\text{VP}}(t) = (3\pi/2) \sin(\pi t)/(5 - 3 \cos(\pi t))$. Evaluating at the Heun nodes gives

$$a(0) = 0, \quad (16)$$

$$a(1/4) = \frac{(9 + 15\sqrt{2})\pi}{82}, \quad (17)$$

$$a(1/2) = \frac{3\pi}{10}, \quad (18)$$

$$a(3/4) = \frac{(-9 + 15\sqrt{2})\pi}{82}, \quad (19)$$

$$a(1) = 0. \quad (20)$$

Positivity of $a(3/4)$ follows from $15\sqrt{2} > 9$, or $15^2 \cdot 2 - 9^2 = 369 > 0$ after squaring. Each Heun factor is therefore strictly positive, so $0 < r_{\text{VP}}$ and $W_2^{\text{VP}} = 2 - r_{\text{VP}}$. The four-step product is an element $P(\pi, \sqrt{2})$ of $\mathbb{Q}[\pi, \sqrt{2}]$. Expanding and reducing $\sqrt{2}^2 = 2$ produces twelve monomials, all with nonnegative coefficients. Substituting any pair of upper bounds for π and $\sqrt{2}$ is therefore valid. The bound $\sqrt{2} < 99/70$ is the integer comparison $99^2 - 2 \cdot 70^2 = 1$. The bound $\pi < 355/113$ follows from Machin's formula $\pi = 16 \arctan(1/5) - 4 \arctan(1/239)$ together with Leibniz remainders: seven Taylor terms upper-bound $\arctan(1/5)$ and four terms

lower-bound $\arctan(1/239)$. The resulting rational is strictly less than $355/113$ by an integer comparison. Substituting these enclosures yields

$$P\left(\frac{355}{113}, \frac{99}{70}\right) = \frac{192113353671412470139303}{102753427879939708813312} < \frac{187}{100}, \quad (21)$$

equivalently the integer comparison

$$19211335367141247013930300 < 19214891013548725548089344. \quad (22)$$

Hence $r_{\text{VP}} < 187/100$ and $W_2^{\text{VP}} > 13/100$. Coefficient records occupy the compact artefact `scalar_counterexample.json`. An independent `mpmath.iv` enclosure at 40 decimal digits, using `mpmath` 1.3.0, and an 80-digit non-interval product, agree with this bound and with the package `float64` implementation to 10^{-14} . They are cross-checks, not the proof.

B Solver formulas

On $[t, t+h]$ for $x' = a(t)x$,

$$r_{\text{Euler}} = 1 + ha(t), \quad (23)$$

$$r_{\text{Heun}} = 1 + \frac{h}{2}(a(t) + a(t+h)(1 + ha(t))), \quad (24)$$

and classical RK4 with stages at t , $t + h/2$ (twice), and $t + h$. Equal NFE requires $B \equiv 0 \pmod{s}$ and sets $S = B/s$.

C Local truncation expansions

Assume a is C^3 . The exact factor admits $1 + ha + (h^2/2)(a' + a^2) + (h^3/6)(a'' + 3aa' + a^3) + O(h^4)$. Subtracting Euler leaves $\frac{1}{2}(a' + a^2)h^2 + O(h^3)$. Substituting $a(t+h)$ into the implemented Heun factor leaves $(-a''/12 + a^3/6)h^3 + O(h^4)$. The aa' terms cancel. The expansion was confirmed with independent symbolic algebra on the implemented expression.

D Grid-aware Euler construction

Proposition 2. *Fix $L > 0$ and an integer $N \geq 1$. Let $h = 1/N$ and let forward Euler use the left-endpoint grid $t_n = nh$ on $[0, 1]$. Set $\varepsilon_N = N(e^{L/N} - 1) - L > 0$ and define $a_0(t) = L$ and $a_{1,N}(t) = L + \varepsilon_N \cos(2\pi Nt)$. For $x(0) = 1$: both exact endpoints equal e^L ; $\int_0^1 a_{1,N}^2 = L^2 + \varepsilon_N^2/2 > \int_0^1 a_0^2$; Euler is exact for $a_{1,N}$; Euler has strictly positive endpoint error for a_0 .*

Proof. The solution of $x' = a(t)x$, $x(0) = 1$, is $x(t) = \exp(\int_0^t a(s) ds)$. The cosine perturbation has integer period, so $\int_0^1 \cos(2\pi Nt) dt = 0$ and both exact endpoints equal e^L . Expanding the square and using the same orthogonality gives $\int_0^1 a_{1,N}^2 = L^2 + \varepsilon_N^2/2$. The inequality $\varepsilon_N > 0$ is $e^z > 1 + z$ at $z = L/N > 0$, hence $\int_0^1 a_{1,N}^2 > \int_0^1 a_0^2 = L^2$.

On the left-endpoint Euler grid, $\cos(2\pi Nt_n) = \cos(2\pi n) = 1$, so $1 + ha_{1,N}(t_n) = 1 + (L + \varepsilon_N)/N = e^{L/N}$. Every oscillatory Euler factor equals $e^{L/N}$ and the product is e^L . For the constant field every Euler factor equals $1 + L/N$, and the product is $(1 + L/N)^N$. Since $\log(1 + z) < z$ for $z = L/N > 0$, one has $(1 + L/N)^N < e^L$. $\square$

The field $a_{1,N}$ is chosen after the grid. The construction is not the mechanism of Proposition 1. An 80-digit check confirms that the oscillatory Euler product equals e^L , but it is not a substitute for the argument above.

E Centered inversion table

Table 2 lists every inverted solver-budget row under continuous $\mathcal{R}$. These rows are a hierarchical listing of the 5 geometry×solver cells, not a population rate.

Table 2: All 14 inverted solver-budget rows among 36 centered blocks, grouped into 5 of 12 geometry×solver cells.

family	d	solver	NFE	R prefers	W_2 prefers	W_2 margin
anisotropic	2	heun	8	VP	linear	1.905310×10^{-3}
anisotropic	2	heun	16	VP	linear	1.939806×10^{-3}
anisotropic	2	heun	32	VP	linear	6.634439×10^{-4}
anisotropic	8	heun	16	VP	linear	1.758942×10^{-3}
anisotropic	8	heun	32	VP	linear	7.628173×10^{-4}
low-rank	2	heun	8	VP	linear	1.836251×10^{-2}
low-rank	2	heun	16	VP	linear	6.039925×10^{-3}
low-rank	2	heun	32	VP	linear	1.698035×10^{-3}
low-rank	2	rk4	8	VP	linear	1.516722×10^{-3}
low-rank	2	rk4	16	VP	linear	1.289981×10^{-4}
low-rank	2	rk4	32	VP	linear	1.118892×10^{-5}
low-rank	8	euler	8	linear	VP	3.543761×10^{-1}
low-rank	8	euler	16	linear	VP	2.046946×10^{-1}
low-rank	8	euler	32	linear	VP	1.096576×10^{-1}

F Non-centered blocks

Table 3: Non-centered family: all 18 blocks, with the signed-difference driver of the W_2^2 path preference ($\Delta_{\text{mean}}, \Delta_{\text{cov}}$).

d	solver	NFE	inversion	W_2 prefers	W_2^2 driver
2	euler	8	no	VP	covariance
2	euler	16	no	VP	covariance
2	euler	32	no	VP	covariance
2	heun	8	yes	linear	mean
2	heun	16	yes	linear	mean
2	heun	32	yes	linear	covariance
2	rk4	8	no	VP	covariance
2	rk4	16	no	VP	covariance
2	rk4	32	yes	linear	covariance
8	euler	8	yes	linear	mean
8	euler	16	yes	linear	mean
8	euler	32	yes	linear	mean
8	heun	8	yes	linear	mean
8	heun	16	yes	linear	mean
8	heun	32	yes	linear	mean
8	rk4	8	no	VP	covariance
8	rk4	16	no	VP	covariance
8	rk4	32	yes	linear	mean

Table 4: Signed path differences for the non-centered family. $\Delta_{\text{mean}} = M_{\text{lin}} - M_{\text{VP}}$, $\Delta_{\text{cov}} = C_{\text{lin}} - C_{\text{VP}}$, $\Delta_{\text{total}} = \Delta_{\text{mean}} + \Delta_{\text{cov}}$.

d	solver	NFE	inv.	Δ_{mean}	Δ_{cov}	Δ_{total}
2	euler	8	no	-1.969780×10^{-2}	7.008429×10^{-2}	5.038649×10^{-2}
2	euler	16	no	-5.010761×10^{-3}	1.895811×10^{-2}	1.394734×10^{-2}
2	euler	32	no	-1.266182×10^{-3}	4.941067×10^{-3}	3.674885×10^{-3}
2	heun	8	yes	-1.401380×10^{-3}	-8.296535×10^{-4}	-2.231034×10^{-3}
2	heun	16	yes	-8.132945×10^{-5}	-7.740260×10^{-5}	-1.587321×10^{-4}
2	heun	32	yes	-4.800285×10^{-6}	-5.428459×10^{-6}	-1.022874×10^{-5}
2	rk4	8	no	-1.931000×10^{-6}	7.169013×10^{-5}	6.975913×10^{-5}
2	rk4	16	no	$-5.235990 \times 10^{-10}$	1.106023×10^{-8}	1.053664×10^{-8}
2	rk4	32	yes	$-7.527744 \times 10^{-12}$	$-5.405709 \times 10^{-11}$	$-6.158484 \times 10^{-11}$
8	euler	8	yes	-3.252232×10^{-1}	2.207633×10^{-1}	-1.044599×10^{-1}
8	euler	16	yes	-8.388778×10^{-2}	5.947257×10^{-2}	-2.441521×10^{-2}
8	euler	32	yes	-2.129903×10^{-2}	1.546831×10^{-2}	-5.830720×10^{-3}
8	heun	8	yes	-5.927520×10^{-3}	-4.969512×10^{-4}	-6.424471×10^{-3}
8	heun	16	yes	-3.450747×10^{-4}	-9.787629×10^{-5}	-4.429510×10^{-4}
8	heun	32	yes	-2.074391×10^{-5}	-7.576731×10^{-6}	-2.832065×10^{-5}
8	rk4	8	no	-3.909251×10^{-6}	3.088406×10^{-4}	3.049314×10^{-4}
8	rk4	16	no	-1.137695×10^{-8}	7.433020×10^{-8}	6.295325×10^{-8}
8	rk4	32	yes	$-7.287617 \times 10^{-11}$	2.093892×10^{-11}	$-5.193726 \times 10^{-11}$

G Quadrature of $\mathcal{R}$ versus $\widehat{\mathcal{R}}_{24}$

Table 5 compares the continuous integral with the historical 24-node trapezoidal estimator on every headline geometry. Path orderings agree. Trapezoidal values at 24, 48, 96, and 192 nodes are recorded in `paper/arxiv/artifacts/geometries.json`. Adaptive quadrature is used for multi-mode $\mathcal{R}$; the scalar integrals above are closed form. Scipy error estimates are not interval certificates.

Table 5: Continuous $\mathcal{R}$ versus $\widehat{\mathcal{R}}_{24}$ on headline geometries, including the non-centered anisotropic family.

geometry	$\mathcal{R}_{\text{lin}}$	$\widehat{\mathcal{R}}_{24}^{\text{lin}}$	$\mathcal{R}_{\text{VP}}$	$\widehat{\mathcal{R}}_{24}^{\text{VP}}$
scalar $\lambda = 4$	0.963495	0.964206	0.616850	0.616850
anisotropic $d = 2, 8$	0.972296	0.972637	0.192270	0.192361
low-rank $d = 2$	1.245531	1.244898	0.625461	0.625454
low-rank $d = 8$	2.944104	2.947652	4.730544	4.729521
non-centered anisotropic $d = 2, 8$	1.093537	1.093851	0.342354	0.342536

H Low-rank robustness grid

The post-hoc low-rank grid has 66 preference blocks: 2 dimensions, 3 solvers, and, for each pair, the three primary NFE budgets $\{8, 16, 32\}$ at noise-floor perturbations $\delta \in \{-0.1, 0, 0.1\}$, plus NFE $\{64, 128\}$ at $\delta = 0$ only. The perturbation multiplies the noise variance 0.05 by $(1 + \delta)$; the factor F is regenerated from seed 271828. Extra budgets are not run at $\delta \neq 0$. Table 6 summarizes the 66 blocks; the full list is `paper/arxiv/artifacts/robustness_lowrank.json`.

Table 6: Post-hoc low-rank robustness: all 66 blocks follow the same solver-path split as the primary grid.

solver	$d = 2$ blocks	$d = 8$ blocks	preferred path
euler	11	11	VP
heun	11	11	linear
rk4	11	11	linear

I High-precision and interval procedures

Scalar linear Heun factors are computed in exact rationals. The VP Heun product is certified by substituting $\pi < 355/113$ and $\sqrt{2} < 99/70$ into a nonnegative element of $\mathbb{Q}[\pi, \sqrt{2}]$; the ranking inequality is the integer comparison in Appendix A. An independent `mpmath.iv` enclosure at 40 decimal digits (`mpmath` 1.3.0) and an 80-digit (non-interval) `mpmath` product are recorded as cross-checks against the package float64 Heun implementation. Frozen Phase 4 Gaussian W_2 values were checked against an 80-digit Gelbrich reference in `outputs/phase4_precision_2026-07-24-v1/`. Default numerical precision for the benchmark runs is float64.

J Three-path comparison

Table 7 reports linear, VP, and per-mode log-covariance on every centered solver-budget block. Log-covariance has the smallest $\mathcal{R}$ and the smallest W_2 in all 36 blocks. This is an observed count on the tested grid, not a population claim.

K Low-rank seed fraction

Fifty independent draws of the rank-2 factor F (seeds 10000, ..., 10049), at $d \in \{2, 8\}$, each tested for at least one linear-versus-VP inversion among the nine solver-budget conditions. Unlike the post-hoc 66-block robustness grid, these draws redraw F . Both dimensions invert in 50 of 50 draws. The Wald interval at $\hat{p} = 1$ is degenerate.

Table 7: Per-mode log-covariance versus linear and VP on the centered grid.

family	d	solver	NFE	R min	W_2 min	$W_{2,\log}$	$W_{2,\text{lin}}$
anisotropic	2	euler	8	log-cov	log-cov	0.01164	0.2441
anisotropic	2	euler	16	log-cov	log-cov	0.005874	0.1266
anisotropic	2	euler	32	log-cov	log-cov	0.002952	0.06454
anisotropic	2	heun	8	log-cov	log-cov	0.0006613	0.03243
anisotropic	2	heun	16	log-cov	log-cov	0.0001682	0.006327
anisotropic	2	heun	32	log-cov	log-cov	4.244e-05	0.001359
anisotropic	2	rk4	8	log-cov	log-cov	3.834e-06	0.01119
anisotropic	2	rk4	16	log-cov	log-cov	2.473e-07	0.000213
anisotropic	2	rk4	32	log-cov	log-cov	1.575e-08	9.781e-06
anisotropic	8	euler	8	log-cov	log-cov	0.0131	0.4509
anisotropic	8	euler	16	log-cov	log-cov	0.006608	0.2337
anisotropic	8	euler	32	log-cov	log-cov	0.003319	0.1191
anisotropic	8	heun	8	log-cov	log-cov	0.0007027	0.04645
anisotropic	8	heun	16	log-cov	log-cov	0.0001786	0.008457
anisotropic	8	heun	32	log-cov	log-cov	4.503e-05	0.001746
anisotropic	8	rk4	8	log-cov	log-cov	3.897e-06	0.01812
anisotropic	8	rk4	16	log-cov	log-cov	2.513e-07	0.0003324
anisotropic	8	rk4	32	log-cov	log-cov	1.6e-08	1.37e-05
low-rank	2	euler	8	log-cov	log-cov	0.03303	0.2933
low-rank	2	euler	16	log-cov	log-cov	0.01683	0.1523
low-rank	2	euler	32	log-cov	log-cov	0.008497	0.07773
low-rank	2	heun	8	log-cov	log-cov	0.002998	0.0585
low-rank	2	heun	16	log-cov	log-cov	0.0007696	0.0122
low-rank	2	heun	32	log-cov	log-cov	0.0001953	0.002707
low-rank	2	rk4	8	log-cov	log-cov	4.7e-05	0.004578
low-rank	2	rk4	16	log-cov	log-cov	2.974e-06	0.000432
low-rank	2	rk4	32	log-cov	log-cov	1.89e-07	1.732e-05
low-rank	8	euler	8	log-cov	log-cov	0.2736	0.8109
low-rank	8	euler	16	log-cov	log-cov	0.1445	0.424
low-rank	8	euler	32	log-cov	log-cov	0.07439	0.2175
low-rank	8	heun	8	log-cov	log-cov	0.05274	0.4094
low-rank	8	heun	16	log-cov	log-cov	0.01399	0.08936
low-rank	8	heun	32	log-cov	log-cov	0.00363	0.02047
low-rank	8	rk4	8	log-cov	log-cov	0.004721	0.06962
low-rank	8	rk4	16	log-cov	log-cov	0.0002675	0.00746
low-rank	8	rk4	32	log-cov	log-cov	1.687e-05	0.0001322

Table 8: Fraction of low-rank geometries with at least one inversion, 50 independent draws of F .

d	inversions / 50	fraction	Wald 95%
2	50/50	1.00	[1.00, 1.00]
8	50/50	1.00	[1.00, 1.00]