

Few-Step Flow-Matching Error Can Be Misranked by Averaged Jacobian Regularity: A Certified Gaussian Counterexample

Clément Callaert
CentraleSupélec and Université Paris-Saclay
callaert.clement@gmail.com

Abstract

Does averaged squared Jacobian regularity rank two interpolants in the same order as equal-NFE endpoint error? Chen, Vanden-Eijnden, and Xu propose minimizing A_2 , the time-integrated squared spatial Jacobian norm of a flow-matching marginal ODE, as a criterion for schedule design, without a proved discretization-error bound. Classical one-step bounds use a Lipschitz constant of the field; A_2 is not that constant. Already for independent $N(0, 1)$ and $N(0, 4)$, the exact regularity integrals of the linear and trigonometric variance-preserving Gaussian interpolants are $5\pi/8 - 1$ and $\pi^2/16$, while Heun at NFE 8 reverses Gaussian Wasserstein-2 distance. The linear Heun product is the rational $6797469/3559400$; a nonnegative element of $\mathbb{Q}[\pi, \sqrt{2}]$ yields $r_{\text{VP}} < 187/100$. The object is a flow matching marginal interpolant ODE, not a score-based probability-flow ODE. The Gaussian drift and W_2 formula are closed form; the regularity integrand is exact, while multimode $\mathcal{R}$ is evaluated deterministically by adaptive quadrature. Pairwise and four-path census comparisons on commuting Gaussian interpolants are reported below; a finite census does not imply that a global A_2 -minimizer minimizes fixed-NFE error. The analysis uses stochastic interpolants, few-step sampling, Runge–Kutta, and interpolation schedules; no learned field is used.

1 Introduction

A few-step sampler engages a probability path, a numerical solver, and a budget of number of function evaluations (NFE) before any endpoint error is observed. Few-step sampling therefore makes the sampling schedule a design choice that must be made a priori. Classical one-step bounds involve a Lipschitz constant of the field [Hairer et al., 1993]. What the practitioner needs is a scalar criterion for schedule design that can be evaluated from the path alone.

In the stochastic interpolants and flow matching setting, Chen et al. [2025] propose minimizing averaged squared Jacobian regularity (their Definition 3.2, denoted A_2 , the time-integrated squared spatial Jacobian norm) as a selection criterion. The integrand is a Jacobian norm. They prove KL equivalence of schedules and a logarithmic scaling law. They prove no bound relating A_2 to discretization error, Wasserstein-2 distance, or sampling error. This paper asks whether the

pairwise order that A_2 induces between two interpolant paths is reliable at equal NFE on a flow-matching marginal ODE.

Commuting Gaussian interpolants are the right probe. The Gaussian drift and W_2 formula are closed form; the regularity integrand is exact, while multimode $\mathcal{R}$ is evaluated deterministically by adaptive quadrature. Any observed divergence is therefore a property of the criterion rather than an estimation artefact. The family includes the linear interpolant and the trigonometric variance-preserving (VP) path, both C^1 shared schedules; the Chen scalar log-covariance schedule, a shared schedule whose square-root coefficients are not C^1 at the endpoints; and the per-mode log-covariance path.

The leading logical result is a deterministic scalar example, proved by an exact integral, an exact rational product of the Heun method, and a rational enclosure of a two-stage Runge–Kutta product (Section 3.2). The same calculation then splits into three comparisons, summarized in Figure 1: $\mathcal{R}$ as a pairwise comparator of two fixed Class S paths, $\mathcal{R}$ as a pairwise comparison of two shared-schedule Class Q paths, and a four-path census that includes the per-mode Class Q path.

The contributions are as follows.

- A certified one-dimensional counterexample: exact $\mathcal{R}$ prefers trigonometric VP while Heun at NFE 8 prefers the linear path in Gaussian W_2 (Section 3.2, Appendix A). An exact integral or an exact Heun product that restored the $\mathcal{R}$ ranking would contradict the claim.
- An exact account of the information discarded by the unsigned average: leading exact-minus-method coefficients, stage locations, and a forced sign change of the linear drift (Section 3.3). A truncation expansion in which those coefficients reduced to $\|A\|_2^2$ would contradict the account.
- A three-comparison characterisation on a complete $4 \times 3 \times 3$ census: pairwise linear-versus-VP inversions in 5 of 12 cells; shared-schedule pairwise VP-versus-scalar-log-covariance inversions in 9 of 36 blocks; among four candidate paths, the per-mode path has the smallest $\mathcal{R}$ and the smallest W_2 in 36 of 36 blocks. This finite-census observation does not imply that a global $\mathcal{R}$ -minimizer minimizes fixed-NFE error. When Σ_1 has at least two distinct eigenvalues, that path is not a Class S shared interpolant. Rank agreement between $\mathcal{R}$ and W_2 changes sign with the solver (Sections 4.3–4.5). A census in which all three comparisons agreed would contradict the split.
- An inversion map over target variance $\lambda \in [0.05, 100]$, reported as log-Lebesgue measure rather than a grid-point count (Section 4.6). Emptiness of the inversion set on that interval would contradict the map.

2 Background

2.1 Gaussian interpolants and the marginal velocity field

Let $X_0 \sim N(\mu_0, \Sigma_0)$ and $X_1 \sim N(\mu_1, \Sigma_1)$ be independent in $\mathbb{R}^d$, with Σ_0, Σ_1 symmetric positive definite. A *shared schedule* is a pair of scalar functions used




Figure 1: Three comparisons for $\mathcal{R}$ on commuting Gaussian interpolants. Each row shows continuous $\mathcal{R}$ (left) and equal-NFE Gaussian W_2 (right; lower is preferred). (a) Pairwise linear versus VP on low-rank $d = 8$, Euler, NFE 8: $\mathcal{R}$ and W_2 invert; this occurs in 5 of 12 geometry×solver cells. (b) Shared-schedule pairwise comparison, VP versus Chen Ex. 3.3 scalar ($M = \lambda_{\max}$) on anisotropic $d = 8$, Euler, NFE 8: inversion, as in 9 of 36 blocks. (c) Four candidate paths on low-rank $d = 8$, Euler, NFE 8, including per-mode $q_i(t) = \lambda_i^t$: smallest $\mathcal{R}$ and smallest W_2 in 36 of 36 blocks on this finite census. This does not imply that a global $\mathcal{R}$ -minimizer minimizes fixed-NFE error. When Σ_1 has at least two distinct eigenvalues, the per-mode path is not a Class S shared interpolant. Exact-moment modal Runge–Kutta; census counts, not estimators.

for every eigenmode, equivalently $Q(t) = \alpha(t)^2 \Sigma_0 + \sigma(t)^2 \Sigma_1$. We distinguish three regularity classes.

- *Class S* (shared C^1 interpolants): $\alpha, \sigma \in C^1([0, 1])$ with

$$\alpha(0) = 1, \quad \sigma(0) = 0, \quad \alpha(1) = 0, \quad \sigma(1) = 1. \quad (1)$$

The linear path $(1 - t, t)$ and trigonometric VP $(\cos(\pi t/2), \sin(\pi t/2))$ belong to Class S, as does the grid-aware embedding of Appendix C.

- *Class Q* (regular covariance paths): a C^1 path of positive definite covariances $Q(t)$ with $Q(0) = \Sigma_0$, $Q(1) = \Sigma_1$, for which $A(t) = \frac{1}{2}Q'(t)Q(t)^{-1}$ extends continuously to $[0, 1]$. Every Class S interpolant induces a Class Q path. The per-mode schedule $q_i(t) = \lambda_i^t$ is Class Q.

- *Class R* (regular squares): $\alpha^2, \sigma^2 \in C^1([0, 1])$, $\alpha^2 \geq 0$, $\sigma^2 \geq 0$, and

$$\alpha(0)^2 = 1, \quad \sigma(0)^2 = 0, \quad \alpha(1)^2 = 0, \quad \sigma(1)^2 = 1. \quad (2)$$

The square roots $\alpha = \sqrt{\alpha^2}$ and $\sigma = \sqrt{\sigma^2}$ need not be differentiable at the endpoints. Chen et al.'s scalar log-covariance schedule $\sigma(t)^2 = (M^t - 1)/(M - 1)$, $\alpha(t)^2 = (M - M^t)/(M - 1)$ for $M > 0$, $M \neq 1$ is Class R: $\sigma(t) \asymp \sqrt{t}$ as $t \rightarrow 0$ and $\alpha(t) \asymp \sqrt{1-t}$ as $t \rightarrow 1$, so it is not Class S. When $M = 1$ the formulas are read in the limit $\sigma(t)^2 = t$, $\alpha(t)^2 = 1 - t$. The covariance is C^∞ and A is continuous, so the path is Class Q.

Two paths are called shared-schedule Class Q paths when both lie in Class Q and both use a single scalar pair (α, σ) . We do not call a Class R path a Class S interpolant.

The interpolant of a Class S or Class R schedule is

$$X_t = \alpha(t)X_0 + \sigma(t)X_1, \quad (3)$$

as in stochastic interpolants and flow matching [Albergo et al., 2025, Lipman et al., 2023, Liu et al., 2023, Tong et al., 2024]. Throughout, $\alpha^2 + \sigma^2$ is not required to equal 1 except on the VP path and on Chen et al.'s scalar log-covariance schedule. The two classical paths compared below are the linear schedule $(\alpha, \sigma) = (1 - t, t)$ and the trigonometric VP schedule $(\cos(\pi t/2), \sin(\pi t/2))$.

The law of X_t is Gaussian with mean $m_t = \alpha\mu_0 + \sigma\mu_1$ and covariance $Q(t) = \alpha^2\Sigma_0 + \sigma^2\Sigma_1$. We assume $Q(t) \succ 0$ on $[0, 1]$. The marginal velocity of (3) is the affine field

$$b(t, x) = A(t)x + c(t), \quad A(t) = C_t Q(t)^{-1}, \quad c(t) = \dot{m}_t - A(t)m_t, \quad (4)$$

with $C_t = \dot{\alpha}\alpha\Sigma_0 + \dot{\sigma}\sigma\Sigma_1$ on $(0, 1)$ for Class S and Class R. Equivalently, $C_t = \frac{1}{2}(\alpha^2)'\Sigma_0 + \frac{1}{2}(\sigma^2)'\Sigma_1$, which extends continuously to $[0, 1]$ whenever $\alpha^2, \sigma^2 \in C^1([0, 1])$. This is a marginal interpolant ODE, not a score-based probability-flow ODE in the sense of Song et al. [2021], although the latter is an instance of a generative ODE. If $\mu_0 = \mu_1 = 0$ and $\Sigma_0 = I$, then $c \equiv 0$ and $A = \frac{1}{2}Q'Q^{-1}$.

If $\Sigma_0 = I$, then $Q(t)$, $Q'(t)$, $A(t)$, and Σ_1 share the eigenbasis of Σ_1 . Simultaneous diagonalization with the identity is not an extra hypothesis. In that basis, mode i with eigenvalue $\lambda_i > 0$ obeys $x'_i = a_i(t)x_i + c_i(t)$ with $q_i(t) = \alpha(t)^2 + \lambda_i\sigma(t)^2$ and $a_i(t) = q'_i(t)/(2q_i(t))$. For a Class Q path that is not shared, the same identity holds modewise with no single pair (α, σ) . The covariance drift depends only on the variance schedule. The exact one-step factor on $[t_n, t_{n+1}]$ is $\sqrt{q_i(t_{n+1})/q_i(t_n)}$. For centered modes the exact endpoint factor is $\sqrt{\lambda_i}$.

2.2 Solvers, evaluation budget, and endpoint error

Euler uses $s = 1$ field evaluation per step, Heun $s = 2$, and classical RK4 $s = 4$. A budget B is admitted only if $B \equiv 0 \pmod{s}$. The step count is $S = B/s$ and the grid is the uniform partition of $[0, 1]$ with S steps. Equal NFE is not equal step count: at $B = 8$, Euler takes 8 steps, Heun 4, and RK4 2.

A solver applied to (4) produces an affine endpoint map $x \mapsto Rx + d$. Source moments are pushed through this map. Error is the Gaussian W_2 of the endpoint laws [Gelbrich, 1990, Peyré and Cuturi, 2019],

$$W_2^2 = \|\hat{m} - \mu_1\|_2^2 + \text{tr}(\hat{C} + \Sigma_1 - 2(\Sigma_1^{1/2}\hat{C}\Sigma_1^{1/2})^{1/2}). \quad (5)$$

This is neither pathwise trajectory error nor local truncation error. In the centered commuting case, $W_2^2 = \sum_i (|r_i| - \sqrt{\lambda_i})^2$.

2.3 The averaged squared Jacobian criterion

Chen et al. [2025, Definition 3.2] introduce

$$A_2 = \int_0^1 \mathbb{E}[\|\nabla b_t(I_t)\|_2^2] dt. \quad (6)$$

For the affine field (4) the integrand is deterministic and

$$\mathcal{R}[b] = \int_0^1 \|A(t)\|_2^2 dt \quad (7)$$

coincides with A_2 . Thus A_2 is the time-integrated squared spatial Jacobian norm, not a classical Lipschitz constant of the field. By definition $\mathcal{R}$ ignores $c(t)$. For commuting $A(t)$, $\|A(t)\|_2 = \max_i |a_i(t)|$. Headline comparisons use the continuous integral (7). The historical 24-node trapezoidal estimator is written $\hat{\mathcal{R}}_{24}$ and is used only as an implementation check. No trapezoidal value is described as an exact integral.

What the criterion does and does not establish. Chen et al. [2025] propose A_2 as a design objective. They prove that schedules related by a time-change have equivalent KL cost, and that the optimized one-dimensional log-covariance schedule attains a logarithmic scaling of A_2 . They prove no bound relating A_2 to a discretization error, a Wasserstein error, or a sampling error. They never compare the linear interpolant to trigonometric VP.

3 Method

3.1 The variance-path invariant and the Cauchy–Schwarz bound

Proposition 1. *For any Class Q one-dimensional variance path from 1 to $\lambda > 0$, $\int_0^1 a(t) dt = \frac{1}{2} \ln \lambda$. Consequently*

$$\mathcal{R} = \int_0^1 a(t)^2 dt = \left(\int_0^1 a(t) dt \right)^2 + \text{Var}_t(a) \geq \left(\frac{1}{2} \ln \lambda \right)^2, \quad (8)$$

with equality if and only if a is constant.

Proof. The integral identity is $\frac{1}{2} \log q(1) - \frac{1}{2} \log q(0)$. Expanding $\int a^2$ as $(\int a)^2 + \text{Var}_t(a)$ gives the bound. $\square$

Thus $\mathcal{R}$ measures only the temporal variance of the log-covariance rate. Its unconstrained Class Q minimizer is the constant drift $a \equiv \frac{1}{2} \ln \lambda$, i.e. $q(t) = \lambda^t$, which is Chen et al. [2025, Example 3.3] in one dimension. In several dimensions the per-mode schedule $q_i(t) = \lambda_i^t$ attains $\mathcal{R} = (\frac{1}{2} \max_i |\ln \lambda_i|)^2$. If Σ_1 has at least two distinct eigenvalues, that path is not a Class S shared interpolant: matching $\alpha^2 + \lambda_i \sigma^2 = \lambda_i^t$ on two distinct eigenvalues forces $\sigma(t) \asymp \sqrt{t}$ as $t \rightarrow 0$. If three or more distinct eigenvalues are present, no shared (α, σ) exists even in Class R, because $\alpha^2 + \lambda \sigma^2$ is affine in λ while λ^t is not. Chen et al. [2025, Proposition 3.9] optimize A_2 over shared scalar schedules under the spectral hypothesis $1 \geq \lambda^{(1)} \geq \dots \geq \lambda^{(d)} > 0$, taking M equal to the smallest eigenvalue. The geometries below have mixed spectra with some $\lambda_i > 1$; the scalar candidate used in the census is Chen Example 3.3 with $M = \lambda_{\max}$. That is a pairwise comparison of two shared-schedule Class Q paths, not a test of Proposition 3.9.

3.2 A certified one-dimensional counterexample

Assumption 1. *The endpoints $X_0 \sim N(0, 1)$ and $X_1 \sim N(0, 4)$ are independent. The compared schedules are linear and trigonometric VP. The solver is explicit Heun with NFE $B = 8$, hence $S = 4$ steps of size $h = 1/4$.*

Under Assumption 1, $q(t) = \alpha(t)^2 + 4\sigma(t)^2$ and $a(t) = q'(t)/(2q(t))$. The exact endpoint factor is 2. All Heun factors below are positive, so $W_2 = ||r| - 2| = |r - 2|$.

Proposition 2. *Under Assumption 1,*

$$\mathcal{R}_{\text{lin}} = \frac{5\pi}{8} - 1, \quad \mathcal{R}_{\text{VP}} = \frac{\pi^2}{16}, \quad (9)$$

so averaged regularity prefers VP. The Heun endpoint factors satisfy

$$r_{\text{lin}} = \frac{6797469}{3559400}, \quad 0 < r_{\text{VP}} < \frac{187}{100}, \quad (10)$$

the second bound being a rational enclosure obtained by substituting $\sqrt{2} < 99/70$ and $\pi < 355/113$ into the exact element of $\mathbb{Q}[\pi, \sqrt{2}]$. Consequently

$$W_2^{\text{lin}} = \frac{321331}{3559400} < 0.091 < 0.130 < W_2^{\text{VP}}, \quad (11)$$

so equal-NFE Gaussian W_2 prefers the linear path. The regularity ordering does not determine the fixed-NFE W_2 ordering.

Proof sketch. The linear drift is $a(t) = (5t - 1)/(1 - 2t + 5t^2)$. Completing the square to $5(t - 1/5)^2 + 4/5$ and using $\arctan 2 + \arctan(1/2) = \pi/2$ yields $\mathcal{R}_{\text{lin}} = 5\pi/8 - 1$. The VP integral evaluates to $\pi^2/16$. The four linear Heun factors are the rationals $47/52$, $6/5$, $497/370$, and $97/74$, whose product is $6797469/3559400$. Then $321331/3559400 < 91/1000$ is an elementary comparison of integers. The VP grid drifts are exact elements of $\mathbb{Q}[\pi, \sqrt{2}]$; the four-step Heun product has only nonnegative coefficients, so substituting the rational upper bounds $\sqrt{2} < 99/70$ and $\pi < 355/113$ is valid and gives $r_{\text{VP}} < 187/100$, hence $W_2^{\text{VP}} > 13/100$. Details occupy Appendix A. $\square$

One certified inversion proves the logical non-implication. The remainder of the paper identifies what $\mathcal{R}$ cannot retain, then measures how the failure depends on the regime in which $\mathcal{R}$ is used.




Figure 2: Scalar drifts $a_{\text{lin}}(t)$ and $a_{\text{VP}}(t)$ for independent $N(0, 1)$ and $N(0, 4)$. The shaded region is $a_{\text{lin}} < 0$. Markers are the four Heun nodes at NFE 8. $a_{\text{lin}}(0) = -1 < 0 < a_{\text{lin}}(1) = +1$; the VP drift does not change sign on $(0, 1)$. Exact evaluation of the closed-form drifts.

3.3 What the unsigned average discards

The content of a ranking inversion is the leading exact-minus-method coefficient of the one-step factor (Appendix B): $\frac{1}{2}(a' + a^2)h^2$ for Euler and $(-a''/12 + a^3/6)h^3$ for the implemented Heun factor $1 + (h/2)(a(t) + a(t+h)(1 + ha(t)))$. These involve different derivatives, signs, and stage locations. Averaging $\|A\|_2^2$ retains none of them.

The telescoping identity

$$\prod_{n=0}^{S-1} r_{i,n} - \prod_{n=0}^{S-1} e_{i,n} = \sum_{n=0}^{S-1} (r_{i,n} - e_{i,n}) \left(\prod_{j < n} r_{i,j} \right) \left(\prod_{j > n} e_{i,j} \right) \quad (12)$$

is the accounting tool that converts those local coefficients into endpoint factor error. Here $e_{i,n} = \sqrt{q_i(t_{n+1})/q_i(t_n)}$ is exact and $r_{i,n}$ is numerical. Prefixes are numerical; suffixes are exact. The identity holds without a non-vanishing hypothesis. Appendix D illustrates the signed transported terms on the strongest centered block.

Fixed-stage Runge–Kutta methods use signed stage evaluations that $\mathcal{R}$ discards; in the tested schemes, this distinction contributes to the observed inversion. For any independent linear coupling, $a_{\text{lin}}(0) = -1 < 0 < a_{\text{lin}}(1) = +1$. The VP drift vanishes at the endpoints and does not change sign on $(0, 1)$.

3.4 Three regimes: comparator, shared-schedule pairwise comparison, four-path census

Proposition 2 is a pairwise comparison of two fixed Class S paths. Comparing two shared-schedule Class Q paths is a different question, and enumerating four candidate paths including the per-mode Class Q path is a third. The rest of the paper answers these three questions on one factorial grid.

1. $\mathcal{R}$ as a *pairwise comparator* between two Class S paths (linear versus trigonometric VP).

Table 1: Experimental grid. Headline $\mathcal{R}$ is the continuous integral. Ranking inversions were identified in the registered Phase 3 gate. Phase 4 is a confirmatory reproduction on configurations frozen before the rerun; it is not a preregistration that predates discovery. Status “confirmatory” means reproduced on that frozen grid; “post-hoc” means added after the Phase 3 observation. Reserve “pre-registered” for a protocol recorded before any observation. The Gaussian drift and W_2 formula are closed form; the regularity integrand is exact, while multimode $\mathcal{R}$ is evaluated deterministically by adaptive quadrature.

experiment	grid	paths	status
pairwise comparator	4 geom. $\times$ 3 solvers $\times$ {8, 16, 32}	linear, VP	confirmatory
shared-schedule pairwise	same 36 blocks	VP, Ex. 3.3 scalar	post-hoc
four-path census	same 36 blocks	$+ q_i = \lambda_i^t$	post-hoc
inversion map	$\lambda \in [0.05, 100] \times 3 \times 3$	linear, VP	post-hoc
geometry draws	$N = 50$ i.i.d. $F, d \in \{2, 8\}$	linear, VP	post-hoc
robustness $\pm 10\%$, NFE 64, 128	66 low-rank blocks	linear, VP	post-hoc
non-centered stress test	$2 \times 3 \times 3$	linear, VP	confirmatory

2. $\mathcal{R}$ as a *pairwise comparison within the shared-schedule class*: trigonometric VP (Class S, hence Class Q) versus the scalar Chen Example 3.3 schedule with $M = \lambda_{\max}$ (Class R and Class Q, not Class S).
3. A *four-path census* that additionally includes the per-mode schedule $q_i(t) = \lambda_i^t$ (Class Q; not Class S when Σ_1 has at least two distinct eigenvalues).

A complementary scalar construction (Appendix C) exhibits an admissible Class S $\mathcal{R}$ -minimizer for which left-endpoint Euler has strictly larger endpoint error than a higher- $\mathcal{R}$ Class S path. That field is chosen after the grid. It is not the mechanism of Proposition 2 or of the Gaussian census. A 36 of 36 count on four candidate paths does not imply that a global $\mathcal{R}$ -minimizer minimizes fixed-NFE error; this construction blocks that generalization.

4 Experiments

4.1 Setup

Source $N(0, I)$. Four centered targets, with seed 271828 for low-rank factors: anisotropic Gaussians with condition number 4 in dimensions 2 and 8 (eigenvalues a geometric sequence from $1/2$ to 2); and rank-2 covariances $FF^\top + 0.05I$. The dimension-2 low-rank eigenvalues are 0.291 and 2.97; the dimension-8 low-rank eigenvalues are 0.05 (six times), 4.65, and 10.7 (three decimals in the body; full float64 values in Appendix D). Paths, solvers, budgets, and registration status are in Table 1. Precision is float64, audited against an 80-digit Gelbrich reference.

4.2 Statistical treatment and what the counts mean

Given a geometry, a path, a solver, and a budget, $\mathcal{R}$ and W_2 are deterministic numerical quantities with no Monte Carlo sampling noise; float64 results are checked against high-precision references. The 36-block grid is a complete enumeration of 4 geometries $\times$ 3 solvers $\times$ 3 budgets, not a draw. Because $\mathcal{R}$ is independent of solver and budget, and because the two anisotropic geometries share one $\mathcal{R}$ pair,

the census contains 3 distinct linear-versus-VP $\mathcal{R}$ comparisons and nests into 12 geometry $\times$ solver cells (Appendix F). Each inversion is reported by the ratio of its W_2 margin to the 80-digit floor; the pre-declared threshold is $\geq 10^3$, and every census inversion lies above it. The only random ingredient, and the only confidence interval in the paper, is the draw of the low-rank factor F (Section 4.7).

4.3 R as a pairwise comparator: linear versus VP

Does $\mathcal{R}[b] < \mathcal{R}[b']$ imply $E[b] < E[b']$ at equal NFE for the linear and trigonometric VP interpolants? The protocol is the confirmatory Phase 4 36-block census of Table 1, with continuous $\mathcal{R}$, on a grid frozen before the rerun. Linear versus VP invert in 5 of 12 geometry $\times$ solver cells; 4 of those cells invert at every NFE in $\{8, 16, 32\}$, from 3 distinct $\mathcal{R}$ comparisons. On this census the paired concordance score $(n_{\text{agree}} - n_{\text{invert}})/n$ between the $\mathcal{R}$ order and the W_2 order equals $2/9$ over 36 blocks. Stratified by solver, with $n = 4$ geometries and a structural tie between the two anisotropic targets, the score is $+1/2$ under Euler and RK4 and $-1/3$ under Heun: rank agreement changes sign with the solver. The strongest block is low-rank, $d = 8$, Euler, NFE 8: $\mathcal{R}$ prefers linear (2.944 versus 4.731) while W_2 prefers VP (0.811 versus 0.456). Appendix D lists every inverted row. This census does not estimate a population frequency, and it does not speak to learned fields.

4.4 R as a pairwise comparison within the shared-schedule class: VP versus the scalar log-covariance schedule

When both paths are shared-schedule Class Q paths, does the one with smaller $\mathcal{R}$ have the smaller W_2 ? The protocol compares trigonometric VP to Chen Example 3.3 with a single ratio $M = \lambda_{\max}$ on the same 36 blocks (post-hoc). This is a pairwise comparison of two candidates, not a global minimization of $\mathcal{R}$ over the shared-schedule class, and not a test of Chen et al. [2025, Proposition 3.9]. The ranking inverts in 9 of 36 blocks and 4 of 12 cells: anisotropic $d = 2$, Euler at NFE 16 and 32; anisotropic $d = 8$, Euler at NFE 8, 16, and 32; low-rank $d = 2$, Euler at NFE 8 and Heun at NFE 8, 16, and 32; low-rank $d = 8$, none. Figure 3 places the two shared-schedule paths next to linear and the per-mode path. Every pairwise inversion margin exceeds the 80-digit floor by a factor greater than 10^3 . The comparison does not say that A_2 fails as a global optimization objective on the shared-schedule class.

4.5 Four-path census: the per-mode log-covariance path

If the per-mode schedule $q_i(t) = \lambda_i^t$ is included among the four candidate paths, does it attain both the smallest $\mathcal{R}$ and the smallest W_2 ? Among the four candidate paths on the stated 36-block census, the per-mode log-covariance path has the smallest continuous $\mathcal{R}$ and the smallest Gaussian W_2 in 36 of 36 blocks. This finite-census observation does not imply that a global $\mathcal{R}$ -minimizer minimizes fixed-NFE error. When Σ_1 has at least two distinct eigenvalues, that path is not a Class S shared interpolant. The observation is a census count on four candidates, not a sampling law, and it does not supply a practical multi-dimensional schedule.




Figure 3: Four paths on the four centered geometries: linear, trigonometric VP, shared Chen Example 3.3 scalar schedule with $M = \lambda_{\max}$, and the per-mode log-covariance schedule. Left: continuous $\mathcal{R}$. Right: Euler Gaussian W_2 at NFE 8. Among these four candidate paths, the per-mode schedule has the smallest $\mathcal{R}$ and the smallest W_2 in 36 of 36 blocks on the stated census. This finite-census observation does not imply that a global $\mathcal{R}$ -minimizer minimizes fixed-NFE error. When Σ_1 has at least two distinct eigenvalues, the per-mode path is not a Class S shared interpolant. Exact-moment modal Runge–Kutta; $\mathcal{R}$ by adaptive quadrature except the per-mode closed form.

4.6 Where the inversions live: a sweep over target variance

Are linear-versus-VP inversions an artefact of $\lambda = 4$? The protocol is a log-uniform sweep $\lambda \in [0.05, 100]$ at 1281 nodes, for each of Euler, Heun, and RK4 and each NFE in $\{8, 16, 32\}$. The reported quantity is the Lebesgue measure in $\log \lambda$ of the inversion set, not a count of inverted nodes. At Heun NFE 8 that measure is 3.337 on an interval of log-length 7.601, hence a fraction 0.439. Doubling the resolution to 2561 nodes changes every cell’s measure by less than 1% (maximum relative change 0.0020). Figure 4 displays the set. At $\lambda = 4$ the inversion is present under Heun at every tested budget $\text{NFE} \in \{4, 8, 12, 16, 24, 32, 64, 128, 256\}$, absent under Euler, and mixed under RK4 (absent at NFE 8 and 12). Euler does invert at more extreme λ . The scalar example is solver-dependent, not an artefact of Heun at budget 8 alone. The sweep does not yield a population law over target covariances.

4.7 Robustness to the random geometry

Does the low-rank inversion survive a redraw of F ? The protocol draws 50 i.i.d. rank-2 factors from `torch.Generator` with seeds 10000, ..., 10049, at $d \in \{2, 8\}$, and records whether each geometry has at least one linear-versus-VP inversion among the nine solver-budget conditions. Unlike the 66-block robustness grid of Appendix G, these draws redraw F . Both dimensions invert in 50 of 50 draws. The 95% Clopper–Pearson interval is $[0.929, 1.000]$ (Table 2). This is the only confidence interval in the paper. It is not a statement that inversion occurs for every low-rank geometry.




Figure 4: Linear versus VP ranking inversion versus target variance λ , for Euler, Heun, and RK4 at NFE 8, 16, and 32. Dark cells: $\mathcal{R}$ and W_2 disagree. The dashed line is $\lambda = 4$ (Proposition 2). The inversion set is reported in the text by its Lebesgue measure in $\log \lambda$ on $[0.05, 100]$, not by a count of dark pixels. Numerical evaluation of a deterministic ranking.

Table 2: Fraction of low-rank geometries with at least one linear-versus-VP inversion, from 50 i.i.d. draws of the rank-2 factor F (seeds 10000 through 10049; `torch.Generator`). The interval is exact Clopper–Pearson 95%. The existing $\pm 10\%$ robustness grid does not redraw F . Inferential interval on a binomial geometry draw; not an interval on $\mathcal{R}$ or W_2 .

d	N	inversions	$\hat{p}$	Clopper–Pearson 95%
2	50	50/50	1.00	[0.929, 1.000]
8	50	50/50	1.00	[0.929, 1.000]

4.8 Numerical validity controls

Adaptive quadrature of $\max_i a_i(t)^2$ agrees in sign with $\hat{\mathcal{R}}_{24}$ on every headline geometry, including the scalar pair of Proposition 2. Trapezoidal estimates at 24, 48, 96, and 192 nodes converge to the same ordering (Appendix F). All 72 float64 W_2 values on the centered census agree with an 80-digit reference to at most 9.7×10^{-10} . Interval procedures for the VP Heun product occupy Appendix F.

A confirmatory non-centered family (Appendix E) shifts the means so that $c(t) \neq 0$. Then $\mathcal{R}$, which ignores c , is blind to a mean contribution in W_2^2 . Of 18 blocks, 11 invert under continuous $\mathcal{R}$, and several inversions including all dimension-8 Euler and Heun inversions are mean-driven. The family is a structural stress test of that blind spot, not an independent replication of the centered mechanism.

Mixture targets are excluded. Dimension-8 mixture endpoint estimators failed a post-result calibration diagnostic; the census accepts Gaussian targets only.

5 Related work

Flow matching and stochastic interpolants supply the Gaussian path family [Lipman et al., 2023, Albergo et al., 2025, Liu et al., 2023, Tong et al., 2024]. Lipschitz-guided schedule design proposes the selection criterion tested here [Chen et al., 2025]. Solver-specific schedules and exact linear parts are documented in diffusion

practice [Sabour et al., 2024, Karras et al., 2022, Lu et al., 2022a,b, Zhang and Chen, 2023].

Stéphanovitch [2026] prove sharp Lipschitz regularity of canonical flow-matching fields and Euler-type W_2 rates of order $\sqrt{d}/N$ up to logs. That is a bound, not a pairwise ranking of two interpolants at equal NFE. Gupta et al. [2026] allocate Euler steps where a learned field is sharp; the path is held fixed. Yuan et al. [2026] learn instance-aware grids. Du et al. [2026] select paths by a least-action principle. Khan [2026] regularizes the material derivative of learned fields.

Tao and Choi [2026] decompose $\nabla v = S + \Omega$ and control Euler error by the logarithmic norm $\mu_2 = \lambda_{\max}(S)$, which is signed, whereas $\mathcal{R}$ integrates $\|A\|_2^2$. In one dimension $\mu_2(a) = a$ and $\|A\|_2 = |a|$. Their Theorem 5 states that McCann displacement interpolation has vanishing material derivative, hence Euler is of order 2 in the Eulerian analysis; Remark 7 records the stronger Lagrangian exactness observed on that interpolant. Neither statement applies to the independent-coupling linear or VP path studied here. The present paper supplies a rationally certified Gaussian instance of ranking failure for the unsigned average $\|A\|_2^2$, together with a three-comparison map of pairwise ranking on this commuting Gaussian family. A counterexample to μ_2 itself is outside our scope. Benton et al. [2024] bound W_2 of a learned flow by an L^2 velocity error and a Lipschitz constant of the approximate field: an error bound, not a pairwise ranking of two interpolants at equal NFE. Consistency models [Song et al., 2023] distill few-step maps. Align-your-steps [Sabour et al., 2024] and DPM-Solver [Lu et al., 2022a] design the solver grid rather than the interpolant. Those objects are distinct from the ranking question tested here.

6 Conclusion

Averaged squared Jacobian regularity does not determine the equal-NFE Gaussian W_2 ranking of two fixed interpolants. The fact is proved by a one-dimensional Heun calculation whose integrals and linear factor are exact and whose VP factor is enclosed by a rational substitution in $\mathbb{Q}[\pi, \sqrt{2}]$.

The same scalar splits into three comparisons. As a pairwise comparator of linear versus VP it inverts on a complete census. As a pairwise comparison of two shared-schedule Class Q paths it still inverts. Among the four candidate paths on the stated 36-block census, the per-mode log-covariance path has the smallest continuous $\mathcal{R}$ and the smallest Gaussian W_2 in 36 of 36 blocks. This finite-census observation does not imply that a global $\mathcal{R}$ -minimizer minimizes fixed-NFE error. When Σ_1 has at least two distinct eigenvalues, that path is not a Class S shared interpolant. Appendix C gives an admissible Class S $\mathcal{R}$ -minimizer whose left-endpoint Euler error is strictly larger than that of a higher- $\mathcal{R}$ Class S path.

7 Limitations

The systems are commuting affine Gaussian interpolants. No learned neural field, image, video, or world-model quantity is reported. The solvers are three fixed-step Runge–Kutta methods at small equal-NFE budgets. Exponential integrators solve affine scalar modes exactly, so the comparisons concern generic RK discretizations

of those modes rather than the best integrator for an affine field. The Euler construction of Appendix C depends on the grid: the field is chosen after the grid. Extended robustness grids (NFE 64 and 128, $\pm 10\%$ noise-floor perturbations) are post-hoc. Census counts are hierarchical enumerations, not population frequencies. No counterexample to the signed logarithmic norm μ_2 is provided.

References

- Michael S. Albergo, Nicholas M. Boffi, and Eric Vanden-Eijnden. Stochastic interpolants: A unifying framework for flows and diffusions. *Journal of Machine Learning Research*, 26(209):1–80, 2025. arXiv:2303.08797.
- Joe Benton, George Deligiannidis, and Arnaud Doucet. Error bounds for flow matching methods. *Transactions on Machine Learning Research*, 2024. arXiv:2305.16860.
- Yifan Chen, Eric Vanden-Eijnden, and Jiawei Xu. Lipschitz-guided design of interpolation schedules in generative models. *arXiv preprint arXiv:2509.01629*, 2025. version 3, 16 May 2026.
- Shukai Du, Junzhe Zhang, and Yiming Li. Lagrangian flow matching: A least-action framework for principled path design. *arXiv preprint arXiv:2605.15419*, 2026.
- Matthias Gelbrich. On a formula for the L^2 Wasserstein metric between measures on Euclidean and Hilbert spaces. *Mathematische Nachrichten*, 147:185–203, 1990. doi: 10.1002/mana.19901470121.
- Aditi Gupta, Soon Hoe Lim, Annan Yu, and N. Benjamin Erichson. Sharpen your flow: Sharpness-aware sampling for flow matching. *arXiv preprint arXiv:2605.11547*, 2026.
- Ernst Hairer, Syvert P. Nørsett, and Gerhard Wanner. *Solving Ordinary Differential Equations I: Nonstiff Problems*. Springer, 2nd edition, 1993.
- Tero Karras, Miika Aittala, Timo Aila, and Samuli Laine. Elucidating the design space of diffusion-based generative models. In *Advances in Neural Information Processing Systems*, 2022. arXiv:2206.00364.
- Tauhid Khan. Isokinetic flow matching for pathwise straightening of generative flows. *arXiv preprint arXiv:2604.04491*, 2026.
- Yaron Lipman, Ricky T. Q. Chen, Heli Ben-Hamu, Maximilian Nickel, and Matthew Le. Flow matching for generative modeling. In *International Conference on Learning Representations (ICLR)*, 2023. arXiv:2210.02747.
- Xingchao Liu, Chengyue Gong, and Qiang Liu. Flow straight and fast: Learning to generate and transfer data with rectified flow. In *International Conference on Learning Representations (ICLR)*, 2023. arXiv:2209.03003.
- Cheng Lu, Yuhao Zhou, Fan Bao, Jianfei Chen, Chongxuan Li, and Jun Zhu. DPM-solver: A fast ODE solver for diffusion probabilistic model sampling in around 10 steps. In *Advances in Neural Information Processing Systems*, 2022a. arXiv:2206.00927.

- Cheng Lu, Yuhao Zhou, Fan Bao, Jianfei Chen, Chongxuan Li, and Jun Zhu. DPM-solver++: Fast solver for guided sampling of diffusion probabilistic models. *arXiv preprint arXiv:2211.01095*, 2022b.
- Gabriel Peyré and Marco Cuturi. *Computational Optimal Transport*. Now Publishers, 2019. arXiv:1803.00567.
- Amirmojtaba Sabour, Sanja Fidler, and Karsten Kreis. Align your steps: Optimizing sampling schedules in diffusion models. In *International Conference on Machine Learning (ICML)*, 2024. arXiv:2404.14507.
- Yang Song, Jascha Sohl-Dickstein, Diederik P. Kingma, Abhishek Kumar, Stefano Ermon, and Ben Poole. Score-based generative modeling through stochastic differential equations. In *International Conference on Learning Representations (ICLR)*, 2021. arXiv:2011.13456.
- Yang Song, Prafulla Dhariwal, Mark Chen, and Ilya Sutskever. Consistency models. In *International Conference on Machine Learning (ICML)*, 2023. arXiv:2303.01469.
- Arthur Stéphanovitch. Lipschitz regularity in flow matching and diffusion models: Sharp sampling rates and functional inequalities. *arXiv preprint arXiv:2604.06065*, 2026.
- Chenxi Tao and Seung-Kyum Choi. On the role of strain and vorticity in numerical integration error for flow matching. *arXiv preprint arXiv:2605.06680*, 2026.
- Alexander Tong, Kilian Fatras, Nikolay Malkin, Guillaume Huguet, Yanlei Zhang, Jarrid Rector-Brooks, Guy Wolf, and Yoshua Bengio. Improving and generalizing flow-based generative models with minibatch optimal transport. *Transactions on Machine Learning Research*, 2024. arXiv:2302.00482.
- Liangyu Yuan, Ruoyu Wang, Tong Zhao, Dingwen Fu, Mingkun Lei, Beier Zhu, and Chi Zhang. Few-step diffusion sampling through instance-aware discretizations. In *Proceedings of the IEEE/CVF Conference on Computer Vision and Pattern Recognition (CVPR)*, pages 35882–35892, 2026. arXiv:2603.17671.
- Qinsheng Zhang and Yongxin Chen. Fast sampling of diffusion models with exponential integrator. In *International Conference on Learning Representations (ICLR)*, 2023. arXiv:2204.13902.

A Proof of Proposition 2

Linear drift and integral. $q_{\text{lin}}(t) = (1-t)^2 + 4t^2 = 1 - 2t + 5t^2$ and $q' = -2 + 10t$, so $a_{\text{lin}}(t) = (5t-1)/q_{\text{lin}}(t)$. Completing the square yields the two equivalent identities

$$q_{\text{lin}}(t) = 5 \left[\left(t - \frac{1}{5} \right)^2 + \left(\frac{2}{5} \right)^2 \right], \quad (13)$$

$$q_{\text{lin}}(t) = 5 \left(t - \frac{1}{5} \right)^2 + \frac{4}{5}. \quad (14)$$

Thus $a_{\text{lin}}(t) = (t - 1/5)/((t - 1/5)^2 + (2/5)^2)$. The substitution $u = t - 1/5 = (2/5) \tan \theta$ gives $du = (2/5) \sec^2 \theta d\theta$ and converts $\int a_{\text{lin}}^2$ into $(5/2) \int \sin^2 \theta d\theta$. The definite limits are $\theta(0) = \arctan(-1/2)$ and $\theta(1) = \arctan 2$. The antiderivative $(5/4)(\theta - \sin \theta \cos \theta)$ therefore contributes $(5/4)(\arctan 2 + \arctan(1/2)) - (5/4)(4/5)$. Both angles lie in $(0, \pi/2)$ and $\tan(\arctan 2 + \arctan(1/2))$ is undefined, so the sum is $\pi/2$. Hence $\mathcal{R}_{\text{lin}} = 5\pi/8 - 1$. (The subtracted 1 is the boundary evaluation of $\sin \theta \cos \theta = \tan \theta / (1 + \tan^2 \theta)$.)

VP drift and integral. $q_{\text{VP}}(t) = \cos^2(\pi t/2) + 4 \sin^2(\pi t/2) = (5 - 3 \cos(\pi t))/2$ and $q' = (3\pi/2) \sin(\pi t)$, hence

$$a_{\text{VP}}(t) = \frac{(3\pi/2) \sin(\pi t)}{5 - 3 \cos(\pi t)} = \frac{(3\pi/4) \sin(\pi t)}{q_{\text{VP}}(t)}. \quad (15)$$

The change of variables $u = \pi t$ yields $\mathcal{R}_{\text{VP}} = (9\pi/4) \int_0^\pi \sin^2 u / (5 - 3 \cos u)^2 du$. The standard Weierstrass evaluation of the integrand gives $\int_0^\pi \sin^2 u / (5 - 3 \cos u)^2 du = \pi/36$, hence $\mathcal{R}_{\text{VP}} = \pi^2/16$. Both closed forms were checked by adaptive quadrature (absolute discrepancy below 10^{-12} ; quadrature error estimates are not used as certificates).

Heun recurrence. On $x' = a(t)x$ the implemented Heun factor is $r = 1 + (h/2)(a(t_n) + a(t_{n+1})(1 + ha(t_n)))$. With $h = 1/4$ the four linear factors are the rationals $47/52$, $6/5$, $497/370$, and $97/74$, all positive, with product $6797469/3559400$. Then $2 - r = 321331/3559400$, and $321331 \cdot 1000 < 91 \cdot 3559400$ is an integer comparison, so $W_2^{\text{lin}} < 0.091$.

Exact VP grid and rational certificate. At the Heun nodes the VP drift takes the values

$$\begin{array}{cccccc} t & 0 & 1/4 & 1/2 & 3/4 & 1 \\ a_{\text{VP}}(t) & 0 & \frac{(9+15\sqrt{2})\pi}{82} & \frac{3\pi}{10} & \frac{(-9+15\sqrt{2})\pi}{82} & 0 \end{array} \quad (16)$$

Positivity of $a(3/4)$ follows from $15\sqrt{2} > 9$, or $15^2 \cdot 2 - 9^2 = 369 > 0$ after squaring. Each Heun factor is therefore strictly positive, so $0 < r_{\text{VP}}$ and $W_2^{\text{VP}} = 2 - r_{\text{VP}}$. The four-step product is the following element $P(\pi, \sqrt{2})$ of $\mathbb{Q}[\pi, \sqrt{2}]$, obtained by expanding the four Heun factors and reducing $\sqrt{2}^2 = 2$:

$$\begin{aligned} P(\pi, \sqrt{2}) = & 1 + \frac{3}{40}\pi + \frac{15}{164}\pi\sqrt{2} + \frac{78579}{10758400}\pi^2 + \frac{45}{5248}\pi^2\sqrt{2} \\ & + \frac{11421}{17213440}\pi^3 + \frac{4671}{17213440}\pi^3\sqrt{2} \\ & + \frac{5103}{275415040}\pi^4 + \frac{567}{55083008}\pi^4\sqrt{2} \\ & + \frac{243}{2203320320}\pi^5 + \frac{729}{2203320320}\pi^5\sqrt{2} + \frac{729}{176265625600}\pi^6. \end{aligned} \quad (17)$$

Each of the twelve coefficients is a ratio of nonnegative integers, hence nonnegative, and the constant term is 1. Substituting any pair of upper bounds for π and $\sqrt{2}$ is therefore valid. The bound $\sqrt{2} < 99/70$ is the integer comparison $99^2 - 2 \cdot 70^2 = 1$. The bound $\pi < 355/113$ follows from Machin's formula $\pi = 16 \arctan(1/5) -$

$4 \arctan(1/239)$ together with Leibniz remainders: seven Taylor terms upper-bound $\arctan(1/5)$ and four terms lower-bound $\arctan(1/239)$. The resulting rational is strictly less than $355/113$ by an integer comparison. Substituting these enclosures yields

$$P\left(\frac{355}{113}, \frac{99}{70}\right) = \frac{192113353671412470139303}{102753427879939708813312} < \frac{187}{100}, \quad (18)$$

equivalently the integer comparison

$$19211335367141247013930300 < 19214891013548725548089344. \quad (19)$$

Hence $r_{\text{VP}} < 187/100$ and $W_2^{\text{VP}} > 13/100$. An independent interval enclosure at 40 decimal digits and an 80-digit non-interval product agree with this bound and with the float64 implementation to 10^{-14} . They are cross-checks, not the proof.

B Solver formulas and local truncation expansions

On $[t, t+h]$ for $x' = a(t)x$,

$$r_{\text{Euler}} = 1 + ha(t), \quad (20)$$

$$r_{\text{Heun}} = 1 + \frac{h}{2}(a(t) + a(t+h)(1 + ha(t))), \quad (21)$$

and classical RK4 with stages at t , $t+h/2$ (twice), and $t+h$. Equal NFE requires $B \equiv 0 \pmod{s}$ and sets $S = B/s$.

Assume a is C^3 . The exact factor admits $1 + ha + (h^2/2)(a' + a^2) + (h^3/6)(a'' + 3aa' + a^3) + O(h^4)$. Subtracting Euler leaves $\frac{1}{2}(a' + a^2)h^2 + O(h^3)$. Substituting $a(t+h)$ into the implemented Heun factor leaves $(-a''/12 + a^3/6)h^3 + O(h^4)$. The aa' terms cancel. The expansion was confirmed with independent symbolic algebra on the implemented expression.

C Grid-aware Euler construction

Proposition 3. *Fix $L > 0$ and an integer $N \geq 1$. Let $h = 1/N$ and let forward Euler use the left-endpoint grid $t_n = nh$ on $[0, 1]$. Set $\varepsilon_N = N(e^{L/N} - 1) - L > 0$ and define $a_0(t) = L$ and $a_{1,N}(t) = L + \varepsilon_N \cos(2\pi Nt)$. For $x(0) = 1$: both exact endpoints equal e^L ; $\int_0^1 a_{1,N}^2 = L^2 + \varepsilon_N^2/2 > \int_0^1 a_0^2$; Euler is exact for $a_{1,N}$; Euler has strictly positive endpoint error for a_0 .*

Proof. The solution of $x' = a(t)x$, $x(0) = 1$, is $x(t) = \exp(\int_0^t a(s) ds)$. The cosine perturbation has integer period, so $\int_0^1 \cos(2\pi Nt) dt = 0$ and both exact endpoints equal e^L . Expanding the square and using the same orthogonality gives $\int_0^1 a_{1,N}^2 = L^2 + \varepsilon_N^2/2$. The inequality $\varepsilon_N > 0$ is $e^z > 1 + z$ at $z = L/N > 0$, hence $\int_0^1 a_{1,N}^2 > \int_0^1 a_0^2 = L^2$.

On the left-endpoint Euler grid, $\cos(2\pi Nt_n) = \cos(2\pi n) = 1$, so $1 + ha_{1,N}(t_n) = 1 + (L + \varepsilon_N)/N = e^{L/N}$. Every oscillatory Euler factor equals $e^{L/N}$ and the product is e^L . For the constant field every Euler factor equals $1 + L/N$, and the product is $(1 + L/N)^N$. Since $\log(1 + z) < z$ for $z = L/N > 0$, one has $(1 + L/N)^N < e^L$. $\square$

The field $a_{1,N}$ is chosen after the grid. The construction is not the mechanism of Proposition 2. An 80-digit check confirms that the oscillatory Euler product equals e^L , but it is not a substitute for the argument above.

The same fields are Class S Gaussian interpolants. Set $\lambda = e^{2L}$ and

$$\begin{aligned} q_k(t) &= \exp\left(2 \int_0^t a_k(s) ds\right), \\ \alpha_k(t) &= \sqrt{q_k(t)} \cos(\pi t/2), \\ \sigma_k(t) &= \sqrt{q_k(t)/\lambda} \sin(\pi t/2). \end{aligned} \tag{22}$$

Then $q_k(0) = 1$, $\int_0^1 a_k = L$ so $q_k(1) = \lambda$, and the endpoint conditions $\alpha_k(0) = \sigma_k(1) = 1$, $\alpha_k(1) = \sigma_k(0) = 0$ hold. Positivity $q_k > 0$ and C^∞ regularity of a_k give $\alpha_k, \sigma_k \in C^\infty([0, 1])$, hence the embedding belongs to Class S. The identity $\alpha_k^2 + \lambda \sigma_k^2 = q_k$ is trigonometric, and $q'_k/(2q_k) = a_k$ by construction. Among scalar fields with fixed integral L , a_0 uniquely minimizes $\int_0^1 a^2$ (Proposition 1). Euler has strictly positive endpoint error for a_0 , while $a_{1,N}$ has strictly larger regularity and left-endpoint Euler is exact for $a_{1,N}$. Thus a_0 is an admissible Class S $\mathcal{R}$ -minimizer among one-dimensional variance paths from 1 to λ , yet left-endpoint Euler has strictly larger endpoint error for a_0 than for the higher- $\mathcal{R}$ path $a_{1,N}$. This construction blocks any inference from a finite-census 36 of 36 count on four candidates to the statement that a global $\mathcal{R}$ -minimizer minimizes fixed-NFE error.

D Full block tables

Table 3 lists every inverted linear-versus-VP solver-budget row under continuous $\mathcal{R}$. These rows are a hierarchical listing of the 5 geometry $\times$ solver cells, not a population rate. Table 4 records the strongest centered block. Table 5 reports the four-path census: among four candidate paths, the per-mode schedule has the smallest $\mathcal{R}$ and the smallest W_2 in 36 of 36 blocks. This finite-census observation does not imply that a global $\mathcal{R}$ -minimizer minimizes fixed-NFE error. When Σ_1 has at least two distinct eigenvalues, that path is not a Class S shared interpolant. Table 6 lists the nine shared-schedule pairwise inversions. Figure 5 illustrates identity (12) on the strongest block; Figure 6 shows the low-rank solver-path split.

E Non-centered structural stress test

The quantity $\mathcal{R}$ depends only on $A(t)$. If $c(t) \neq 0$, the mean still contributes $\|\hat{m} - \mu_1\|_2^2$ to W_2^2 . Dimensions $d \in \{2, 8\}$, coordinates indexed by $i = 0, \dots, d-1$. Source $N(\mu_0, I)$ with $\mu_{0,i} = 0.75(-1)^i$. Target $N(\mu_1, D)$ with $\mu_{1,i} = 1 + 0.25i$ and $D = \text{diag}(\lambda_i)$, where the λ_i form a geometric sequence from $6^{-1/2}$ to $6^{1/2}$ (condition number 6). Then $c(t) \neq 0$. Paths are linear and trigonometric VP. Solvers are Euler, Heun, and RK4 at NFE $B \in \{8, 16, 32\}$. Write $M = \|\hat{m} - \mu_1\|_2^2$ and $C = W_2^2 - M$, and

$$\Delta_{\text{mean}} = M_{\text{lin}} - M_{\text{VP}}, \quad \Delta_{\text{cov}} = C_{\text{lin}} - C_{\text{VP}}, \quad \Delta_{\text{total}} = \Delta_{\text{mean}} + \Delta_{\text{cov}}. \tag{23}$$

Table 3: All 14 inverted solver-budget rows among 36 centered blocks, grouped into 5 of 12 geometry×solver cells. Census listing, not a sample.

family	d	solver	NFE	R prefers	W_2 prefers	W_2 margin
anisotropic	2	heun	8	VP	linear	1.905310×10^{-3}
anisotropic	2	heun	16	VP	linear	1.939806×10^{-3}
anisotropic	2	heun	32	VP	linear	6.634439×10^{-4}
anisotropic	8	heun	16	VP	linear	1.758942×10^{-3}
anisotropic	8	heun	32	VP	linear	7.628173×10^{-4}
low-rank	2	heun	8	VP	linear	1.836251×10^{-2}
low-rank	2	heun	16	VP	linear	6.039925×10^{-3}
low-rank	2	heun	32	VP	linear	1.698035×10^{-3}
low-rank	2	rk4	8	VP	linear	1.516722×10^{-3}
low-rank	2	rk4	16	VP	linear	1.289981×10^{-4}
low-rank	2	rk4	32	VP	linear	1.118892×10^{-5}
low-rank	8	euler	8	linear	VP	3.543761×10^{-1}
low-rank	8	euler	16	linear	VP	2.046946×10^{-1}
low-rank	8	euler	32	linear	VP	1.096576×10^{-1}

Table 4: Strongest centered inversion: low-rank, $d = 8$, Euler, NFE 8. $\mathcal{R}$ is the continuous integral; $\widehat{\mathcal{R}}_{24}$ is shown only as a check. Eigenvalues: 0.05 (six times), 4.64827725629505, 10.675638226546441. Exact calculation.

path	$\mathcal{R}$	$\widehat{\mathcal{R}}_{24}$	Gaussian W_2
linear	2.9441044083	2.9476523251	0.8108540111
VP	4.7305438136	4.7295206355	0.4564779075

A block is mean-driven if $\Delta_{\text{mean}}\Delta_{\text{total}} > 0$ and $|\Delta_{\text{mean}}| > |\Delta_{\text{cov}}|$, and covariance-driven if $\Delta_{\text{cov}}\Delta_{\text{total}} > 0$ and $|\Delta_{\text{cov}}| > |\Delta_{\text{mean}}|$. Otherwise it is mixed. The same paths, solvers, and budgets give 18 blocks, of which 11 invert under continuous $\mathcal{R}$. Table 7 reports the driver of each path preference. Several inversions, including all dimension-8 Euler and Heun inversions, are mean-driven: $\mathcal{R}$ cannot see them. The family was specified before execution as a stress test of this structural blind spot, not as an independent replication of the centered mechanism.

F Quadrature, precision, and interval procedures

Table 9 compares the continuous integral with the historical 24-node trapezoidal estimator on every headline geometry. Path orderings agree. Trapezoidal values at 24, 48, 96, and 192 nodes converge to the same sign. Adaptive quadrature is used for multi-mode $\mathcal{R}$; the scalar integrals of Appendix A are closed form. Scipy error estimates are not interval certificates.

Because $\mathcal{R}$ depends on neither solver nor budget, the 36 linear-versus-VP rows nest into 12 geometry×solver cells. The two anisotropic geometries share one $\mathcal{R}$ pair, so the census contains three distinct $\mathcal{R}$ comparisons:




Figure 5: Modal contributions to squared Wasserstein error and transported signed defects for the strongest centered inversion. The bars reconstruct the telescoping identity (12). Exact modal identity on one block; illustration, not an independent causal test.

geometry	$(\mathcal{R}_{\text{lin}}, \mathcal{R}_{\text{VP}})$	distinct $\mathcal{R}$ pair
anisotropic $d = 2$ and $d = 8$	(0.972, 0.192)	1 (shared)
low-rank $d = 2$	(1.246, 0.625)	1
low-rank $d = 8$	(2.944, 4.731)	1

Frozen Gaussian W_2 on the centered census was checked against an 80-digit Gelbrich reference. The maximum observed gap is the numerical floor 9.7×10^{-10} . The paired concordance score $(n_{\text{agree}} - n_{\text{invert}})/n$ between the $\mathcal{R}$ order and the W_2 order is a descriptive census statistic, not a test (Table 12). On the 36 linear-versus-VP blocks, the score is 2/9. Stratified by solver, with $n = 4$ geometries and a structural tie between the two anisotropic targets, the score is $+1/2$ under Euler and RK4 and $-1/3$ under Heun.

Scalar linear Heun factors are computed in exact rationals. The VP Heun product is certified by substituting $\pi < 355/113$ and $\sqrt{2} < 99/70$ into a nonnegative element of $\mathbb{Q}[\pi, \sqrt{2}]$; the ranking inequality is the integer comparison in Appendix A. An independent interval enclosure at 40 decimal digits (`mpmath` 1.3.0) and an 80-digit (non-interval) product are recorded as cross-checks against the float64 Heun implementation. Default numerical precision for the benchmark runs is float64.

Table 10 reports the log-Lebesgue measure of the linear-versus-VP inversion set on $[0.05, 100]$. The declared resolution is 1281 log-uniform nodes; doubling to 2561 nodes changes every cell by less than 1%.

G Robustness grids

The post-hoc low-rank grid has 66 preference blocks: 2 dimensions, 3 solvers, and, for each pair, the three primary NFE budgets $\{8, 16, 32\}$ at noise-floor perturbations $\delta \in \{-0.1, 0, 0.1\}$, plus NFE $\{64, 128\}$ at $\delta = 0$ only. The perturbation multiplies the noise variance 0.05 by $(1 + \delta)$; the factor F is regenerated from seed 271828 and




Figure 6: Signed low-rank W_2 margin at equal NFE. Positive values prefer VP; negative values prefer linear. Vertical axis: symmetric-log with linear threshold 10^{-4} . Exact-moment modal Runge–Kutta.

is not redrawn. Extra budgets are not run at $\delta \neq 0$. Table 11 summarizes the 66 blocks. All 66 follow the same solver-path split as the primary low-rank census: VP under Euler, linear under Heun and RK4.

The geometry-draw experiment of Section 4.7 is the one place F is redrawn. The paired concordance score on the centered census is discussed in Appendix F and recorded in Table 12.

H Reproducibility, artifacts, and provenance

Default numerical precision for the benchmark runs is float64. The scalar calculation uses exact rational and algebraic expressions together with validated bounds; the precision audit uses 80-digit arithmetic. Compact artifacts used by this PDF live under `paper/arxiv/artifacts/`. They are committed files. Commands that rebuild tables, figures, and checksums are listed in the repository `README.md`. A GitHub Release tag is a later owner publication gate and is not required to read the numbers in this PDF. Experiment-code commit of the frozen Phase 4 grid: `e48c9390e62b38f206342e6aeb7f160122ccc79c`. The Hydra `outputs/` tree is not part of the public clone; eigenvalues, exact-moment modal Runge–Kutta, and the compact JSON files suffice for Table 4 and Proposition 2. Coefficient records for the VP Heun polynomial occupy `paper/arxiv/artifacts/scalar_counterexample.json`. Covariance matrices occupy `paper/arxiv/artifacts/geometries.json`. The 80-digit Gelbrich audit of the centered census occupies the frozen run `outputs/phase4_precision_2026-07-24-v1/` when that tree is present. Source and compact artifacts: <https://github.com/clement-callaert/fewstep-field-regularity>. The placeholder check must pass with no retired commit token remaining. The release check verifies compact checksums. Tag existence is a manual publication gate, not a scientific failure of this manuscript.




Figure 7: Signed path differences on all 18 non-centered blocks. $\Delta_{\text{mean}} = M_{\text{lin}} - M_{\text{VP}}$ and $\Delta_{\text{cov}} = C_{\text{lin}} - C_{\text{VP}}$. A positive bar means the linear path has the larger contribution; a negative bar means VP has the larger contribution. The axis is symmetric-log with linear threshold 10^{-6} . Stars mark ranking inversions. Exact numerical values appear in Table 8.

Table 5: Four candidate paths on the centered grid. Among these four paths, the per-mode schedule has the smallest $\mathcal{R}$ and the smallest W_2 in 36 of 36 blocks. This finite-census observation does not imply that a global $\mathcal{R}$ -minimizer minimizes fixed-NFE error. When Σ_1 has at least two distinct eigenvalues, that path is not a Class S shared interpolant.

family	d	solver	NFE	R min	W_2 min	$W_{2,\log}$	$W_{2,\text{lin}}$
anisotropic	2	euler	8	log-cov	log-cov	0.01164	0.2441
anisotropic	2	euler	16	log-cov	log-cov	0.005874	0.1266
anisotropic	2	euler	32	log-cov	log-cov	0.002952	0.06454
anisotropic	2	heun	8	log-cov	log-cov	0.0006613	0.03243
anisotropic	2	heun	16	log-cov	log-cov	0.0001682	0.006327
anisotropic	2	heun	32	log-cov	log-cov	4.244e-05	0.001359
anisotropic	2	rk4	8	log-cov	log-cov	3.834e-06	0.01119
anisotropic	2	rk4	16	log-cov	log-cov	2.473e-07	0.000213
anisotropic	2	rk4	32	log-cov	log-cov	1.575e-08	9.781e-06
anisotropic	8	euler	8	log-cov	log-cov	0.0131	0.4509
anisotropic	8	euler	16	log-cov	log-cov	0.006608	0.2337
anisotropic	8	euler	32	log-cov	log-cov	0.003319	0.1191
anisotropic	8	heun	8	log-cov	log-cov	0.0007027	0.04645
anisotropic	8	heun	16	log-cov	log-cov	0.0001786	0.008457
anisotropic	8	heun	32	log-cov	log-cov	4.503e-05	0.001746
anisotropic	8	rk4	8	log-cov	log-cov	3.897e-06	0.01812
anisotropic	8	rk4	16	log-cov	log-cov	2.513e-07	0.0003324
anisotropic	8	rk4	32	log-cov	log-cov	1.6e-08	1.37e-05
low-rank	2	euler	8	log-cov	log-cov	0.03303	0.2933
low-rank	2	euler	16	log-cov	log-cov	0.01683	0.1523
low-rank	2	euler	32	log-cov	log-cov	0.008497	0.07773
low-rank	2	heun	8	log-cov	log-cov	0.002998	0.0585
low-rank	2	heun	16	log-cov	log-cov	0.0007696	0.0122
low-rank	2	heun	32	log-cov	log-cov	0.0001953	0.002707
low-rank	2	rk4	8	log-cov	log-cov	4.7e-05	0.004578
low-rank	2	rk4	16	log-cov	log-cov	2.974e-06	0.000432
low-rank	2	rk4	32	log-cov	log-cov	1.89e-07	1.732e-05
low-rank	8	euler	8	log-cov	log-cov	0.2736	0.8109
low-rank	8	euler	16	log-cov	log-cov	0.1445	0.424
low-rank	8	euler	32	log-cov	log-cov	0.07439	0.2175
low-rank	8	heun	8	log-cov	log-cov	0.05274	0.4094
low-rank	8	heun	16	log-cov	log-cov	0.01399	0.08936
low-rank	8	heun	32	log-cov	log-cov	0.00363	0.02047
low-rank	8	rk4	8	log-cov	log-cov	0.004721	0.06962
low-rank	8	rk4	16	log-cov	log-cov	0.0002675	0.00746
low-rank	8	rk4	32	log-cov	log-cov	1.687e-05	0.0001322

Table 6: Shared-schedule pairwise inversions: trigonometric VP versus Chen Example 3.3 with $M = \lambda_{\max}$. Nine of 36 blocks; four of 12 cells. Post-hoc pairwise census, not a global family optimum.

family	d	solver	NFE	R prefers	W_2 prefers
anisotropic	2	euler	16	scalar	VP
anisotropic	2	euler	32	scalar	VP
anisotropic	8	euler	8	scalar	VP
anisotropic	8	euler	16	scalar	VP
anisotropic	8	euler	32	scalar	VP
low-rank	2	euler	8	VP	scalar
low-rank	2	heun	8	VP	scalar
low-rank	2	heun	16	VP	scalar
low-rank	2	heun	32	VP	scalar

Table 7: Non-centered family: all 18 blocks, with the signed-difference driver of the W_2^2 path preference $(\Delta_{\text{mean}}, \Delta_{\text{cov}})$.

d	solver	NFE	inversion	W_2 prefers	W_2^2 driver
2	euler	8	no	VP	covariance
2	euler	16	no	VP	covariance
2	euler	32	no	VP	covariance
2	heun	8	yes	linear	mean
2	heun	16	yes	linear	mean
2	heun	32	yes	linear	covariance
2	rk4	8	no	VP	covariance
2	rk4	16	no	VP	covariance
2	rk4	32	yes	linear	covariance
8	euler	8	yes	linear	mean
8	euler	16	yes	linear	mean
8	euler	32	yes	linear	mean
8	heun	8	yes	linear	mean
8	heun	16	yes	linear	mean
8	heun	32	yes	linear	mean
8	rk4	8	no	VP	covariance
8	rk4	16	no	VP	covariance
8	rk4	32	yes	linear	mean

Table 8: Signed path differences for the non-centered family. $\Delta_{\text{mean}} = M_{\text{lin}} - M_{\text{VP}}$, $\Delta_{\text{cov}} = C_{\text{lin}} - C_{\text{VP}}$, $\Delta_{\text{total}} = \Delta_{\text{mean}} + \Delta_{\text{cov}}$.

d	solver	NFE	inv.	Δ_{mean}	Δ_{cov}	Δ_{total}
2	euler	8	no	-1.969780×10^{-2}	7.008429×10^{-2}	5.038649×10^{-2}
2	euler	16	no	-5.010761×10^{-3}	1.895811×10^{-2}	1.394734×10^{-2}
2	euler	32	no	-1.266182×10^{-3}	4.941067×10^{-3}	3.674885×10^{-3}
2	heun	8	yes	-1.401380×10^{-3}	-8.296535×10^{-4}	-2.231034×10^{-3}
2	heun	16	yes	-8.132945×10^{-5}	-7.740260×10^{-5}	-1.587321×10^{-4}
2	heun	32	yes	-4.800285×10^{-6}	-5.428459×10^{-6}	-1.022874×10^{-5}
2	rk4	8	no	-1.931000×10^{-6}	7.169013×10^{-5}	6.975913×10^{-5}
2	rk4	16	no	$-5.235990 \times 10^{-10}$	1.106023×10^{-8}	1.053664×10^{-8}
2	rk4	32	yes	$-7.527744 \times 10^{-12}$	$-5.405709 \times 10^{-11}$	$-6.158484 \times 10^{-11}$
8	euler	8	yes	-3.252232×10^{-1}	2.207633×10^{-1}	-1.044599×10^{-1}
8	euler	16	yes	-8.388778×10^{-2}	5.947257×10^{-2}	-2.441521×10^{-2}
8	euler	32	yes	-2.129903×10^{-2}	1.546831×10^{-2}	-5.830720×10^{-3}
8	heun	8	yes	-5.927520×10^{-3}	-4.969512×10^{-4}	-6.424471×10^{-3}
8	heun	16	yes	-3.450747×10^{-4}	-9.787629×10^{-5}	-4.429510×10^{-4}
8	heun	32	yes	-2.074391×10^{-5}	-7.576731×10^{-6}	-2.832065×10^{-5}
8	rk4	8	no	-3.909251×10^{-6}	3.088406×10^{-4}	3.049314×10^{-4}
8	rk4	16	no	-1.137695×10^{-8}	7.433020×10^{-8}	6.295325×10^{-8}
8	rk4	32	yes	$-7.287617 \times 10^{-11}$	2.093892×10^{-11}	$-5.193726 \times 10^{-11}$

Table 9: Continuous $\mathcal{R}$ versus $\widehat{\mathcal{R}}_{24}$ on headline geometries, including the non-centered anisotropic family. Quadrature, not an exact integral except on the scalar pair.

geometry	$\mathcal{R}_{\text{lin}}$	$\widehat{\mathcal{R}}_{24}^{\text{lin}}$	$\mathcal{R}_{\text{VP}}$	$\widehat{\mathcal{R}}_{24}^{\text{VP}}$
scalar $\lambda = 4$	0.963495	0.964206	0.616850	0.616850
anisotropic $d = 2, 8$	0.972296	0.972637	0.192270	0.192361
low-rank $d = 2$	1.245531	1.244898	0.625461	0.625454
low-rank $d = 8$	2.944104	2.947652	4.730544	4.729521
non-centered anisotropic $d = 2, 8$	1.093537	1.093851	0.342354	0.342536

Table 10: Lebesgue measure in $\log \lambda$ of the linear-versus-VP inversion region on $[0.05, 100]$. Resolution 1281 nodes; doubling changes every entry by less than 1%. Not a count of inverted grid points.

solver	NFE	meas $_{\log \lambda}$	fraction of $[\ln 0.05, \ln 100]$
euler	8	2.444	0.321
euler	16	2.556	0.336
euler	32	2.669	0.351
heun	8	3.337	0.439
heun	16	3.723	0.490
heun	32	3.985	0.524
rk4	8	2.212	0.291
rk4	16	3.082	0.405
rk4	32	3.432	0.452

Table 11: Post-hoc low-rank robustness: all 66 blocks follow the same solver-path split as the primary grid. The factor F is not redrawn.

solver	$d = 2$ blocks	$d = 8$ blocks	preferred path
euler	11	11	VP
heun	11	11	linear
rk4	11	11	linear

Table 12: Paired concordance score $(n_{\text{agree}} - n_{\text{invert}})/n$ between the $\mathcal{R}$ order and the W_2 order on the linear-versus-VP census. Descriptive statistic on a complete enumeration, not Kendall's τ and not an inferential test.

stratum	paired concordance
census (36 blocks)	2/9
anisotropic $d = 2$	1/3
anisotropic $d = 8$	5/9
low-rank $d = 2$	-1/3
low-rank $d = 8$	1/3
Euler	1/2
Heun	-1/3
RK4	1/2