M and Z Estimators

Clément Lalanne

Notation for this lecture

Parameter and data. The parameter set is $\Theta \subset \mathbb{R}^p$ and the true parameter is θ_0. The observations $X_1, \ldots, X_n$ are i.i.d. copies of a random vector X of $\mathbb{R}^d$, of law $\mathcal{L}_{\theta_0}$ inside a model $\{\mathcal{L}_\theta; \theta \in \Theta\}$.

Criteria. $M_n : \Theta \to \mathbb{R}$ is a random function and $M : \Theta \to \mathbb{R}$ is its deterministic limit. For Z estimators, we use $Z_n : \Theta \to \mathbb{R}^q$ and $Z : \Theta \to \mathbb{R}^q$.

Random function means that, for every θ, $M_n(\theta)$ is a random variable, and that all the $M_n(\theta), \theta \in \Theta$, are defined on one probability space.

Subscripts. $\mathbb{E}_\theta$ is the expectation computed as if the parameter were θ, while $\mathbb{E}_X = \mathbb{E}_{\theta_0}$ is the real one. Suprema carry their index set, as in $\sup_{\theta \in \Theta}$.

Derivatives. $J\phi(\theta)$ denotes the Jacobian matrix of ϕ at θ, and $\nabla\phi(\theta)$ the gradient when ϕ is real valued.

Part I. Two new ways to define an estimator

M estimators

Definition. Let $(M_n)_{n\in\mathbb{N}}$ be a sequence of random functions from Θ to $\mathbb{R}$. An **M-estimator** is a sequence $(\hat{\theta}_n)_{n\in\mathbb{N}}$ of random vectors taking values in Θ such that, for every n,

$$\hat{\theta}_n \in \arg\max_{\theta\in\Theta} M_n(\theta) \ .$$

The letter M stands for maximizer. Since replacing M_n by $-M_n$ turns a maximization into a minimization, we lose nothing by always maximizing.

In practice, the almost sure requirement is too strong for anything computed numerically. This is why it is relaxed twice, first to holding with probability going to 1, and then to maximizing up to an additive $o_{\mathbb{P}}(1)$.

Exercise, the empirical mean

Exercise. Given random vectors $X_1, \dots, X_n$ of $\mathbb{R}^d$, express the empirical mean $\bar{X}_n = \frac{1}{n}\sum_{i\leq n} X_i$ as an M-estimator.

Solution. We take $\Theta = \mathbb{R}^d$ and

$$M_n(\theta) = -\frac{1}{n}\sum_{i\leq n} \|\theta - X_i\|^2 \ .$$

Then $\nabla M_n(\theta) = -\frac{2}{n}\sum_{i\leq n}(\theta - X_i)$ vanishes only at $\theta = \bar{X}_n$. Furthermore, since M_n is strictly concave and tends to $-\infty$ at infinity, it has a unique maximizer, which is thus $\bar{X}_n$.

Empirical risk minimization

Definition. Given a loss $\ell : \mathbb{R}^d \times \Theta \to \mathbb{R}$, the **empirical risk** and the **risk** of a parameter are

$$\widehat{R}_S(\theta) = \frac{1}{n} \sum_{i \leq n} \ell(X_i, \theta) \ ,$$
$$R(\theta) = \mathbb{E}_X[\ell(X, \theta)] \ ,$$

and the **empirical risk minimizer** is the M-estimator of $M_n = -\widehat{R}_S$, whose limit criterion is $M = -R$.

Languages from two communities. The two names come from two communities rather than from two objects, with $\ell = -m$ and $M = -R$. In supervised learning, $X = (x, y)$ and $\ell(X, \theta) = \ell(y, f_\theta(x))$ for a predictor f_θ.

Maximum likelihood

Assumption. For every θ, $\mathcal{L}_\theta$ has a density f_θ with respect to a reference measure ν.

The **likelihood** and the **log-likelihood** of the sample are $L_n(\theta) = \prod_{i \le n} f_\theta(X_i)$ and $\log L_n(\theta) = \sum_{i \le n} \log f_\theta(X_i)$.

Definition. The **maximum likelihood estimator** is the M-estimator associated with $M_n = \frac{1}{n} \log L_n$, that is, with

$$M_n(\theta) = \frac{1}{n} \sum_{i \le n} \log f_\theta(X_i) \ .$$

Equivalently, it is the empirical risk minimizer for the log loss $\ell(x, \theta) = -\log f_\theta(x)$, since $M_n = -\widehat{R}_S$ for that loss.

Idea. We choose the member of the model that makes the observed sample most probable. Note that the factor $\frac{1}{n}$ does not change the maximizer, and that it makes M_n converge to something finite.

Exercise, the gaussian model

Exercise. What is the maximum likelihood estimator in the gaussian model $\Theta = \mathbb{R} \times (0, \infty), \theta = (\mu, \sigma^2), \mathcal{L}_\theta = \mathcal{N}(\mu, \sigma^2)$?

Solution. With $f_{(\mu,\sigma^2)}(x) = \frac{1}{\sigma\sqrt{2\pi}} e^{-(x-\mu)^2/(2\sigma^2)}$, we obtain

$$M_n(\mu, \sigma^2) = \frac{1}{n} \sum_{i \leq n} \left\{ -\log \sigma - \frac{1}{2} \log(2\pi) - \frac{(X_i - \mu)^2}{2\sigma^2} \right\} .$$

$\partial_\mu M_n = \frac{1}{n\sigma^2} \sum_{i \leq n} (X_i - \mu)$ vanishes only at $\mu = \bar{X}_n$, and

$$\partial_{\sigma^2} M_n = -\frac{1}{2\sigma^2} + \frac{1}{2\sigma^4} \cdot \frac{1}{n} \sum_{i \leq n} (X_i - \mu)^2 ,$$

which vanishes at $\mu = \bar{X}_n$ exactly when $\sigma^2 = \frac{1}{n} \sum_{i \leq n} (X_i - \bar{X}_n)^2$.

We recover the pair of the method of moments of the previous lecture, although the two methods differ in general.

Z estimators

Definition. Let $(Z_n)_{n\in\mathbb{N}}$ be a sequence of random functions from Θ to $\mathbb{R}^q$. A **Z-estimator** is a sequence $(\hat{\theta}_n)_{n\in\mathbb{N}}$ of random vectors taking values in Θ such that, for every n, almost surely

$$Z_n(\hat{\theta}_n) = 0 \; .$$

The letter Z stands for zero.

Every smooth M-estimator is a Z-estimator. Indeed, if Θ is open and M_n is differentiable, a maximizer in the interior satisfies $\nabla M_n(\hat{\theta}_n) = 0$, so that $\hat{\theta}_n$ is a Z-estimator with $Z_n = \nabla M_n$ and $q = p$.

The method of moments is a Z-estimator, with $Z_n(\theta) = e_n - e(\theta)$ in the notation of the previous lecture.

Part II. Consistency

The two ingredients of consistency

We want to show that $\hat{\theta}_n \xrightarrow{\mathbb{P}} \theta_0$, a statement that mixes two things of completely different natures.

A statement about the criterion. M_n is close to M, uniformly in θ. All the randomness lies there.

A statement about the geometry. M separates θ_0 from the rest of Θ. There is no randomness in it at all.

Independence of the two statements. Each can hold while the other fails.

Stage one, the criterion error

Proposition. Let $\Delta_n = \sup_{\theta \in \Theta} |M_n(\theta) - M(\theta)|$ and assume that $\hat{\theta}_n$ nearly maximizes M_n, meaning

$$M_n(\hat{\theta}_n) \geq \sup_{\theta \in \Theta} M_n(\theta) + o_{\mathbb{P}}(1) \; .$$

Then

$$\sup_{\theta \in \Theta} M(\theta) - M(\hat{\theta}_n) \leq 2\Delta_n + o_{\mathbb{P}}(1) \; .$$

Proof. For every fixed $\theta \in \Theta$, we have

$$\begin{aligned} M(\hat{\theta}_n) &\geq M_n(\hat{\theta}_n) - \Delta_n && \text{by definition of } \Delta_n \\ &\geq M_n(\theta) - \Delta_n + o_{\mathbb{P}}(1) && \text{by near-maximization} \\ &\geq M(\theta) - 2\Delta_n + o_{\mathbb{P}}(1) \; . && \text{by definition of } \Delta_n \end{aligned}$$

Since the left side does not depend on θ, taking the supremum over θ on the right concludes the proof.

Generality of stage one

If $\Delta_n \xrightarrow{\mathbb{P}} 0$, the proposition takes the form

$$\sup_{\theta \in \Theta} M(\theta) - M(\hat{\theta}_n) \xrightarrow{\mathbb{P}} 0 \ .$$

No topology, no identifiability, no compactness, no true parameter. Θ is an arbitrary set, M is an arbitrary function on it, and we have not even used θ_0.

A statement about real numbers. The conclusion says that the value of the criterion at $\hat{\theta}_n$ approaches its optimal value, and nothing about where $\hat{\theta}_n$ is.

This is already the useful statement in many settings. In empirical risk minimization, the excess risk is this quantity, which generalization bounds control.

Empirical risk minimization as stage one

For the empirical risk minimizer of Part I, where $M = -R$ and $M_n = -\widehat{R}_S$, stage one takes the form

$$R(\hat{\theta}_n) - \inf_{\theta \in \Theta} R(\theta) \leq 2 \sup_{\theta \in \Theta} \left| \widehat{R}_S(\theta) - R(\theta) \right| + o_{\mathbb{P}}(1) \; .$$

Names of the terms. The left side is the estimation error, Δ_n is the uniform deviation, and the $o_{\mathbb{P}}(1)$ is the optimisation error (i.e. the gap between what the optimizer returns and the true empirical minimum).

Statistical learning theory as a whole is the estimation of Δ_n, by Rademacher complexities, VC dimension or metric entropy.

Irrelevance of stage two in learning

The parameter is not identifiable, on purpose. Permuting the units of a hidden layer, or rescaling one layer against the next, leaves f_θ and hence M unchanged. As a result, the label switching counterexample of Part II is the normal situation here rather than a pathology, and $\delta \equiv 0$ below the distance between two symmetric copies.

Over-parameterized models have a continuum of exact empirical minimizers, so that $\hat{\theta}_n$ is not well defined as a point and no statement about its convergence could hold.

The model is misspecified deliberately. There is no θ_0 with $\mathcal{L}_{\theta_0} = \mathbb{P}$, and $\inf_\theta R(f_\theta) - R^\star$ is the approximation error, i.e. a deterministic quantity that the estimator cannot affect.

What is asked of a learning algorithm is the risk of the predictor it returns

Now Stage Two: Well separated maximum

Definition. A point $\theta_0 \in \Theta$ is a **well separated maximizer** of M if $M(\theta_0) = \sup_{\theta \in \Theta} M(\theta)$ and $\delta(\epsilon) > 0$ for every $\epsilon > 0$, where

$$\delta(\epsilon) := M(\theta_0) - \sup_{\theta \in \Theta : \|\theta - \theta_0\| \geq \epsilon} M(\theta) \, .$$

Reading δ. It measures, in criterion, the loss incurred by standing at distance ϵ from θ_0 in parameter, and the condition says that this loss is positive at every distance.

It is deterministic. Indeed, δ depends on M and on Θ and on nothing else (no sample, no n, no probability).

Stage two, from the criterion to the parameter

Proposition. Let θ_0 be a well separated maximizer of M, and let $(\hat{\theta}_n)$ satisfy $M(\theta_0) - M(\hat{\theta}_n) \xrightarrow{\mathbb{P}} 0$. Then $\hat{\theta}_n \xrightarrow{\mathbb{P}} \theta_0$.

Proof. Let us fix $\epsilon > 0$. If $\|\hat{\theta}_n - \theta_0\| \geq \epsilon$, then $\hat{\theta}_n$ belongs to the set over which the supremum defining $\delta(\epsilon)$ is taken, so that

$$\begin{aligned} M(\hat{\theta}_n) &\leq \sup_{\theta \in \Theta : \|\theta - \theta_0\| \geq \epsilon} M(\theta) \\ &= M(\theta_0) - \delta(\epsilon) \ , \end{aligned}$$

that is, $M(\theta_0) - M(\hat{\theta}_n) \geq \delta(\epsilon)$.

Hence,

$$\mathbb{P}\left(\|\hat{\theta}_n - \theta_0\| \geq \epsilon\right) \leq \mathbb{P}\left(M(\theta_0) - M(\hat{\theta}_n) \geq \delta(\epsilon)\right) \ ,$$

and the right side tends to 0 since the criterion error tends to 0 in probability and $\delta(\epsilon)$ is a fixed positive number, which concludes the proof.

The consistency theorem for M estimators


Theorem. Let (M_n) be random functions from $\Theta \subset \mathbb{R}^p$ to $\mathbb{R}$ and $M : \Theta \to \mathbb{R}$ deterministic. Assume

1. $\sup_{\theta \in \Theta} |M_n(\theta) - M(\theta)| \xrightarrow{\mathbb{P}} 0$,
2. there is $\theta_0 \in \Theta$ such that for every $\epsilon > 0$, $\sup_{\theta \in \Theta : \|\theta - \theta_0\| \geq \epsilon} M(\theta) < M(\theta_0)$,
3. $(\hat{\theta}_n)$ satisfies $M_n(\hat{\theta}_n) \geq \sup_{\theta \in \Theta} M_n(\theta) + o_{\mathbb{P}}(1)$.

Then $\hat{\theta}_n \xrightarrow{\mathbb{P}} \theta_0$.

Proof. Hypotheses 1 and 3 yield stage one and hypothesis 2 is stage two, so that composing the two propositions concludes the proof.

The two stages, in one picture



Two gaps, two stages. The blue arrow is Δ_n (i.e. how far M_n sits from M uniformly), and stage one turns it into a bound on the vertical distance from the top. The black arrow is $\delta(\epsilon)$, the exchange rate between the axes, since leaving the interval lowers the criterion by $\delta(\epsilon)$, so that a criterion error below $\delta(\epsilon)$ keeps $\hat{\theta}_n$ inside it. A flat curve makes $\delta(\epsilon)$ zero and the exchange impossible.

Role of each hypothesis

Hypothesis 1 carries all the probability. It is a law of large numbers, uniform in θ, which does not mention θ_0 and to which the final part of the lecture is entirely devoted.

Hypothesis 2 carries all the geometry. It does not mention M_n, the sample, or n.

Hypothesis 3 is the definition of the estimator, weakened as far as possible.

A sufficient condition for well separation

Lemma. If Θ is compact, M is continuous on Θ, and θ_0 is the unique maximizer of M, then θ_0 is a well separated maximizer.

Proof. Let us fix $\epsilon > 0$ and let $K_\epsilon = \{\theta \in \Theta : \|\theta - \theta_0\| \geq \epsilon\}$. If K_ϵ is empty, the supremum is $-\infty$ and there is nothing to prove. Otherwise, K_ϵ is closed in the compact Θ, hence compact, and the continuous M attains its supremum on it at some $\theta_\epsilon \neq \theta_0$, so that

$$\begin{aligned} \sup_{\theta \in K_\epsilon} M(\theta) &= M(\theta_\epsilon) \\ &< M(\theta_0)\,. \quad \text{since } \theta_0 \text{ is the unique maximizer} \end{aligned}$$

From separation to a rate

Let us suppose that the separation is polynomial, i.e. $\delta(\epsilon) \geq c\epsilon^{\alpha}$ for some $c, \alpha > 0$ and all small ϵ.

If $\|\hat{\theta}_n - \theta_0\| = r$, then $\|\hat{\theta}_n - \theta_0\| \geq \epsilon$ for every $\epsilon \leq r$, so that the argument of stage two yields $M(\theta_0) - M(\hat{\theta}_n) \geq \delta(\epsilon) \geq c\epsilon^{\alpha}$ for every $\epsilon \leq r$. Hence, $M(\theta_0) - M(\hat{\theta}_n) \geq cr^{\alpha}$ and

$$\begin{aligned} \|\hat{\theta}_n - \theta_0\| &\leq \left(\frac{M(\theta_0) - M(\hat{\theta}_n)}{c} \right)^{1/\alpha} \\ &\leq \left(\frac{2\Delta_n + o_{\mathbb{P}}(1)}{c} \right)^{1/\alpha} . \qquad \text{by stage one} \end{aligned}$$

The generic case is $\alpha = 2$. If M is twice differentiable at θ_0 with negative definite Hessian, the separation is quadratic near θ_0, and the exponent $1/\alpha = 1/2$ loses a square root, since a typical Δ_n of order $n^{-1/2}$ only yields $n^{-1/4}$.

A pessimistic rate. The bound above uses the uniform Δ_n over all of Θ to control a local quantity. We latter show that replacing it by a modulus over a shrinking neighbourhood of θ_0 recovers the rate $n^{-1/2}$.

The same two stages for Z estimators

Proposition (stage one). If $\sup_{\theta \in \Theta} \|Z_n(\theta) - Z(\theta)\| \xrightarrow{\mathbb{P}} 0$ and $Z_n(\hat{\theta}_n) = o_{\mathbb{P}}(1)$, then $\|Z(\hat{\theta}_n)\| = o_{\mathbb{P}}(1)$.

Proof. By the triangle inequality, $\|Z(\hat{\theta}_n)\| \leq \|Z_n(\hat{\theta}_n)\| + \sup_{\theta \in \Theta} \|Z_n(\theta) - Z(\theta)\|$, and both terms are $o_{\mathbb{P}}(1)$.

Proposition (stage two). If $Z(\theta_0) = 0$ and $\delta(\epsilon) := \inf_{\|\theta - \theta_0\| \geq \epsilon} \|Z(\theta)\| > 0$ for every $\epsilon > 0$, then $\|Z(\hat{\theta}_n)\| = o_{\mathbb{P}}(1)$ implies $\hat{\theta}_n \xrightarrow{\mathbb{P}} \theta_0$.

Proof. $\|\hat{\theta}_n - \theta_0\| \geq \epsilon$ forces $\|Z(\hat{\theta}_n)\| \geq \delta(\epsilon)$, so $\mathbb{P}(\|\hat{\theta}_n - \theta_0\| \geq \epsilon) \leq \mathbb{P}(\|Z(\hat{\theta}_n)\| \geq \delta(\epsilon)) \to 0$.

The consistency theorem for Z estimators

Theorem. Let (Z_n) be random functions from $\Theta \subset \mathbb{R}^p$ to $\mathbb{R}^q$ and $Z : \Theta \to \mathbb{R}^q$ deterministic. Assume

1. $\sup_{\theta \in \Theta} \|Z_n(\theta) - Z(\theta)\| \xrightarrow{\mathbb{P}} 0$,
2. $Z(\theta_0) = 0$ and, for every $\epsilon > 0$, $\inf_{\theta \in \Theta : \|\theta - \theta_0\| \geq \epsilon} \|Z(\theta)\| > 0$,
3. $(\hat{\theta}_n)$ satisfies $Z_n(\hat{\theta}_n) = o_{\mathbb{P}}(1)$.

Then $\hat{\theta}_n \xrightarrow{\mathbb{P}} \theta_0$.


Remark. The course notes write hypothesis 2 as one chain, $\inf \|Z(\theta)\| > 0 = \|Z(\theta_0)\|$. It consists of two separate requirements, namely that the limit function vanishes at θ_0 and that it stays away from zero everywhere else.

Easy case, a monotone function in one dimension

Theorem. Let $\Theta = \mathbb{R}$, let (Z_n) be random functions from $\mathbb{R}$ to $\mathbb{R}$ and $Z : \mathbb{R} \to \mathbb{R}$ deterministic. Assume

1. for every **fixed** θ, $Z_n(\theta) \xrightarrow{\mathbb{P}} Z(\theta)$, with no uniformity,
2. almost surely, Z_n is non-decreasing,
3. there is θ_0 such that $Z(\theta_0 - \epsilon) < 0 < Z(\theta_0 + \epsilon)$ for every $\epsilon > 0$.

Then any $(\hat{\theta}_n)$ with $Z_n(\hat{\theta}_n) = o_{\mathbb{P}}(1)$ satisfies $\hat{\theta}_n \xrightarrow{\mathbb{P}} \theta_0$.



Proof in the monotone case

Let us fix $\epsilon > 0$. Splitting on the two sides, we have

$$
\begin{aligned}
&\mathbb{P}\left(|\hat{\theta}_n - \theta_0| \geq \epsilon\right) \\
&= \mathbb{P}(\hat{\theta}_n \leq \theta_0 - \epsilon) + \mathbb{P}(\hat{\theta}_n \geq \theta_0 + \epsilon) \\
&\leq \mathbb{P}\left(Z_n(\hat{\theta}_n) \leq Z_n(\theta_0 - \epsilon)\right) + \mathbb{P}\left(Z_n(\hat{\theta}_n) \geq Z_n(\theta_0 + \epsilon)\right) && \text{since } Z_n \text{ is non-decreasing} \\
&= \mathbb{P}\left(o_{\mathbb{P}}(1) \leq Z(\theta_0 - \epsilon) + o_{\mathbb{P}}(1)\right) + \mathbb{P}\left(o_{\mathbb{P}}(1) \geq Z(\theta_0 + \epsilon) + o_{\mathbb{P}}(1)\right) \ . && \text{by hypotheses 3 and 1}
\end{aligned}
$$

The two numbers $Z(\theta_0 - \epsilon) < 0$ and $Z(\theta_0 + \epsilon) > 0$ are fixed, and each probability is the probability that a quantity tending to 0 in probability exceeds a fixed nonzero threshold in the wrong direction, so that both tend to 0, which concludes the proof.

The trade-off. Uniform convergence, the demanding hypothesis of the general theorem, is replaced by pointwise convergence at two points per ϵ. In exchange, the statement needs monotonicity and dimension one.

Exercise, the empirical median

Exercise. Let $X_1, X_2, \ldots$ be i.i.d. copies of a real X with a continuous density bounded away from 0 on a neighbourhood of the population median. Show that the empirical median $\hat{\theta}_n$, defined by $\sum_{i \le n} \operatorname{sign}(\hat{\theta}_n - X_i) = 0$, is consistent.

Solution. We take $Z_n(\theta) = \frac{1}{n} \sum_{i \le n} \operatorname{sign}(\theta - X_i)$, so that $Z_n(\hat{\theta}_n) = 0$, and

$$\begin{aligned} Z_n(\theta) &= \frac{1}{n} \sum_{i \le n} \mathbf{1}\{X_i < \theta\} - \frac{1}{n} \sum_{i \le n} \mathbf{1}\{X_i > \theta\} \\ &\xrightarrow{\mathbb{P}} \mathbb{P}(X < \theta) - \mathbb{P}(X > \theta) && \text{by the law of large numbers} \\ &= 2F_X(\theta) - 1\,, && \text{since } \mathbb{P}(X = \theta) = 0 \end{aligned}$$

which we denote $Z(\theta)$.

Hypothesis 2 holds since $\theta \mapsto \operatorname{sign}(\theta - x)$ is non-decreasing for each x, and hypothesis 3 holds with θ_0 the median ($F_X(\theta_0) = 1/2$), since F_X is strictly increasing where the density is positive. Hence, the theorem applies.

Part III. Uniform convergence

The uniform deviation

For both consistency theorems to apply, we need the uniform deviation of stage one to tend to 0,

$$\Delta_n = \sup_{\theta \in \Theta} |M_n(\theta) - M(\theta)| \xrightarrow{\mathbb{P}} 0 \ .$$

Simplification. Let $X_1, X_2, \ldots$ i.i.d... We consider $M_n(\theta) = \frac{1}{n} \sum_{i \leq n} m(X_i, \theta)$ and $M(\theta) = \mathbb{E}_X[m(X, \theta)]$ for a function $m : \mathbb{R}^d \times \Theta \to \mathbb{R}$.

Pointwise VS Uniform convergence. For each θ, the law of large numbers yields $M_n(\theta) \xrightarrow{\mathbb{P}} M(\theta)$, but uniformity in θ is out of its reach.

Rewriting. With $\mathcal{F} = \{m(\cdot, \theta); \theta \in \Theta\}$, the uniform deviation becomes

$$\Delta_n = \sup_{f \in \mathcal{F}} \left| \frac{1}{n} \sum_{i \leq n} f(X_i) - \mathbb{E}_X[f(X)] \right| \ .$$

Glivenko-Cantelli classes. With $\mathcal{L}$ the law of X, we say that $\mathcal{F}$ is **$\mathcal{L}$-Glivenko-Cantelli** if $\Delta_n \xrightarrow{\mathbb{P}} 0$.

First idea: Covering in supremum norm

If $\mathcal{F}$ is finite, the law of large numbers applied to each element and a maximum over finitely many terms ssolves the problem by union bound.

Otherwise, an idea could be to approximate $\mathcal{F}$ by a finite family of functions. If every $f \in \mathcal{F}$ has some f' in a finite $\mathcal{F}'$ with $\|f - f'\|_\infty \leq \epsilon$, then

$$\sup_{f \in \mathcal{F}} \left| \frac{1}{n} \sum_{i \leq n} f(X_i) - \mathbb{E}_X[f(X)] \right| \leq \sup_{f' \in \mathcal{F}'} \left| \frac{1}{n} \sum_{i \leq n} f'(X_i) - \mathbb{E}_X[f'(X)] \right| + 2\epsilon \ .$$


Exercise. Compute the smallest cardinality of such an $\mathcal{F}'$ for $\mathcal{F} = \{\mathbf{1}_{(-\infty,t]}; t \in [0,1]\}$.

Solution. For $s \neq t$, the two indicators differ by 1 somewhere, so that $\|\mathbf{1}_{(-\infty,t]} - \mathbf{1}_{(-\infty,s]}\|_\infty = 1$. Hence, for $\epsilon < 1/2$, a ball of radius ϵ contains at most one element of $\mathcal{F}$, and since $\mathcal{F}$ is uncountable, no finite $\mathcal{F}'$ works. The covering number is infinite on the simplest class there is, which is nevertheless $\mathcal{L}$-Glivenko-Cantelli for every law $\mathcal{L}$ by the classical Glivenko-Cantelli theorem.

Refinement: Brackets

Definition. For $l \le u$ pointwise, the **bracket** $[l, u]$ is the set of functions $f : \mathbb{R}^d \to \mathbb{R}$ with $l \le f \le u$ pointwise. For $\epsilon > 0$, $q > 0$ and a law $\mathcal{L}$ on $\mathbb{R}^d$, the **bracketing number** $N_{[]}(\mathcal{F}, L^q(\mathcal{L}), \epsilon)$ is the smallest N for which some brackets $[l_1, u_1], \ldots, [l_N, u_N]$ cover $\mathcal{F}$ and satisfy, for every j,

$$\left(\int (u_j - l_j)^q d\mathcal{L} \right)^{1/q} \le \epsilon \, .$$




Two relaxations. The width is measured in $L^q(\mathcal{L})$ rather than in supremum norm, and l, u (drawn dashed) need not belong to $\mathcal{F}$.

Exercise, the indicator class

Exercise. Let $\mathcal{L} = \mathrm{Unif}([0,1])$ and $\mathcal{F} = \{f_t = \mathbf{1}_{(-\infty,t]}; t \in [0,1]\}$. Find $N_{[]}(\mathcal{F}, L^2(\mathcal{L}), \epsilon)$ up to multiplicative constants.

Solution, upper bound. For $s \leq t$ in $[0,1]$, $\|f_t - f_s\|_{L^2(\mathcal{L})} = \sqrt{t-s}$, and $f_r \in [f_s, f_t]$ for every $r \in [s,t]$. Hence, the brackets $[f_0, f_{\epsilon^2}], [f_{\epsilon^2}, f_{2\epsilon^2}], \ldots$ cover $\mathcal{F}$, have width at most ϵ, and there are $\lceil \epsilon^{-2} \rceil$ of them.



Solution, lower bound. If $f_s, f_t \in [l,u]$, then $u - l \geq |f_t - f_s|$ pointwise, so $\|u-l\|_{L^2(\mathcal{L})} \geq \sqrt{|t-s|}$. A bracket of width ϵ thus contains f_t only for t in a set of diameter at most ϵ^2, and covering $[0,1]$ requires at least ϵ^{-2} of them. Hence, $N_{[]}(\mathcal{F}, L^2(\mathcal{L}), \epsilon) \asymp \epsilon^{-2}$.

Consistency theorem: Finite bracketing complexity => Glivenko-Cantelli

Theorem. Let $\mathcal{F}$ be a class of $\mathcal{L}$-integrable functions with $N_{[]}(\mathcal{F}, L^1(\mathcal{L}), \epsilon) < \infty$ for every $\epsilon > 0$. Then $\mathcal{F}$ is $\mathcal{L}$-Glivenko-Cantelli.

Proof. Let us fix $\epsilon > 0$, let $N = N_{[]}(\mathcal{F}, L^1(\mathcal{L}), \epsilon)$ and let $[l_j, u_j], j \leq N$, be brackets covering $\mathcal{F}$ with $\int |u_j - l_j| d\mathcal{L} \leq \epsilon$. Discarding the brackets that meet no element of $\mathcal{F}$, each l_j and u_j is within ϵ in L^1 of an integrable function, hence integrable.

Given $f \in \mathcal{F}$, we choose j with $l_j \leq f \leq u_j$. Then, $\mathbb{E}_X[l_j(X)] \leq \mathbb{E}_X[f(X)] \leq \mathbb{E}_X[u_j(X)] \leq \mathbb{E}_X[l_j(X)] + \epsilon$, and averaging the same inequalities over the sample,

$$\frac{1}{n}\sum_{i\leq n} l_j(X_i) \leq \frac{1}{n}\sum_{i\leq n} f(X_i) \leq \frac{1}{n}\sum_{i\leq n} u_j(X_i)\ .$$

Proof, continued

Subtracting $\mathbb{E}_X[f(X)]$ and using the two displayed chains, we obtain

$$\frac{1}{n}\sum_{i\leq n} l_j(X_i) - \mathbb{E}_X[l_j(X)] - \epsilon \leq \frac{1}{n}\sum_{i\leq n} f(X_i) - \mathbb{E}_X[f(X)] \leq \frac{1}{n}\sum_{i\leq n} u_j(X_i) - \mathbb{E}_X[u_j(X)] + \epsilon\,.$$

Since the bound only depends on f through j, taking the supremum over $\mathcal{F}$ yields

$$\sup_{f\in\mathcal{F}} \left| \frac{1}{n}\sum_{i\leq n} f(X_i) - \mathbb{E}_X[f(X)] \right| \leq \max_{g\in\{l_1,u_1,\ldots,l_N,u_N\}} \left| \frac{1}{n}\sum_{i\leq n} g(X_i) - \mathbb{E}_X[g(X)] \right| + \epsilon\,.$$

The maximum is over $2N$ fixed integrable functions. By the law of large numbers, each term tends to 0 almost surely, and a maximum of finitely many sequences tending to 0 tends to 0. Hence, $\limsup_n \mathbb{P}(\sup_{\mathcal{F}} |\cdot| \geq 2\epsilon) = 0$ for every $\epsilon > 0$, which concludes the proof.

Corollary: Continuously parametrized functions over metric spaces.

Proposition. Let $\mathcal{F} = \{g_\theta; \theta \in \Theta\}$ with $g_\theta : \mathbb{R}^d \to \mathbb{R}$, and assume

1. Θ is a compact metric space,
2. $\theta \mapsto g_\theta(x)$ is continuous for every x,
3. $\int \sup_{\theta \in \Theta} |g_\theta(x)| d\mathcal{L}(x) < \infty$.

Then $\mathcal{F}$ is $\mathcal{L}$-Glivenko-Cantelli.

The three hypotheses are those that a model satisfies in practice, namely a compact parameter set, a criterion continuous in the parameter, and one integrable envelope.

Proof, building the brackets

Let us fix $\epsilon > 0$. For $\theta \in \Theta$ and $N \in \mathbb{N}$, let $B_{\theta,N}$ be the open ball of radius $1/N$ around θ in Θ, and

$$\tilde{l}_{\theta,N}(x) = \inf_{\tilde{\theta} \in B_{\theta,N}} g_{\tilde{\theta}}(x) \ ,$$
$$\tilde{u}_{\theta,N}(x) = \sup_{\tilde{\theta} \in B_{\theta,N}} g_{\tilde{\theta}}(x) \ .$$

For fixed x, $\tilde{u}_{\theta,N}(x) - \tilde{l}_{\theta,N}(x) \to 0$ as $N \to \infty$ by continuity of $\theta \mapsto g_\theta(x)$ at θ. Since the difference is dominated by $2 \sup_{\theta \in \Theta} |g_\theta|$, which is integrable by hypothesis 3, dominated convergence yields

$$\int \left| \tilde{u}_{\theta,N} - \tilde{l}_{\theta,N} \right| d\mathcal{L} \xrightarrow[N \to \infty]{} 0 \ .$$

We choose N_θ with $\int |\tilde{u}_{\theta,N_\theta} - \tilde{l}_{\theta,N_\theta}| d\mathcal{L} \leq \epsilon$, and fix it.

Proof, a finite subcover

The balls $B_{\theta,N_\theta}, \theta \in \Theta$, form an open cover of Θ, and by compactness, finitely many of them suffice:

$$\Theta \subset \bigcup_{j \leq m} B_{\theta_j, N_{\theta_j}} \ .$$

We set $l_j = \tilde{l}_{\theta_j,N_{\theta_j}}$ and $u_j = \tilde{u}_{\theta_j,N_{\theta_j}}$. Every $\theta \in \Theta$ lies in some $B_{\theta_j,N_{\theta_j}}$, and then $l_j \leq g_\theta \leq u_j$ by construction, so that the m brackets $[l_j, u_j]$ cover $\mathcal{F}$ and each has $L^1(\mathcal{L})$ width at most ϵ.

Hence $N_{[]}(\mathcal{F}, L^1(\mathcal{L}), \epsilon) \leq m < \infty$ for every $\epsilon > 0$, and an application of the previous theorem concludes the proof.

Role of compactness. It only appears here, to pass from a cover indexed by Θ to a finite subcover, which is the role it played in the well separation lemma of Part II, and the reason why both appear in the maximum likelihood theorem.

Consistency of maximum likelihood

We now apply Parts II and III to our running example, and prove that maximum likelihood is consistent.

Theorem. Let $\{\mathcal{L}_\theta; \theta \in \Theta\}$ have densities f_θ with respect to Lebesgue measure, and let $X_1, X_2, \ldots$ be i.i.d. with density f_{θ_0}. Assume

1. Θ is compact in $\mathbb{R}^p$,
2. $f_\theta(x) > 0$ for every θ and x,
3. $\theta \mapsto f_\theta(x)$ is continuous on Θ for every x,
4. $\int \sup_{\theta \in \Theta} |\log f_\theta(x)| f_{\theta_0}(x) dx < \infty$,
5. **identifiability**: $\mathcal{L}_\theta \neq \mathcal{L}_{\theta_0}$ for every $\theta \neq \theta_0$.

Then the maximum likelihood estimator satisfies $\hat{\theta}_n \xrightarrow{\mathbb{P}} \theta_0$.

Plan of the proof

Maximum likelihood is the M-estimator with $M_n(\theta) = \frac{1}{n}\sum_{i\leq n} \log f_\theta(X_i)$, whose pointwise limit is $M(\theta) = \mathbb{E}_X[\log f_\theta(X)]$ by the law of large numbers. We apply the consistency theorem for M estimators, whose three hypotheses are the three steps of the proof.

1. **Uniform convergence**, i.e. $\{\log f_\theta; \theta \in \Theta\}$ is $\mathcal{L}_{\theta_0}$-Glivenko-Cantelli. This is where Part III comes into play, since items 1 to 4 yield the hypotheses of the corollary on continuously parametrized functions.

2. **Well separation of θ_0.** Identifiability (item 5) makes θ_0 the unique maximizer of M, and the lemma of Part II turns it into well separation, with compactness (item 1) and the continuity of M.

3. **Near maximization**, which holds by definition of $\hat{\theta}_n$.

Proof, uniform convergence

For the first step, we apply the corollary on continuously parametrized functions with $g_\theta = \log f_\theta$. Its hypothesis 1 is item 1, its hypothesis 2 follows from items 2 and 3, and its hypothesis 3 is item 4, since $d\mathcal{L}_{\theta_0}(x) = f_{\theta_0}(x)dx$. Hence, $\{\log f_\theta; \theta \in \Theta\}$ is $\mathcal{L}_{\theta_0}$-Glivenko-Cantelli, which means that

$$\sup_{\theta \in \Theta} \left| \frac{1}{n} \sum_{i \leq n} \log f_\theta(X_i) - \mathbb{E}_X[\log f_\theta(X)] \right| \xrightarrow{\mathbb{P}} 0 \ .$$

We thus obtain hypothesis 1 of the consistency theorem for M estimators, which concludes the first step.

Proof, the separation

For the second step, we first show that identifiability makes θ_0 the unique maximizer of M, using the inequality $\log t \leq 2(\sqrt{t} - 1)$ for $t > 0$. For $\theta \neq \theta_0$, we have

$$
\begin{aligned}
M(\theta) - M(\theta_0) &= \int \log\left(\frac{f_\theta(x)}{f_{\theta_0}(x)}\right) f_{\theta_0}(x)dx && \text{by definition of } M \\
&\leq 2\int \left(\sqrt{\frac{f_\theta(x)}{f_{\theta_0}(x)}} - 1\right) f_{\theta_0}(x)dx && \text{by the inequality above} \\
&= -\int \left(\sqrt{f_\theta(x)} - \sqrt{f_{\theta_0}(x)}\right)^2 dx\,, && \text{since } \textstyle\int f_\theta = \int f_{\theta_0} = 1
\end{aligned}
$$

which is negative by item 5. Hence, θ_0 is the unique maximizer of M.

M is continuous on Θ by dominated convergence (items 2 and 3 yield the continuity of the integrand and item 4 the domination). With item 1, the lemma of Part II shows that θ_0 is a well separated maximizer. Since the third step holds by definition, the consistency theorem concludes the proof.

Relaxing identifiability: The risk gap of maximum likelihood

Without identifiability, the first step alone still says something about the estimated law $\mathcal{L}_{\hat{\theta}_n}$. Since maximum likelihood is the empirical risk minimizer for the log loss, stage one controls its risk gap, which we now identify.

Proposition. If the model is well specified, then, with $R(\theta) = \mathbb{E}_X[-\log f_\theta(X)]$, the risk gap of the log loss is a relative entropy:

$$R(\theta) - \inf_{\theta' \in \Theta} R(\theta') = \mathrm{KL}(\mathcal{L}_{\theta_0} \| \mathcal{L}_\theta) \ .$$

Proof. We have

$$\begin{aligned} R(\theta) - R(\theta_0) &= \mathbb{E}_X \left[\log \frac{f_{\theta_0}(X)}{f_\theta(X)} \right] \\ &= \mathrm{KL}(\mathcal{L}_{\theta_0} \| \mathcal{L}_\theta) \\ &\geq 0 \ , \qquad \text{by Jensen's inequality} \end{aligned}$$

Consistency of the estimated law

Plugging this gap into empirical risk minimization as stage one, with Δ_n the uniform deviation of the class $\{\log f_\theta ; \theta \in \Theta\}$, we obtain

$$\mathrm{KL}(\mathcal{L}_{\theta_0} \| \mathcal{L}_{\hat{\theta}_n}) \leq 2\Delta_n + o_{\mathbb{P}}(1) \ .$$

The estimated law is consistent in relative entropy as soon as this class is $\mathcal{L}_{\theta_0}$-Glivenko-Cantelli, without identifiability, compactness or separation. In particular, fitting a non-identifiable model still estimates something, since the predicted distribution converges and only the parameter is meaningless.

Role of identifiability. It only serves to pass from the law to the parameter, in the second step of the proof.

Part IV. Limit laws

Beyond consistency

Parts II and III prove that $\hat{\theta}_n \xrightarrow{\mathbb{P}} \theta_0$. However, consistency only says where the estimator goes, and nothing about the size of the error $\hat{\theta}_n - \theta_0$ at a given n.

The empirical mean as a guide. For $\bar{X}_n$, consistency is the law of large numbers, and the central limit theorem goes one step further. For real X with mean μ and variance σ^2, it states that

$$\sqrt{n}(\bar{X}_n - \mu) \xrightarrow{d} \mathcal{N}(0, \sigma^2) \ .$$

Hence, the error is of order $\sigma/\sqrt{n}$, and $[\bar{X}_n \pm 1.96\sigma/\sqrt{n}]$ contains μ with probability going to 0.95.

Goal of this part. We look for the analogue of the central limit theorem for $\hat{\theta}_n$, called **asymptotic normality**, i.e. for a covariance matrix Σ such that

$$\sqrt{n}(\hat{\theta}_n - \theta_0) \xrightarrow{d} \mathcal{N}(0, \Sigma) \ .$$

The heuristic, linearization

Unlike $\bar{X}_n$, the estimator $\hat{\theta}_n$ is not an average. For the method of moments of the previous lecture, it was an explicit function $e^{-1}(e_n)$ of an average, and the delta method concluded. In general, $\hat{\theta}_n$ is only defined implicitly by $Z_n(\hat{\theta}_n) = 0$, and the idea is to linearize Z_n around θ_0 to come back to averages.

Let us take $Z_n(\theta) = \frac{1}{n}\sum_{i \leq n} z(X_i, \theta)$ with X_i i.i.d. and $\mathbb{E}_X[z(X, \theta_0)] = 0$. Since $\hat{\theta}_n \xrightarrow{\mathbb{P}} \theta_0$ by Parts II and III, we have

$$\begin{aligned} 0 &= Z_n(\hat{\theta}_n) && \text{by definition of } \hat{\theta}_n \\ &\approx Z_n(\theta_0) + (JZ_n)(\theta_0)(\hat{\theta}_n - \theta_0)\ . && \text{by a first order expansion at } \theta_0 \end{aligned}$$

Solving for the error and multiplying by $\sqrt{n}$, we obtain

$$\sqrt{n}(\hat{\theta}_n - \theta_0) \approx -((JZ_n)(\theta_0))^{-1}\sqrt{n}Z_n(\theta_0)\ ,$$

where $(JZ_n)(\theta_0)$ and $Z_n(\theta_0)$ are both averages of i.i.d. terms.

The heuristic, the limit law

Each of the two averages is handled by a classical theorem.

Law of large numbers. $(JZ_n)(\theta_0) \xrightarrow{\mathbb{P}} \mathbb{E}_X[J_z(X, \theta_0)] =: V$, and if V is invertible, continuous mapping inverts it.

Central limit theorem. $\sqrt{n} Z_n(\theta_0) \xrightarrow{d} \mathcal{N}(0, \mathrm{Cov}_X(z(X, \theta_0)))$, where no centering is needed since $\mathbb{E}_X[z(X, \theta_0)] = 0$.

Slutsky. Combining the two, we obtain asymptotic normality,

$$\sqrt{n}(\hat{\theta}_n - \theta_0) \xrightarrow{d} \mathcal{N}\left(0, V^{-1} \mathrm{Cov}_X(z(X, \theta_0)) V^{-\top}\right) \ .$$

Limits of the heuristic

Warning. The expansion requires $\theta \mapsto z(x, \theta)$ to be differentiable for each fixed x. However, for the empirical median, $z(x, \theta) = \operatorname{sign}(\theta - x)$ is not differentiable at $\theta = x$, and not even continuous there, so that the main example of this lecture is out of the reach of the heuristic.

What survives. Only $\theta \mapsto \mathbb{E}_X[z(X, \theta)]$ needs to be smooth. For the median, this expectation is $2F_X(\theta) - 1$, which is as smooth as the density. In other words, averaging over x repairs what happens at each fixed x.

What replaces the remainder control. The heuristic implicitly controls a Taylor remainder at the random point $\hat{\theta}_n$. We replace it by a uniform bound over a shrinking neighbourhood of θ_0, obtained with the maximal inequality of the next slide, and the same bound yields the limit law of the value.

The main tool: the maximal inequality

Theorem. Let $X_1, X_2, \ldots$ be i.i.d. of law $\mathcal{L}$ on $\mathbb{R}^d$ and let $\mathcal{F}$ be a class of functions $\mathbb{R}^d \to \mathbb{R}$ with an **envelope** F, meaning $|f| \leq F$ $\mathcal{L}$-almost everywhere for every $f \in \mathcal{F}$, and $\mathbb{E}_X[F(X)^2] < \infty$. Then, denoting $\mathbb{G}_n f = \frac{1}{\sqrt{n}} \sum_{i \leq n} (f(X_i) - \mathbb{E}_X[f(X)])$, we have

$$\mathbb{E}^\star \left[\sup_{f \in \mathcal{F}} \mathbb{G}_n f \right] \leq C_{\mathrm{MI}} \int_0^{\sqrt{\mathbb{E}_X[F(X)^2]}} \sqrt{\log N_{[]}(\mathcal{F}, L^2(\mathcal{L}), \epsilon)} d\epsilon \ ,$$

for a universal constant C_{MI}.

Proof omitted, see van der Vaart, Corollary 19.35. Here, $\mathbb{E}^\star$ is the outer expectation, since a supremum over an uncountable class need not be measurable.

Absolute values add a 1. We have $\sup_f |\mathbb{G}_n f| = \sup_{g \in \mathcal{F} \cup (-\mathcal{F})} \mathbb{G}_n g$, and negating a bracket keeps its width, so bracketing numbers at most double. Hence, the theorem on $\mathcal{F} \cup (-\mathcal{F})$ yields the same bound with $\sqrt{1 + \log N_{[]}}$, as $\log 2 \leq 1$. **We use this form below.**

Sketch of the proof

We sketch where the bound on $\mathbb{E}^\star[\sup_{f\in\mathcal{F}} \mathbb{G}_n f]$, the left side of the theorem, comes from.

A single scale. For a bracket $l \le f \le u$, the Glivenko-Cantelli argument of Part III yields

$$\frac{1}{n}\sum_{i\le n} f(X_i) - \mathbb{E}_X[f(X)] \le \underbrace{\frac{1}{n}\sum_{i\le n} u(X_i) - \mathbb{E}_X[u(X)]}_{\text{fluctuation}} + \underbrace{\|u-l\|_{L^1(\mathcal{L})}}_{\text{bias} \le \epsilon}.$$

Without $\sqrt{n}$, in Part III. The fluctuation tends to 0 by the law of large numbers, so that only the bias remains, which is an expectation, hence the L^1 width.

With $\sqrt{n}$, here. The fluctuation becomes $\mathbb{G}_n u$, of order $\|u\|_{L^2(\mathcal{L})}$, and the bias becomes $\sqrt{n}\epsilon$, which forces $\epsilon \lesssim n^{-1/2}$ and far too many brackets for a union bound.

Chaining. We approximate f at every scale $\epsilon_k = 2^{-k}\epsilon_0$ at once and telescope, $f = f_0 + \sum_{k\ge 1}(f_k - f_{k-1})$. The bias terms cancel between consecutive scales, and only fluctuations of increments remain.

Bernstein's inequality

Each increment $h = f_k - f_{k-1}$ appears through the centered average $\mathbb{G}_n h$, whose tail we control as follows.

Lemma (Bernstein's inequality). Let h be bounded with $\mathbb{E}_X[h(X)] = 0$. Then, for every $t > 0$,

$$\mathbb{P}\left(|\mathbb{G}_n h| \geq t\right) \leq 2 \exp\left(-\frac{t^2/2}{\|h\|^2_{L^2(\mathcal{L})} + t\|h\|_\infty/(3\sqrt{n})}\right).$$

We omit the proof, which relies on the Chernoff method (Boucheron, Lugosi and Massart, Section 2.8).

Variance term. $\|h\|^2_{L^2(\mathcal{L})} = \mathrm{Var}_X(h(X))$ is the variance of $\mathbb{G}_n h$ and of its gaussian limit.

Range term. $t\|h\|_\infty/(3\sqrt{n})$ accounts for the largest value of h, and vanishes as $n \to \infty$.

Two regimes. If $t \leq 3\sqrt{n}\|h\|^2_{L^2(\mathcal{L})}/\|h\|_\infty$, the variance term dominates and the bound is at most the gaussian tail $2\exp(-t^2/(4\|h\|^2_{L^2(\mathcal{L})}))$. Beyond, the range term dominates and the tail is only exponential.

Origin of the L^2 control

Since the threshold of the gaussian regime grows like $\sqrt{n}$, Bernstein's inequality makes $\mathbb{G}_n h$ approximately sub-gaussian for large n, with variance proxy $\|h\|^2_{L^2(\mathcal{L})}$. Hence, a maximum over N such increments satisfies

$$\mathbb{E}_S\left[\max_{j\le N} \mathbb{G}_n h_j\right] \lesssim \max_{j\le N} \|h_j\|_{L^2(\mathcal{L})}\sqrt{\log N}\ .$$

Reading the bound. A maximum of N increments grows like their largest standard deviation times $\sqrt{\log N}$, and the standard deviation of $\mathbb{G}_n h$ is $\|h\|_{L^2(\mathcal{L})}$. This is where the L^2 norm comes from.

L^1 against L^2. The L^1 norm of Part III measured the bias of a bracket, which chaining cancels, whereas the L^2 norm measures the fluctuations of the increments, which chaining keeps.

The entropy integral, resolution by resolution

At resolution k, with $N_k = N_{[]}(\mathcal{F}, L^2(\mathcal{L}), \epsilon_k)$, the increments $f_k - f_{k-1}$ form a set $\mathcal{H}_k$ of at most N_k^2 functions of L^2 norm at most $3\epsilon_k$. At the coarsest resolution $\epsilon_0 \asymp \sqrt{\mathbb{E}_X[F(X)^2]}$, the single bracket $[-F, F]$ suffices. Hence,

$$\mathbb{E}^\star\left[\sup_{f\in\mathcal{F}} \mathbb{G}_n f\right] \le \underbrace{\mathbb{E}_S[\mathbb{G}_n f_0]}_{=0 \text{ at resolution } 0} + \sum_{k\ge 1} \underbrace{\mathbb{E}_S\left[\max_{h\in\mathcal{H}_k} \mathbb{G}_n h\right]}_{\lesssim \epsilon_k\sqrt{\log N_k} \text{ at resolution } k}$$

$$\lesssim \int_0^{\epsilon_0} \sqrt{\log N_{[]}(\mathcal{F}, L^2(\mathcal{L}), \epsilon)}\, d\epsilon \,. \qquad \text{since } \epsilon_k = 2(\epsilon_k - \epsilon_{k+1})$$

Reading the sum. Each resolution contributes the size ϵ_k of its increments times the square root of the logarithm of their number, and the entropy integral adds up these contributions.

Exercise, the empirical distribution function

Exercise. Let $X_1, X_2, \ldots$ be i.i.d. $\mathrm{Unif}([0,1])$ and $\mathcal{F} = \{\mathbf{1}_{(-\infty,t]}; t \in [0,1]\}$. Bound $\mathbb{E}^\star[\sup_{f\in\mathcal{F}} \frac{1}{\sqrt{n}} |\sum_{i\le n}(f(X_i) - \mathbb{E}_X[f(X)])|]$.

Solution. The constant function $F \equiv 1$ is an envelope, so that $\mathbb{E}_X[F(X)^2] = 1$, and the exercise of Part III yields $N_{[]}(\mathcal{F}, L^2(\mathcal{L}), \epsilon) \lesssim \epsilon^{-2}$, hence

$$
\begin{aligned}
&\int_0^1 \sqrt{1 + \log N_{[]}(\mathcal{F}, L^2(\mathcal{L}), \epsilon)}\, d\epsilon \\
&\lesssim \int_0^1 \left(1 + \sqrt{2\log(1/\epsilon)}\right) d\epsilon && \text{since } \sqrt{1+x} \le 1 + \sqrt{x} \\
&= 1 + \sqrt{2}\int_0^\infty \sqrt{u} e^{-u} du && \text{with } \epsilon = e^{-u} \\
&= 1 + \frac{\sqrt{2\pi}}{2}\,. && \text{since } \sqrt{2}\Gamma(3/2) = \sqrt{2\pi}/2
\end{aligned}
$$

As a result, the empirical distribution function deviates from the true one by $O_{\mathbb{P}}(n^{-1/2})$ uniformly in t, which is the Kolmogorov and Smirnov rate.

Rate of the uniform deviation

Since $\sqrt{n}\Delta_n = \sup_{f \in \mathcal{F}} |\mathbb{G}_n f|$, the maximal inequality with absolute values yields a rate for the uniform deviation of Part III,

$$\mathbb{E}_S[\Delta_n] \leq \frac{C_{\mathrm{MI}}}{\sqrt{n}} \int_0^{\sqrt{\mathbb{E}_X[F(X)^2]}} \sqrt{1 + \log N_{[]}(\mathcal{F}, L^2(\mathcal{L}), \epsilon)}\, d\epsilon \, .$$

The parametric rate. As soon as the integral is finite, Δ_n is of order $n^{-1/2}$. For instance, a class with $\log N_{[]}(\epsilon) \lesssim p \log(1/\epsilon)$ has an integral of order $\sqrt{p}$, so that Δ_n is of order $\sqrt{p/n}$, and the indicators of the exercise are of this kind.

A question. The integral is finite when the integrand is integrable close to 0, roughly when $\log N_{[]}(\epsilon)$ grows more slowly than ϵ^{-2}. What happens when it is not?

Entropy of Lipschitz functions

To answer the question, we need a class whose entropy grows faster than ϵ^{-2}. The typical one is the nonparametric class $\mathcal{F} = \{f : [0,1]^d \to \mathbb{R}; \|f\|_\infty \le 1, \|f\|_{\mathrm{Lip}} \le 1\}$.

Lemma. There is C_d, depending only on d, such that, for every $\epsilon \in (0,1]$,

$$\log N(\mathcal{F}, \|\cdot\|_\infty, \epsilon) \le C_d \epsilon^{-d} .$$

Proof. We cut $[0,1]^d$ into $K \le (2\sqrt{d}/\epsilon)^d$ cubes of side at most $\epsilon/\sqrt{d}$, with centers $x_1, \dots, x_K$ in lexicographic order, so that each x_j with $j \ge 2$ is adjacent to an earlier center. To $f \in \mathcal{F}$, we associate the function g equal on the j-th cube to the multiple q_j of ϵ closest to $f(x_j)$. Since every point of that cube is within $\epsilon/2$ of x_j, we get $|f - g| \le \epsilon/2 + \epsilon/2 = \epsilon$ there.

It remains to count the functions g. There are at most $4/\epsilon$ choices for q_1. For adjacent cubes, $|q_j - q_{j'}| \le \epsilon/2 + \epsilon + \epsilon/2 = 2\epsilon$, so that each next q_j has at most 5 choices given the earlier ones. Hence, the logarithm of the number of functions g is at most $\log(4/\epsilon) + K \log 5 \le (4 + (2\sqrt{d})^d \log 5)\epsilon^{-d}$, which concludes the proof.

Divergence of the entropy integral

For the Lipschitz class, the lemma bounds the integrand by $\sqrt{1 + C_d \epsilon^{-d}}$, which is integrable at 0 only for $d < 2$. For $d \geq 2$, the maximal inequality thus yields an infinite bound, and we stop the chaining at a scale δ instead.

Proposition (truncated maximal inequality). Let $\mathcal{F}$ have an envelope F with $\mathbb{E}_X[F(X)^2] < \infty$. Then, for every $\delta > 0$ with $4\delta \leq \sqrt{\mathbb{E}_X[F(X)^2]}$,

$$\mathbb{E}^\star[\Delta_n] \leq C \left(\delta + \frac{1}{\sqrt{n}} \int_\delta^{\sqrt{\mathbb{E}_X[F(X)^2]}} \sqrt{1 + \log N(\mathcal{F}, \|\cdot\|_\infty, \epsilon)} d\epsilon \right) ,$$

for a universal constant C.

Reading it. The term δ accounts for the scales below δ, where we approximate in supremum norm without chaining, and the integral for the chaining above δ. The cutoff δ is free, and we optimize it.

Polynomial entropy. When $\log N(\mathcal{F}, \|\cdot\|_\infty, \epsilon) \lesssim \epsilon^{-\alpha}$, as for the Lipschitz class with $\alpha = d$, optimizing in δ leads to three regimes, depending on α.

Proof of the truncated maximal inequality

Let $\eta = 4\delta$. Picking one element of $\mathcal{F}$ in each ball of a supremum norm $\eta/2$-cover, we get a finite $\mathcal{F}_\eta \subset \mathcal{F}$ with at most $N(\mathcal{F}, \|\cdot\|_\infty, 2\delta)$ elements, such that every $f \in \mathcal{F}$ is within η of some $g \in \mathcal{F}_\eta$. Since replacing f by g changes both the average and the expectation by at most η, we have

$$\Delta_n \le \sup_{g \in \mathcal{F}_\eta} \left| \frac{1}{n} \sum_{i \le n} g(X_i) - \mathbb{E}_X[g(X)] \right| + 8\delta \; .$$

The maximal inequality with absolute values applies to $\mathcal{F}_\eta$, which has envelope F. Its bracketing numbers at scale ϵ are at most $N(\mathcal{F}, \|\cdot\|_\infty, 2\delta)$, since each g is a bracket of width 0, and at most $N(\mathcal{F}, \|\cdot\|_\infty, \epsilon/2)$, since a ball of radius $\epsilon/2$ is a bracket of L^2 width ϵ.

We use the first bound for $\epsilon \le 4\delta$ and the second above. Since $N(\mathcal{F}, \|\cdot\|_\infty, \epsilon/2) \ge N(\mathcal{F}, \|\cdot\|_\infty, 2\delta)$ for $\epsilon \le 4\delta$, the part below 4δ is at most twice the integral over $[2\delta, 4\delta]$, and the substitution $\epsilon = 2u$ yields the result with $C = \max(8, 6C_{\mathrm{MI}})$, which concludes the proof.

Nonparametric rates, the three regimes

Let $\mathcal{F}$ have envelope 1 and satisfy $\log N(\mathcal{F}, \|\cdot\|_\infty, \epsilon) \lesssim \epsilon^{-\alpha}$ for some $\alpha > 0$. The truncated maximal inequality yields $\mathbb{E}^\star[\Delta_n] \lesssim \delta + n^{-1/2} \int_\delta^1 \epsilon^{-\alpha/2} d\epsilon$, and optimizing in δ leads to

entropy exponent	$\int_\delta^1 \epsilon^{-\alpha/2} d\epsilon$	best δ	rate of $\mathbb{E}^\star[\Delta_n]$
$\alpha < 2$	converges at 0	0	$n^{-1/2}$
$\alpha = 2$	$\log(1/\delta)$	$n^{-1/2}$	$n^{-1/2} \log n$
$\alpha > 2$	$\delta^{1-\alpha/2}$	$n^{-1/\alpha}$	$n^{-1/\alpha}$

Reading the table. Below $\alpha = 2$, we recover the parametric rate. Above, the rate $n^{-1/\alpha}$ is slower, and the larger the class, the slower the rate.