

Bachelor Thesis
Decoding the Color Code

Clemens Schumann,
Advised by Peter-Jan Derks

November 22, 2022

Contents

1	Introduction to the Algebra	3
1.1	Schroedinger and Heisenberg picture	3
1.1.1	Schroedinger picture	3
2	Conclusion	6
3	Appendix	6

1 Introduction to the Algebra

1.1 Schroedinger and Heisenberg picture

1.1.1 Schroedinger picture

In the Schroedinger picture, we focus on the time evolution of states:

$$|\psi\rangle = |\psi\rangle(t) \quad (1)$$

In this picture we can introduce quantum circuit diagram notation, whereby:

- States progress in time along horizontal parallel lines
- Time goes from left to right
- Gates denoted X,Y,Z are the pauli matrices $\sigma_x, \sigma_y, \sigma_z$
- Gates can act on one or multiple qubits, whereby an X gate on qubit 1 in a 3-qubit system should be interpreted as:
 $(X \otimes \mathbb{I} \otimes \mathbb{I})(|\psi_1\rangle \otimes |\psi_2\rangle \otimes |\psi_3\rangle)$




Figure 1: A Quantum Circuit, where $|0\rangle$ is the +1 eigenstate in σ_z -basis

In the quantum circuit depicted in figure 1 the input state can be written as $|\psi_{control}\rangle \otimes |0\rangle \otimes |\psi_{target}\rangle$ and the measurement in the first timestep can be expressed as $\mathbb{I} \otimes X \otimes X$.

To simplify calculation we can write $|\psi_{target}\rangle$ as $\alpha|0\rangle + \beta|1\rangle$

The input state is thus: $|\psi_{control}\rangle \otimes |0\rangle \otimes (\alpha|0\rangle + \beta|1\rangle)$.

The initial state $|\phi_{t=0}\rangle = |\psi_{control}\rangle \otimes |\psi_{ancilla}\rangle \otimes |\psi_{target}\rangle$

where

$$|\psi_{control}\rangle = \alpha|0\rangle + \beta|1\rangle$$

$$|\psi_{ancilla}\rangle = |0\rangle$$

$$|\psi_{target}\rangle = \gamma|0\rangle + \delta|1\rangle$$

therefore:

$$|\phi_{t=0}\rangle = \alpha (\gamma|000\rangle + \delta|001\rangle) + \beta (\gamma|100\rangle + \delta|101\rangle) \quad (2)$$

If the first measurement result is +1, the state becomes:

$$\begin{aligned} |\phi_{t=1}^+\rangle &= \frac{1}{2} (\mathbb{I} \otimes \mathbb{I} \otimes \mathbb{I} + \mathbb{I} \otimes X \otimes X) |\phi_{t=0}\rangle \\ &= \alpha (\gamma (|000\rangle + |011\rangle) + \delta (|001\rangle + |010\rangle)) \\ &\quad + \beta (\gamma (|100\rangle + |111\rangle) + \delta (|101\rangle + |110\rangle)) \end{aligned}$$

if the result is -1, it becomes:

$$\begin{aligned} |\phi_{t=1}^-\rangle &= \frac{1}{2} (\mathbb{I} \otimes \mathbb{I} \otimes \mathbb{I} - \mathbb{I} \otimes X \otimes X) |\phi_{t=0}\rangle \\ &= \alpha (\gamma (|000\rangle - |011\rangle) + \delta (|001\rangle - |010\rangle)) \\ &\quad + \beta (\gamma (|100\rangle - |111\rangle) + \delta (|101\rangle - |110\rangle)) \end{aligned}$$

In the case of the +1 Measurement $\rightarrow a=0$:

$$\begin{aligned} |\phi_{t=2}^{++}\rangle &= \frac{1}{2} (\mathbb{I} \otimes \mathbb{I} \otimes \mathbb{I} + Z \otimes Z \otimes \mathbb{I}) |\phi_{t=1}^+\rangle \\ &= (|000\rangle\langle 000| + |001\rangle\langle 001| + |110\rangle\langle 110| + |111\rangle\langle 111|) |\phi_{t=1}^+\rangle \\ &= \alpha (\gamma|000\rangle + \delta|001\rangle) + \beta (\gamma|111\rangle + \delta|110\rangle) \end{aligned}$$

$$\begin{aligned} |\phi_{t=2}^{+-}\rangle &= \frac{1}{2} (\mathbb{I} \otimes \mathbb{I} \otimes \mathbb{I} - Z \otimes Z \otimes \mathbb{I}) |\phi_{t=1}^+\rangle \\ &= (|010\rangle\langle 010| + |011\rangle\langle 011| + |100\rangle\langle 100| + |101\rangle\langle 101|) |\phi_{t=1}^+\rangle \\ &= \alpha (\gamma|011\rangle + \delta|010\rangle) + \beta (\gamma|100\rangle + \delta|101\rangle) \end{aligned}$$

In the case of the -1 Measurement $\rightarrow a=1$:

$$\begin{aligned} |\phi_{t=2}^{-+}\rangle &= \frac{1}{2} (\mathbb{I} \otimes \mathbb{I} \otimes \mathbb{I} + Z \otimes Z \otimes \mathbb{I}) |\phi_{t=1}^{-}\rangle \\ &= \alpha (\gamma|000\rangle + \delta|001\rangle) - \beta (\gamma|111\rangle + \delta|110\rangle) \end{aligned}$$

$$\begin{aligned} |\phi_{t=2}^{--}\rangle &= \frac{1}{2} (\mathbb{I} \otimes \mathbb{I} \otimes \mathbb{I} - Z \otimes Z \otimes \mathbb{I}) |\phi_{t=1}^{-}\rangle \\ &= -\alpha (\gamma|011\rangle + \delta|010\rangle) + \beta (\gamma|100\rangle + \delta|101\rangle) \end{aligned}$$

Now the applied measurement is $\mathbb{I} \otimes X \otimes \mathbb{I}$, which means:

$$\begin{aligned} |\phi_{t=3}^{+++}\rangle &= \frac{1}{2} (\mathbb{I} \otimes \mathbb{I} \otimes \mathbb{I} + \mathbb{I} \otimes X \otimes \mathbb{I}) |\phi_{t=2}^{++}\rangle \\ &= ((|010\rangle + |000\rangle)\langle 000| + (|011\rangle + |001\rangle)\langle 001| \\ &\quad + (|000\rangle + |010\rangle)\langle 010| + (|001\rangle + |011\rangle)\langle 011| \\ &\quad + (|110\rangle + |100\rangle)\langle 100| + (|111\rangle + |101\rangle)\langle 101| \\ &\quad + (|100\rangle + |110\rangle)\langle 110| + (|101\rangle + |111\rangle)\langle 111|) |\phi_{t=2}^{++}\rangle \\ &= \alpha (\gamma(|000\rangle + |010\rangle) + \delta(|011\rangle + |001\rangle)) \\ &\quad + \beta (\gamma(|101\rangle + |111\rangle) + \delta(|100\rangle + |110\rangle)) \end{aligned}$$

In this case, a,b and c would each be zero, therefore no further gate would be applied.

As intended, this state is equivalent to $CNOT|\phi_{t=0}\rangle$

2 Conclusion

3 Appendix