

6.2: Binomial Probabilities

Binomial Experiments

Definition

A **binomial experiment** is an experiment satisfying the following four conditions:

- There is a fixed number of trials, denoted n .

Definition

A **binomial experiment** is an experiment satisfying the following four conditions:

- There is a fixed number of trials, denoted n .
- The n trials are independent and repeated under identical conditions.

Definition

A **binomial experiment** is an experiment satisfying the following four conditions:

- There is a fixed number of trials, denoted n .
- The n trials are independent and repeated under identical conditions.
- There are exactly two possible outcomes for each trial. These outcomes can be considered *success* and *failure*.

Definition

A **binomial experiment** is an experiment satisfying the following four conditions:

- There is a fixed number of trials, denoted n .
- The n trials are independent and repeated under identical conditions.
- There are exactly two possible outcomes for each trial. These outcomes can be considered *success* and *failure*.
- For each trial, the probability of success is the same. We denote the probability of success by p and the probability of failure by q . Because each trial results in either success or failure, $p + q = 1$.

Definition

A **binomial experiment** is an experiment satisfying the following four conditions:

- There is a fixed number of trials, denoted n .
- The n trials are independent and repeated under identical conditions.
- There are exactly two possible outcomes for each trial. These outcomes can be considered *success* and *failure*.
- For each trial, the probability of success is the same. We denote the probability of success by p and the probability of failure by q . Because each trial results in either success or failure, $p + q = 1$.

The central problem of a binomial experiment is to find the probability of r successes out of n trials.

Binomial Experiments

Example

Determine if the following experiment is a binomial experiment. If it is not a binomial experiment, explain why.

- Selecting 20 university students and recording their class rank.

Example

Determine if the following experiment is a binomial experiment. If it is not a binomial experiment, explain why.

- Selecting 20 university students and recording their class rank.

This is not a binomial experiment because there are more than two outcomes for the variable.

Example

Determine if the following experiment is a binomial experiment. If it is not a binomial experiment, explain why.

- Selecting 20 university students and recording whether they are on the Dean's list.

Example

Determine if the following experiment is a binomial experiment. If it is not a binomial experiment, explain why.

- Selecting 20 university students and recording whether they are on the Dean's list.

This is a binomial experiment.

Example

Determine if the following experiment is a binomial experiment. If it is not a binomial experiment, explain why.

- Drawing five cards from a standard deck of cards without replacement and recording whether they are red or black.

Example

Determine if the following experiment is a binomial experiment. If it is not a binomial experiment, explain why.

- Drawing five cards from a standard deck of cards without replacement and recording whether they are red or black.

This is not a binomial experiment because the probability of success will change with each draw.

Binomial Experiments

Example

A survey from Teenage Research Unlimited found that 30% of teenage consumers receive their spending money from part-time jobs. We select 10 teenagers at random to determine the probability that exactly 4 of them will have part-time jobs. Find the values p , q , n , and r .

Example

A survey from Teenage Research Unlimited found that 30% of teenage consumers receive their spending money from part-time jobs. We select 10 teenagers at random to determine the probability that exactly 4 of them will have part-time jobs. Find the values p , q , n , and r .

- We will consider having a part-time job a success.

Example

A survey from Teenage Research Unlimited found that 30% of teenage consumers receive their spending money from part-time jobs. We select 10 teenagers at random to determine the probability that exactly 4 of them will have part-time jobs. Find the values p , q , n , and r .

- We will consider having a part-time job a success.
- Since p is the probability of success, the example states that $p = 0.3$.

Example

A survey from Teenage Research Unlimited found that 30% of teenage consumers receive their spending money from part-time jobs. We select 10 teenagers at random to determine the probability that exactly 4 of them will have part-time jobs. Find the values p , q , n , and r .

- We will consider having a part-time job a success.
- Since p is the probability of success, the example states that $p = 0.3$.
- We can compute $q = 1 - p = 0.7$. Recall that q is the probability of failure.

Example

A survey from Teenage Research Unlimited found that 30% of teenage consumers receive their spending money from part-time jobs. We select 10 teenagers at random to determine the probability that exactly 4 of them will have part-time jobs. Find the values p , q , n , and r .

- We will consider having a part-time job a success.
- Since p is the probability of success, the example states that $p = 0.3$.
- We can compute $q = 1 - p = 0.7$. Recall that q is the probability of failure.
- We consider each selected teenager a trial. So $n = 10$.

Example

A survey from Teenage Research Unlimited found that 30% of teenage consumers receive their spending money from part-time jobs. We select 10 teenagers at random to determine the probability that exactly 4 of them will have part-time jobs. Find the values p , q , n , and r .

- We will consider having a part-time job a success.
- Since p is the probability of success, the example states that $p = 0.3$.
- We can compute $q = 1 - p = 0.7$. Recall that q is the probability of failure.
- We consider each selected teenager a trial. So $n = 10$.
- Since we want to consider the probability that exactly 4 of the selected teenagers will have a part-time job, $r = 4$.

Binomial Probability Distribution Formula

Formula

In a binomial experiment, the probability of r successes out of n trials is given by the formula

$$\Pr(r) = \frac{n!}{r!(n-r)!} p^r \cdot q^{n-r} = (C_{n,r}) \cdot p^r \cdot q^{n-r}$$

where p is the probability of success in each trial and q is the probability of failure in each trial.

Binomial Probability Distribution Formula

Example

A survey from Teenage Research Unlimited found that 30% of teenage consumers receive their spending money from part-time jobs. If we select 10 teenagers at random, what is the probability that exactly 4 of them will have part-time jobs?

- In the previous example we found the following values

$$p = 0.3$$

$$q = 0.7$$

$$n = 10$$

$$r = 4$$

Example

A survey from Teenage Research Unlimited found that 30% of teenage consumers receive their spending money from part-time jobs. If we select 10 teenagers at random, what is the probability that exactly 4 of them will have part-time jobs?

- In the previous example we found the following values

$$p = 0.3$$

$$q = 0.7$$

$$n = 10$$

$$r = 4$$

- Using the binomial probability distribution formula

$$\Pr(4) = \frac{10!}{4!(10-4)!} (0.3)^4 (0.7)^{10-4}$$

$$\approx 0.2$$

Binomial Probability Distribution Formula

Binomial Probability Distribution Formula

Example

If a die is rolled 20 times, what is the probability that exactly half of the rolls will land on 3?

Example

If a die is rolled 20 times, what is the probability that exactly half of the rolls will land on 3?

- We begin by noticing that this is a binomial experiment. Although there are six possible values on the die, we consider landing on a 3 a success and anything else a failure.

Example

If a die is rolled 20 times, what is the probability that exactly half of the rolls will land on 3?

- We begin by noticing that this is a binomial experiment. Although there are six possible values on the die, we consider landing on a 3 a success and anything else a failure.
- Next we identify $n = 20$, $r = 10$, $p = 1/6$ and $q = 5/6$.

Example

If a die is rolled 20 times, what is the probability that exactly half of the rolls will land on 3?

- We begin by noticing that this is a binomial experiment. Although there are six possible values on the die, we consider landing on a 3 a success and anything else a failure.
- Next we identify $n = 20$, $r = 10$, $p = 1/6$ and $q = 5/6$.
- Using the binomial probability distribution formula

$$\Pr(\text{Ten 3s}) = \frac{20!}{10!(20-10)!} \left(\frac{1}{6}\right)^{10} \cdot \left(\frac{5}{6}\right)^{20-10}$$

Example

If a die is rolled 20 times, what is the probability that exactly half of the rolls will land on 3?

- We begin by noticing that this is a binomial experiment. Although there are six possible values on the die, we consider landing on a 3 a success and anything else a failure.
- Next we identify $n = 20$, $r = 10$, $p = 1/6$ and $q = 5/6$.
- Using the binomial probability distribution formula

$$\begin{aligned}\Pr(\text{Ten 3s}) &= \frac{20!}{10!(20-10)!} \left(\frac{1}{6}\right)^{10} \cdot \left(\frac{5}{6}\right)^{20-10} \\ &\approx 0.00049\end{aligned}$$

Mean and Standard Deviation

Formula

In a binomial experiment

- $\mu = np$
- $\sigma = \sqrt{npq}$.

Formula

In a binomial experiment

- $\mu = np$
- $\sigma = \sqrt{npq}$.

The mean value μ can be thought of as the **expected number of successes** in the experiment.

Formula

In a binomial experiment

- $\mu = np$
- $\sigma = \sqrt{npq}$.

The mean value μ can be thought of as the **expected number of successes** in the experiment.

Example

If we roll a single die 20 times, how many times can we expect 3 to roll?

Formula

In a binomial experiment

- $\mu = np$
- $\sigma = \sqrt{npq}$.

The mean value μ can be thought of as the **expected number of successes** in the experiment.

Example

If we roll a single die 20 times, how many times can we expect 3 to roll?

- Using the binomial experiment formula for μ , we can expect the number of 3s rolled to be

$$\mu = 20 \cdot \left(\frac{1}{6}\right) = 3.\bar{3}$$

Formula

In a binomial experiment

- $\mu = np$
- $\sigma = \sqrt{npq}$.

The mean value μ can be thought of as the **expected number of successes** in the experiment.

Example

If we roll a single die 20 times, how many times can we expect 3 to roll? Find the standard deviation for the number of 3s rolled.

Formula

In a binomial experiment

- $\mu = np$
- $\sigma = \sqrt{npq}$.

The mean value μ can be thought of as the **expected number of successes** in the experiment.

Example

If we roll a single die 20 times, how many times can we expect 3 to roll? Find the standard deviation for the number of 3s rolled.

- Using the binomial experiment formula for σ , find the standard deviation for the number of 3s rolled to be

$$\sigma = \sqrt{20 \cdot \left(\frac{1}{6}\right) \cdot \left(\frac{5}{6}\right)} = 1.\bar{6}$$