

1. Let R be the set of positive real numbers and define addition, denoted by \oplus , and multiplication, denoted by \otimes , as follows. For every $a, b \in R$, $a \oplus b = ab$, and $a \otimes b = a^{\log b}$. Please prove or disprove (R, \oplus, \otimes) .

axiom -1: since $R \in R^+$, \oplus is closed in R . There is no positive real number for which $ab \in R^+$

axiom 0: since $R \in R^+$ and \otimes is defined as $a \otimes b = a^{\log b}$, starting with the lowest possible value, $a, b = 1$, $1^{\log 1}$ results in 1, which is included in R^+ . as both values approach infinity the result also approaches infinity which is a real number.

axiom 1: addition is commutative, for all values a, b , $a \cdot b = b \cdot a$

axiom 2: addition is associative, for all values a, b, c , $(a \cdot b) \cdot c = a \cdot (b \cdot c)$

axiom 3: there exists an additive identity. since the \otimes is defined as $a \oplus b = ab$, the additive identity is 1. $1 \cdot x = x \cdot 1 = x$.

axiom 4: There is an additive inverse. Since the additive identity is 1, there must be a value for every $x \in R^+$ where $x \cdot y = 1$. This would be the inverse of the number x . For example: $x = 3$, $x \cdot x^{-1} = 3 \cdot 1/3 = 1$

axiom 5: multiplication is commutative. $a \otimes b = a^{\log b}$. This is commutative because if you take \log_a of both sides it results in $\log b = \log_a b^{\log a}$. Breaking apart the right side, $\frac{\log a \cdot \log b}{\log a}$ in which both $\log a$ cancel each other out and you're left with $\log b = \log b$. Since they are equal, \otimes is commutative.

axiom 6: Multiplication is associative. Given a, b, c : $(a \otimes b) \otimes c = a \otimes (b \otimes c)$ on the left, the resultant is $(a^{\log b})^{\log c}$, and on the right side, $(a^{\log b})^{\log c}$. Since these are equal, multiplication is associative

axiom 7: there exists a multiplicative identity. This axiom holds true for the value 10. $(a^{\log 10} = 10^{\log a} = a)$

axiom 8: there exists a multiplicative inverse. since the multiplicative identity is 10, the multiplicative inverse will be a number where $a^{\log b} = 10$ solving for b , $\log b = \log_a 10$, and solving for b , $b = 10^{\log_a 10}$. Since $a \in R^+$, $\log_a 10$ is positive for all values a . This means that 10^{R^+} and will result in a positive real number meaning that for all values a , there exists another number $b \in R^+$ for the multiplicative inverse.

axiom 9: Multiplication distributes over addition. If $x, y, z \in R$, then $x \cdot (y + z) = x \cdot y + x \cdot z$ and $(x + y) \cdot z = z \cdot x + z \cdot y$. With our redefined definition of both addition and multiplication, $x \cdot (y + z) = x \otimes (y \cdot z) = x^{\log y \cdot \log z}$ and on the right side of the equation $x \cdot y + x \cdot z = x^{\log y} \cdot x^{\log z} = x^{\log y \cdot \log z}$. Since they are equal on both sides of addition, axiom 9 holds.

axiom 10: the additive identity and multiplicative identity are not equal. For our redefined \oplus, \otimes the additive and multiplicative identities were 1 and 10 respectively. since $1 \neq 10$ axiom 10 holds.

since all axioms hold, our redefined field (R, \oplus, \otimes) is a field.

2. Denote the set $0,1,2,3$ in Z_4 and define addition, denoted by $+$, and multiplication, denoted by \cdot or juxtaposition, via the following tables:

axiom 8 states that for every number in Z_4 there is a multiplicative inverse where $a \cdot a^{-1} = 1$. 2 from the range Z_4 has no multiplicative inverse because there exists no other number, b , in Z_4 where $a \cdot b = 1$.

4. Please prove or disprove: if $n \in Z^+$, then $n^2 + n + 41$ is prime:

question 4 asserts that if $n \in Z^+$, then for all values n , $n^2 + n + 41$ will be prime. In order to disprove this, there must be some value for n where the output isn't prime. take $n = 40$, this results in $40^2 + 40 + 41 = 1681$. 1681 is not prime because 41 is a factor of 1681. Therefore $n^2 + n + 41$ is not prime for all $n \in Z^+$.