ECG’s Step by Step


2021

Outline

- History, Terminologies and Lead Placement
- Interpretation Approach
- Practice Cases
- EKG Patterns and Buzzwords

 Clickable Links !!

History of the 12 Lead





- Einthoven built first clinical EKG in 1983 earning 1924 Nobel Prize
- In 1954, AHA recommended for standardization of 12-lead EKG.


Terms


1. Wave: +/- deflection from baseline indicating electrical event.
2. Interval: time btn two specific ECG events.
3. Segment: length btn two points at baseline (not +/- amplitude)
4. Complex: combination of multiple waves
5. Point: only one point on ECG,
   - J point, which is where QRS complex ends and ST segment begins

Where The Leads Go









RA – right forearm or wrist
LA – left forearm or wrist
LL – left lower leg, proximal to ankle
RL – right lower leg, proximal to ankle
V1 – 4-th intercostal space, right sternal edge
V2 – 4-th intercostal space, left sternal edge
V3 – midway between V2 and V4
V4 – 5-th intercostal space, mid-clavicular line
V5 – anterior axillary line in straight line with V4
V6 – mid-axillary line in straight line with V4 and V5


Me, interpreting this EEG
Me, realizing it's actually an ECG

Approach (a systematic approach is vital)

- So that you are not distracted by findings or bias!
- Rate (? tachy or brady) + rhythm (? P waves, regularity, P & QRS relationship)

- Intervals (PR, QRS, QT) and axis (? LAD or RAD)
- Chamber abnormality (? LAA and/or RAA, ? LVH and/or RVH)

- QRST changes (? Q waves, poor R-wave progression V1–V6, ST ↑/↓ or T-wave Δs)

Step 1: Rate

If Reg rhythm determined by “Rule of 300”.

- Divide 300 by no of boxes in R-R Interval

OR

- Strip Method
  - R waves x 6

Use **the every-other-beat method** when the rhythm is regular and the rate is fast”



Not with that rhythm.



Step 2: Rhythm — the 5 Parameters

- **R**ate ?
- Is there a single, **R**egular **P** before every Q wave?
  - Sinus Rhythm
- **Q**RS width?
- P waves **R**elated to QRS, or Varying?
  - WAP or MAT
- No P waves
  - Buried* A fib, junctional or ventricular origins






Figure 02.1-1: Is the QRS wide? Normally in adults — the QRS should *not* be more than *half* a large box in duration (= *not more than 0.10 second*).


Normal Sinus Rythm



"If the rhythm is sinus — then P wave should always be upright in lead II (Panel A). Only exception is IF there is dextrocardia or lead misplacement. Therefore — Panel B is not a sinus rhythm"

1. Regular rhythm rate of 60-100 bpm (or age-appropriate rate)
2. All QRS preceded by a normal P wave
3. Normal P wave axis: P waves upright in leads I & II, inverted in aVR
4. The PR interval remains constant
5. QRS complexes < 100ms wide (unless co-existent interventricular conduction delay present)



Arrhythmia
Bradyarrythmia (<60 bpm)
• Sinus bradycardia
• Sick sinus syndrome
• Escape rhythms
– Junctional
– Ventricular
Conduction delay
• AV nodal conduction blocks
– 1°, 2°, 3°
• Fascicular block
• Bundle branch block
Tachyarrhythmia (> 100 bpm)
Regular
Narrow complex
• Supraventricular tachycardia
• Atrial flutter
• AVNRT
• WAP (retrograde conduction through bypass tract)
Wide complex
• SVT with aberrancy or BBB
• Ventricular
Irregular
• Atrial Fibrillation
• Atrial Flutter (variable block)
• Frequent APBs,VPBS


Arrhythmia
Abnormal impulse formation
Abnormal impulse conduction
Automaticity
Triggered activity
Anatomic reentry
Functional reentry (spiral/scroll wave or rotor)
Overdrive suppressed
Not overdrive suppressed (parasystole)
Early after depolarizations
Delayed after depolarizations
Leading circle
Anisotropic reentry
Spatially discordant APD alternans-induced reentry
Phase 2 reentry
Reflection

Step 3: Axis

Look at I and AVF:

- Up Up = Normal
- Up Down = Left
  - LVH, LBBB, inferior MI, WPW, LAF
- Down Up = Right
  - RVH, PE, COPD, septal defects, lateral MI, WPW, LPFB
- Down Down = Extreme




Step 4: Intervals: PR

PR interval:

From onset of atrial depolarization (beginning of P wave) to onset of vent. depolarization


1. Normal = 0.12 -0.2 (~1 big box)
2. Short in WPW, where accessory pathway bypasses AV node
3. If >0.2 then its 1“st-degree AV block”



Step 4: Intervals: QRS

Time it takes for ventricular depolarization to occur.

- Normal < 0.12 (3 small boxes or “1/2” a large box ”
- If QRS complex is wide ensure 1st the patient is stable!
- If unstable (and rhythm is VT need to immediately shock)
- If QRS is wide & rhythm is supraventricular — then determine WHY QRS is wide.




QRS Widening


There are only 3 reasons for QRS widening with sinus rhythm:

1. Typical RBBB
2. Typical LBBB
3. If QRS is wide but neither typical RBBB nor LBBB is present. In this case, the reason the QRS is wide will be IVCD (IntraVentricular Conduction Delay).

The **only 3 key leads** needed to determine type of conduction defect (RBBB, LBBB, or IVCD) — are **leads I, V1 & V6**



"The reason you want to recognize conduction defects before you get too far in the interpretation process — is that criteria for axis, LVH, RVH, ischemia and infarction are all different when there is BBB or IVCD!"




Has there been anterior infarction (slanted arrows)? Is there lateral ischemia (red circle in lead V6)?

The patient is hemodynamically stable. The rhythm is sinus (vertical arrow in the lead II insert) — and the QRS complex is wide. Thus, although one might be tempted to call anterior MI (slanted arrows in V1,V2,V3) and ongoing ischemia (red circle in lead V6) — the Algorithm in prior slide advises us to first STOP and diagnose WHY the QRS is wide (?) before proceeding further.

The reason for QRS widening is complete LBBB (As a result —deep anterior QS complexes + lateral ST-T wave changes are expected consequences of LBBB and not indicative of prior infarction or ischemia.

Left Bundle Branch Block

- In normal conduction, impulses travel equally down left & right bundles, with septum activated from left to right hence the formation of small Q waves in lateral leads
- In LBBB, conduction delay means impulses travel first via right bundle branch to RV, and then to the LV via the septum
- Septal activation is thus reversed eliminating lateral Q waves
    - *The overall depolarisation vector from the right to left ventricle produces tall R waves in lateral leads (I, V5-6) and deep S waves in the right precordial leads (V1-3). The delay between activation of the RV and LV produces the characteristic "M-shaped" R wave seen in lateral leads*
- Delayed conduction time to LV extends QRS to > 120ms.



Sequence of conduction in LBBB:

1) Conduction delay means impulses travel first via the right bundle branch (black arrow)
2) Septum is activated from right-to-left (yellow arrows)
3) Overall depolarisation vector is directed towards lateral leads (red arrow)


…contd.

1. QRS duration > 120ms
2. Dominant S wave in V1, rS complex (small R, deep S)
3. Broad monophasic R wave in lateral leads (I, aVL, V5-6)
4. Absence of Q in lateral leads
5. Prolonged R wave peak time > 60ms in leads V5-6

Appropriate discordance: Lateral leads with tall, broad R waves will often have associated ST-segment depression and T-wave inversion, and those with deep S waves can have an allowable amount of ST elevation that does not indicate ischaemia




LBBB: Left Bundle Branch Block
V1: Dominant S wave
V6: broad, notched ('M'-shaped) R wave



Right Bundle Branch Block

- In RBBB, the left ventricle is activated normally, thus the early part of the QRS complex correlating to septal depolarisation is unchanged
- There is delayed activation of the right ventricle as depolarisation originates from the left ventricle across the septum.
- This produces a secondary R wave (R') in the precordial leads, and a wide, slurred S wave in lateral leads
- Normal activation of left ventricle means that cardiac axis remains normal in isolated RBBB



Sequence of conduction in RBBB:
1) Left ventricular activation via the left bundle (black arrow) occurs normally
2) Septal depolarisation (yellow arrows) is thus unaffected, producing a normal early QRS complex
3) Activation of the RV originates across the septum. The resultant depolarisation vector (red arrow) produces delayed R waves in leads V1-3, and S waves in lateral leads

…contd.

1. QRS duration > 120ms
2. RSR’ pattern in V1-3 (“M-shaped” QRS complex)
3. Wide, slurred S wave in lateral leads (I, aVL, V5-6)


Appropriate discordance with ST depression and/or T-wave inversion in right precordial leads (V1-3).



RBBB: Right Bundle Branch Block


V1: RSR' pattern in V1, with (appropriate) discordant T wave changes

V6: Widened, slurred S wave in V6




Step 4: Intervals: Prolonged QT?

- QT/QTc interval
  - Screen with T < 50% between Q-Q
  - QT varies w/ HR → corrected w/ Bazett formula: QTc = QT/(RR in sec), overcorrects at high HR,
- drugs: amiodarone, genetic: cardiac ion channel mutation, myocardial disease: MI, RF, 3rd degree, low Ca2+, low K+, low Mg2+





(NEJM 2008;358:169; www.torsades.org)


HEY GIRL, ARE YOU AN EKG SEGMENT?
BECAUSE YOU ARE A QT

Step 5: The P's (Atriums)

- ECG is neither sensitive nor specific for atrial enlargement
- Prefered terms, "LAA" + "RAA" (where "A" stands for Abnormality)
- P wave in lead V1 may be positive, negative or biphasic (Panel B)
- Where in lead II, P is upright.






Step 5: The P's


P Pulmonale

- RAA (Right Atrial Abnormality) — is diagnosed by the finding of tall Peaked and Pointed P waves in the Pulmonary leads. Because patients with COPD often have low diaphragms (therefore an inferior axis) — we think of inferior leads (II,III,aVF) as "pulmonary" leads.

P Mitrale

- LAA (Left Atrial Abnormality) — is diagnosed by finding an m-shaped (notched) and widened P wave (≥0.12 second) in a "mitral" lead (I,II,aVL) — and/or a deep negative component to P wave in lead V1 = P Mitrale"





Step 5: The Qs (RVH)

- Criteria [all insensitive, but specific (except in COPD); all w/ poor PPV in general population]




- **Diagnostic criteria**
  1. Right axis deviation of +110° or more.
  2. Dominant R wave in V1 (> 7mm tall or R/S ratio > 1).
  3. Dominant S wave in V5 or V6 (> 7mm deep or R/S ratio < 1).
  4. *QRS duration < 120ms (i.e. changes not due to RBBB).*

Circ 2009;119:e251; JACC 2014;63:672

Step 5: The Qs (LVH)


- Sensitivity of ECG for detecting LVH does not exceed 60%



1. There are numerous **voltage criteria** for diagnosing LVH, summarised below
2. Most common used is ***Sokolov-Lyon criteria***: S wave depth in V1 + tallest R wave height in V5-V6 > 35 mm
3. Voltage criteria must be accompanied by **non-voltage** criteria to be considered diagnostic of LVH


Step 6: ST/Ts

- Use of PR segment baseline (Panel A) — to determine if there is ST elevation (B) or ST depression (C)
- Panel D — shows J-point notching that is characteristic of early repolarization




…contd.

- Ischemia: lack of oxygenation
  - ST segment depression or T wave inversion in consecutive leads
- Injury: prolonged ischemia
  - ST segment elevation peaked T waves or T wave inversions in consecutive leads
- Infarct: death of tissue
  - may or may not show a Q wave
  - Q waves > 1/3 hight of QRS or > 1 small box width in consecutive leads (not in III and aVR)



NEJM 2003;349:2128; Circ 2009;119:e241 & e262



Coronary Anatomy & ECG Leads

Lateral Leads	I, aVL, V5 – V6	LCx or Diagonal of LAD
Inferior Leads	II, III, aVF	RCA and/or LCx
Anterior/Septal Leads	V1 – V4	LAD

Step 7: Special Cases

- Targeted to WHY you ordered an ECG in the first place, i.e
  - Right and left bundle branches- “William Marrow”, bunny ears or RsR’
  - AV Blocks- 1st-3rd degree blocks and treatment
  - Delta Waves- Shark fins sign for Wolf Parkinson White
  - Peaked Ts or U waves- Hyper and Hypokalemia
  - S1Q3T3- Not sensitive but most specific ECG sign of PE
  - Brugada Syndrome- AD cause of long QT and sudden cardiac death
  - Epsilon Waves- Specific finding for ARVC

Step 8: Overall Review


- Review findings and write the final summery statement of the EKG:
  - Ex: Abnormal EKG with sinus tachycardia, LAD, prolonged QRS and LVH.



Practice Case 1

59 year old smoker with a history of DM2, HLD and HTN Presents for acute onset Crushing chest pain.

What is your diagnosis? vessel supplies this area? best treatments acutely and long term? acutely and long term? his risk factors? short and long term complications?




Ans.

1. The ECG shows classic findings of acute/hyperacute anterior wall Q wave myocardialinfarction (MI), with reciprocal inferior ST depressions.
2. The distribution of changes is consistent with a proximal left anterior descending (LAD) occlusion
3. MONA BASSH- Morphine, Oxygen, Nitroglycerine, Aspirin/Clopidogrel and Beta Blocker, ACE Inhibitor/ARB, Statin, Short acting Nitrates, Heparin/LMWH (ExTRACT TIMI 25 and ESSENSE trials)
4. CABG: L main, 3 vessels, 2 vessels + DM (>70% occlusion), pain despite maximum medical tx, or post-infarction angina
5. Always quit smoking, manage comorbidities aggressively (HTN, DM2 etc.). Premature atherosclerosis here was associated with multiple risk factors for coronary artery disease: hypertension, hyperlipidemia, Diabetes, and tobacco.

Practice Case 2

A 39-year-old patient with ESRD due to SLE presents for N/V after missing his last dialysis appointment. He was trying a new juicing cleanse with lots of fruits. EKG was obtained.

What electrolyte disturbance is likely? treatment options are available and in what order? can you tell what level his K is based on EKG changes? Why or why not? What fruit has the highest level of this electrolyte?




Ans.

1. Hyperkalemia (K+ = 8.7 mEq/L) secondary to acute on chronic renal failure. The ECG shows symmetrically peaked ("tented") T waves associated with potassium levels in excess of 6 mEq/L.
2. The tracing also shows broad and flattened sinus P waves that may precede frank sinoventricular conduction seen with severe hyperkalemia
3. CaGluconate/CaCl, Insulin D5, Albuterol, Lactulose, Lasix, SPS
4. Note that T wave peaking with hyperkalemia is a relative finding: the absolute magnitude of the T waves cannot be used to rule in or rule out hyperkalemia.
5. It’s all about the rate that the K+ goes up. Thus specific levels of serum potassium cannot be linked to unique ECG patterns!
6. Avocados, Guavas and Kiwis all have more K per 100g than a banana

Practice Case 3

A 35 year old with a history of lupus and recent URI presents for chest pain. They are sitting up in bed and wince with every breath. EKG is shown.

What is your diagnosis? lab workup and imaging after this EKG? common causes of this? What are your treatment options for this patient?




Ans.

1. The ECG shows classic findings for acute pericarditis.
2. These include diffuse ST segment elevations (I, II aVF, V2-V6). In addition, there are subtle PR segment deviations (elevated in aVR and depressed in the inferolateral leads). There are also non-diagnostic inferior Q waves.
3. The S1Q3 pattern here is a non-specific finding.
4. CBC, CMP, ESR, CRP, Trop, CXR, TTE, COVID, +/- serologies
5. Post-infectious (viral, protozoa, mycobacteria), autoimmune (SLE, RA, sarcoid, Amyloidosis), post-infarct, hemopericardium, neoplastic, uremia, drug induced, radiation
6. NSAIDS, Colchicine, prednisone, pericardiocentesis, pericardial window.

Practice Case 4

A 61 year old female with a history of rheumatic heart disease as a child presents to her PCP with increasing dyspnea on exertion. Her exam is notable for 4/6 systolic murmur.

What is your diagnosis? what is your lab workup and imaging after this EKG? common causes of this? treatment options for this rhythm?




Ans.

1. The ECG shows very coarse atrial fibrillation with RAD along with T wave inversions in V1-V4 (strain pattern), all consistent with right ventricular overload.
2. Atrial Fibrillation and possible right heart strain/pulmonary hypertension.
3. CBC, CMP, TSH, Troponin, BNP. Echocardiogram and possible RHC.
4. AV nodal blockers such as BB, CCB, Digoxin or conversion with Amiodarone or cardioversion.

EKG Patterns and Buzzwords

- Pattern recognition and illness scripts
  - Wide = Ventricular
  - Narrow = Atria
  - Regular =Sinus Tach, AVNRT,
  - WPW, Aflutter or Vtach
  - Irregular = MAT, Afib or Vfib




Sinus Tachycardia

A 49 year old man presents to the ED because of fever and a change in mental status. He works as a construction worker and had cut himself incidentally earlier in the week but did not clean the wound. Vital signs are BP 105/63, pulse 123 and regular, Temperature 38.8 and reparations 22. His skin is flushed and diaphoretic. Initial labs show a leukocytosis with left shift.


Look for: Narrow regular pattern with discernible P waves (lookout for camel hump)

- Causes- Think: Pain/Anxiety, Hypoxia/Hypovolemia, PE – most common EKG finding for PE, Sepsis +Acidemia
- x: Fix the underlying cause/Ivabradine



SVT or AVNRT

A 38 year old women suddenly felt dizzy and calls 911 because “her heart is going to beat out of her chest”. She has had 2 episodes of this before that resolved spontaneously. When she presents to the hospital, vital signs are BP 128/72, pulse 158 and regular, respirations 22. EKG is shown below.



Look for: Narrow + regular pattern without discernible P waves


- Most common cause of SVT is AVRT where 2 pathways (one fast and one slow) within AV node forming reentrant loop usually Initiated or terminated by PACs
- Tx: Adenosine, verapamil, vagal stimulation (carotid massage etc.)

Wolf-Parkinson-White

23 year old medical student experiences sudden onset palpitations and shortness of breath while studying for final exams. He had stayed up all night and consumed multiple cups of coffee and energy drinks. On presentation to the ED, his EKG is shown below.

Look for: Swooping R waves, shortened PR interval and pseudo- widened QRS

- Pathophysiology: An accessory pathway from atria to ventricles through the bundle of Kent early ventricular activation because the AV node does not slow conduction.
- Tx: unstable - synchronized cardioversion, stable- procainamide. Long term/definitive Tx is radiofrequency ablation




Atrial Flutter

A 67 year old gentleman with episodic dizziness and shortness of breath is placed on a Holter monitor. On analysis of his read out you see the rhythm shown below.

Look for: Sawtooth Pattern with regular rate 150-200 bpm


- Causes: Heart disease: Heart failure (most common association), rheumatic heart disease, CAD, COPD, Atrial septal defect (ASD)
- Treat: AV nodal blocking agents or ablation (IVC/tricuspid isthmus)
- https://litfl.com/atrial-flutter-ecg-library/



Atrial Fibrillation

She lives alone and is unsure how long she was unconscious but denies tongue biting, incontinence or confusion upon awakening. She also notices her left leg weak. She feel her heart beat very fast at times but did not seek treatment for this. A code stroke is called and EKG shows:

Look for: “Undulating baseline”, no discernible p-waves, irregular R-R interval




- Causes- Heart disease: CAD, MI, HTN, mitral regurge, Pericarditis, rheumatic disease, Pulmonary disease, including PE, Hyperthyroidism, hypothyroidism or Pheo, Excessive alcohol intake “holiday heart syndrome”
- Treatment: AV nodal blockers or synchronized cardioversion if unstable. CHA2DS2-VASc score to determine anticoagulation.

Ventricular Tachycardia

A page brings you to the room of a 67 year old who is post op day 1 from an emergent CABG status post STEMI. He is unresponsive with this EKG rhythm:

Look for: > 30 seconds of wide QRS, with a regular rhythm

- Causes: CAD with prior MI- most common cause, Active ischemia, hypotension, Cardiomyopathies, Congenital defects, Prolonged QT syndrome
- Tx: Amiodarone or Sotalol if hemodynamically stable. Cardioversion if unstable. ACLS if pulseless.
- All those with impaired EF and sustained Vtach need an ICD!



Ventricular Fibrillation

A 57 year old patient on the general medicine floor is suddenly found to be unconscious. She is being treated for pneumonia with azithromycin. Her history includes amitriptyline for depression and fibromyalgia, quetiapine for bipolar depression, ondansetron for nausea and methadone for pain. Upon further review of the telemetry reading you notice this rhythm.




Look for: Wide irregular rhythm. Torsades is polymorphic

- Causes-Ischemic heart disease- most common cause + Antiarrhythmic drugs, especially those that cause prolonged QT intervals

Multifocal Atrial Tachycardia


A 78 year old veteran presents to the hospital for shortness of breath. He has a 60 pack year smoking history and has a history of severe COPD requiring 5L continuous oxygen support. His medication reconciliation shows multiple beta agonist inhalers and theophylline. Vital sign show BP 112/63, pulse of 129 and irregular, temperature of 37.2, and respirations are 18 per minute and prolonged expirations. On chest exam, diffuse wheezes are heart in all lung fields with prolonged respirations and an increased AP diameter. There is a loud P2 and a right heart heave. EKG shows the rhythm below:



Look for: Three + morphologically distinct P waves in the same lead Varying PR intervals.

Ectopic Beats

A 23 year old medical student presents for palpitations and tachycardia. He Just finished an overnight call and has an exam upcoming. He consumed countless cups of coffee in an effort to study. An EKG performed shows random segments of this rhythm.




What to look for and causes of ectopic foci of depolarization

1. PAC: A premature QRS complex with normal to abnormal P wave due to sympathomimetics, anxiety, caffeine, hypokalemia, or hypomagnesemia
2. PJC: QRS complex without a preceding P wave
3. PVC: A premature, broad QRS complex due t sympathomimetics, anxiety, caffeine, hypokalemia, hypomagnesemia, dioxin toxicity, or ischemia. 3 more more sustained PVCs are defined as NSVT

Cardiac Tamponade

45 year old with a history of Marfan's Disease presents with acute chest pain that feels "like a knife" radiating to his back. On exam, he has unequal BP of 154/92 and 132/87 in his arms and pulse of 110. Suddenly, he develops JVD, hypotension and pulsus paradoxus. Heart sounds are distant. The EKG taken at that time shows:



Look for: Alternating QRS complex amplitude "electrical alternans"


- Causes: Aortic dissection (hypertension, smoking, collagen vascular, and syphilis), TB, metastatic tumors
- Tx: Pericardiocentesis (US guided in subxiphoid or 5th intercostal space)

Sinus Bradycardia

A 22 year old marathon runner presents for NCAA pre-participation physical. A screening EKG is performed showing this rhythm.

Look for: regular P waves rate < 60 bpm


- Causes: Think normal finding in athletes, or ischemia, increased vagal tone, antiarrhythmic drugs (AV blockers), brainstem herniation
- Tx: Atropine or a cardiac pacemaker



1st° AV Block + Mobitz type I (Wenckebach)

A 23 year old medical student from New Jersey has complaints of palpitations and joint pain following a hiking trip. He is unsure if there was a rash or tick bite history.

- **1st Degree AV Block**
  - Look for: prolonged PR interval with no dropped beats
- **2nd Degree Heart Block Mobitz Type 1**
  - Look for: progressive, prolongation of the PR interval followed by a dropped beat
- Tx: Doesn't require treatment unless unexplained symptoms https://litfl.com/first-degree-heart-block-ecg-library






Mobitz type II + 3rd° AV Block

A 78 year old women presents for syncope and a feeling of “my heart skipping a beat” for 1 week. She has a history of MI and CHF. Vital signs are BP 133/79, pulse 61 bpm with a regular rate. On exam, irregular jugular venous wave forms “cannon-a waves” are seen.


- **2nd Degree Mobitz Type**
  - Look for: prolonged PR interval with dropped beats
- **3rd Degree or Complete Heart Block**
  - Look for: regular P-Pinterval and regularR-R interval
- Tx: Pacemaker implantation



Electrolytes and Intoxications


A 73 year old with renal failure presents for chest pain and palpitations after missing their last 3 dialysis appointments.

- **Hyperkalemia - “peaked” T-waves, widened QRS (sine wave), shortQTcand prolonged PR.**
  - Causes: Think crush injury, burn injury, renal failure
  - Tx: IV CaGluconate, D5 and insulin, beta agonists, kayexalate, furosemide or dialysis



A 17 year old with a history of anorexia and diuretic use presents for inpatient rehabilitation program. She is severely malnourished and labs/ecg are performed.

- **Hypokalemia -“U wave” and long QTc**
  - Tx: replete K (oral>IV) and Mg




Prolonged QT

A 76 year old with a history of depression and Afib is admitted to an inpatient service for pneumonia. She is treated with azithromycin and levofloxacin and she develops nausea treated with ondansetron. Overnight she develops chest discomfort and an EKG is below.

- Causes: Hypokalemia, hypomagnesemia, hypocalcemia, hypothermia, DRUGS


Look for: QT is more than half the preceding RR interval

- Different Measures: Bazett: Overcorrects above 100 bpm. Fridericia /Framingham are more accurate



HOCM

A 17 year old basketball player suddenly collapses at a local AAU tournament. EMS arrives and the patient regains consciousness. In ED the patient is well appearing. A 3/6 systolic murmur is auscultated in RUSB that decreases with hand grip. EKG shown.




Caused by AD mutation in sarcomere proteins (beta-myosin heavy chain, troponin T) that causes an enlarged, obstructing septum. Tx with ICD or BB

Look for: LVH, or deep, narrow “daggerlike”. Q waves in V5-6, I, aVL and II, III or giant T-wave inversion

Syncope: PE

A 24 year old with a history of SLE on OCPs presents with shortness of breath, pleuritic chest pain, calf swelling and dizziness after a long distance flight.

Look for an S1Q3T3 pattern a prominent S wave in lead I a Q wave and inverted T wave in lead III, Sinus tachycardia, T wave inversion in leads V1 - V3 and Right Bundle Branch Block




Syncope: Brugada Syndrome

A 33 year old with sleep apnea comes into the office for a life insurance screening. His father and uncle died suddenly in their 40s. He has pasted out while running but thought nothing of it. EKG shown:

- Causes: Genetic defects in cardiac sodium transporters
- Treatment: ICD

Look for: Coved ST segment elevation >2mm in >1 of V1-V3 followed by a negative T wave or >2mm of saddleback shaped ST elevation



Sick Sinus Syndrome

A 91 year old women presents with a 2 month history of periotic pre-syncope, dizziness, confusion, fatigue and palpitations. She has had no previous cardiac history and only takes a multivitamin. During your interview her pulse ranges from 40-120 bpm.

Look for: Alternating bradycardia, tachycardia and sinus arrest

- Causes: Often idiopathic and occurs in elderly, sarcoidosis, ischemia. Tx: Pacemaker

References

- Roden, Dan M. “Long-QT Syndrome.” *New England Journal of Medicine*, vol. 358, no. 2, Jan. 2008, pp. 169–76. *Taylor and Francis+NEJM*, https://doi.org/10.1056/NEJMcp0706513.
- *Home :: Crediblemeds*. https://crediblemeds.org/. Accessed 11 Sept. 2022.
- Hancock, E. William, et al. “AHA/ACCF/HRS Recommendations for the Standardization and Interpretation of the Electrocardiogram.” *Circulation*, vol. 119, no. 10, Mar. 2009, pp. e251–61. *ahajournals.org (Atypon)*, https://doi.org/10.1161/CIRCULATIONAHA.108.191097.
- Whitman, Isaac R., et al. “Validity of the Surface Electrocardiogram Criteria for Right Ventricular Hypertrophy: The MESA-RV Study (Multi-Ethnic Study of Atherosclerosis-Right Ventricle).” *Journal of the American College of Cardiology*, vol. 63, no. 7, Feb. 2014, pp. 672–81. *PubMed*, https://doi.org/10.1016/j.jacc.2013.08.1633.
- Gami, Apoor S., et al. “Electrocardiographic Poor R-Wave Progression: Analysis of Multiple Criteria Reveals Little Usefulness.” *American Heart Journal*, vol. 148, no. 1, July 2004, pp. 80–85. *DOI.org (Crossref)*, https://doi.org/10.1016/j.ahj.2004.02.005.
- Wang, Kyuhyun, et al. “ST-Segment Elevation in Conditions Other Than Acute Myocardial Infarction.” *New England Journal of Medicine*, vol. 349, no. 22, Nov. 2003, pp. 2128–35. *Taylor and Francis+NEJM*, https://doi.org/10.1056/NEJMra022580.