

# Outline

- ▶ Introduction
- ▶ Methods for controlling error rate(s)
- ▶ Conclusions

# Introduction

- ▶ You have previously considered the significance of a single gene

# Introduction

- ▶ You have previously considered the significance of a single gene
- ▶ The analysis of high-dimensional data is concerned with assessing the significance of multiple loci/genes
  - ▶ Microarray : 20,000-50,000 probe sets
  - ▶ GWAS: 500,000-5,000,000 typed SNPs
  - ▶ RNA-Seq: 22,000 genes (humans)

# Introduction

- ▶ You have previously considered the significance of a single gene
- ▶ The analysis of high-dimensional data is concerned with assessing the significance of multiple loci/genes
  - ▶ Microarray : 20,000-50,000 probe sets
  - ▶ GWAS: 500,000-5,000,000 typed SNPs
  - ▶ RNA-Seq: 22,000 genes (humans)
- ▶ This leads to the *Multiple Testing* problem

# Introduction

- ▶ You have previously considered the significance of a single gene
- ▶ The analysis of high-dimensional data is concerned with assessing the significance of multiple loci/genes
  - ▶ Microarray : 20,000-50,000 probe sets
  - ▶ GWAS: 500,000-5,000,000 typed SNPs
  - ▶ RNA-Seq: 22,000 genes (humans)
- ▶ This leads to the *Multiple Testing* problem
- ▶ Before we address this problem, let's quickly review the single hypothesis case

# Introduction: Type I and II Errors

Recall that in hypothesis testing with a single hypothesis (gene), errors can be classified as:

- ▶ Type I error - rejecting  $H_0$  when it is true
- ▶ Type II error - not rejecting  $H_0$  when it is false

# Introduction: Type I and II Errors

Recall that in hypothesis testing with a single hypothesis (gene), errors can be classified as:

- ▶ Type I error - rejecting  $H_0$  when it is true
- ▶ Type II error - not rejecting  $H_0$  when it is false

We can define probabilities associated with each of these errors:

$$\alpha = Pr(\text{reject } H_0 | H_0 \text{ true})$$

$$\beta = Pr(\text{do not reject } H_0 | H_0 \text{ false})$$

# Introduction: Type I and II Errors

Decision	$H_0$ True	$H_0$ False
Do not reject $H_0$	$1 - \alpha$	$\beta$
Reject $H_0$	$\alpha$	$1 - \beta$



# Introduction: Type I and II Errors

- ▶ Always a compromise between type I and type II error

# Introduction: Type I and II Errors

- ▶ Always a compromise between type I and type II error
- ▶ An  $\alpha$ -level test has probability  $\alpha$  of rejecting  $H_0$  when  $H_0$  is true

# Introduction: Type I and II Errors

- ▶ Always a compromise between type I and type II error
- ▶ An  $\alpha$ -level test has probability  $\alpha$  of rejecting  $H_0$  when  $H_0$  is true
- ▶ Usually try to use the most powerful (smallest  $\beta$ ) test for a given  $\alpha$ -level

# Introduction: Type I and II Errors

- ▶ Always a compromise between type I and type II error
- ▶ An  $\alpha$ -level test has probability  $\alpha$  of rejecting  $H_0$  when  $H_0$  is true
- ▶ Usually try to use the most powerful (smallest  $\beta$ ) test for a given  $\alpha$ -level  $\rightarrow$  *control* type I error

# Introduction: Multiple Tests

- ▶ Single-hypothesis type-I-error-control proves inadequate when experiment is characterized by a collection of hypotheses

# Introduction: Multiple Tests

- ▶ Single-hypothesis type-I-error-control proves inadequate when experiment is characterized by a collection of hypotheses  
← Genomics

# Introduction: Multiple Tests

- ▶ Single-hypothesis type-I-error-control proves inadequate when experiment is characterized by a collection of hypotheses  
← Genomics
- ▶ Why?

# Introduction: Multiple Tests

- ▶ Assume we have  $m$  hypotheses we wish to test as part of a given experiment



# Introduction: Multiple Tests

- ▶ Assume we have  $m$  hypotheses we wish to test as part of a given experiment
  - ▶ These hypotheses could correspond to  $m$  genes that we are investigating for differential expression between two groups

# Introduction: Multiple Tests

- ▶ Assume we have  $m$  hypotheses we wish to test as part of a given experiment
  - ▶ These hypotheses could correspond to  $m$  genes that we are investigating for differential expression between two groups
- ▶ Assume that each hypothesis is tested using a  $\alpha$ -level test

# Introduction: Multiple Tests

- ▶ Assume we have  $m$  hypotheses we wish to test as part of a given experiment
  - ▶ These hypotheses could correspond to  $m$  genes that we are investigating for differential expression between two groups
- ▶ Assume that each hypothesis is tested using a  $\alpha$ -level test
- ▶ Assume that the tests are *INDEPENDENT* and that the null is true for each of the hypothesis

# Introduction: Multiple Tests

What is the probability of rejecting a null hypothesis even though all hypotheses are actually null?

# Introduction: Multiple Tests

What is the probability of rejecting a null hypothesis even though all hypotheses are actually null?

- ▶ For a given test, we have an  $\alpha$  chance of rejecting its null

# Introduction: Multiple Tests

What is the probability of rejecting a null hypothesis even though all hypotheses are actually null?

- ▶ For a given test, we have an  $\alpha$  chance of rejecting its null
- ▶ Therefore we have a  $1 - \alpha$  chance of not rejecting

# Introduction: Multiple Tests

What is the probability of rejecting a null hypothesis even though all hypotheses are actually null?

- ▶ For a given test, we have an  $\alpha$  chance of rejecting its null
- ▶ Therefore we have a  $1 - \alpha$  chance of not rejecting
- ▶ We have  $m$  independent  $\alpha$ -level tests, so the chance of not rejecting any of them is:

$$(1 - \alpha)^m$$

# Introduction: Multiple Tests

What is the probability of rejecting a null hypothesis even though all hypotheses are actually null?

- ▶ For a given test, we have an  $\alpha$  chance of rejecting its null
- ▶ Therefore we have a  $1 - \alpha$  chance of not rejecting
- ▶ We have  $m$  independent  $\alpha$ -level tests, so the chance of not rejecting any of them is:

$$(1 - \alpha)^m$$

- ▶ Therefore, the chance of rejecting at least one hypothesis is:

$$1 - (1 - \alpha)^m$$



# Introduction: Multiple Tests

# Introduction: Notation

- ▶ Gene  $j$  (among the  $m$  genes) is either associated with the outcome or not
- ▶ The truth is unknown to us
- ▶ The null hypothesis for gene  $j$  is denoted by  $H_j$
- ▶  $H_j$ : gene  $j$  is not associated with the outcome of interest
- ▶ The alternative hypothesis is denoted by  $\bar{H}_j$
- ▶  $\bar{H}_j$ : gene  $j$  is associated with the outcome of interest
- ▶  $H_j$  and  $\bar{H}_j$  are called *marginal* or *local* hypotheses

# Introduction: Marginal and Global Hypotheses

- ▶  $H_j$  and  $\bar{H}_j$  are called *marginal* or *local* hypotheses
- ▶ A global null hypothesis: None of the  $m$  genes is associated with the outcome
- ▶ A global alternative: At least one of the  $m$  genes is associated with the outcome
- ▶ Using notation
  - ▶ Global Null:  $\mathbb{H}_0 : H_1 \text{ and } H_2 \text{ and } \dots H_m$
  - ▶ Global Alternative:  $\mathbb{H}_1 : \bar{H}_1 \text{ or } \bar{H}_2 \text{ or } \dots \bar{H}_m$

# Introduction: Unadjusted vs Adjusted $P$ -values

- ▶ Suppose that we only test a single gene, say gene  $j$ , among the  $m$  genes
- ▶ Let  $p_j$  (lower case  $p$ ) denote  $P$ -value corresponding to  $H_j$
- ▶  $p_j$  is called the *marginal* or *unadjusted*  $P$ -value
- ▶ If  $m$  hypotheses are tested, inference on  $H_j$  on the basis of  $p_j$  is inappropriate
- ▶ The  $P$ -value for  $H_j$  has to account for testing the other  $m - 1$  hypotheses
- ▶ We will denote the *adjusted*  $P$ -value by  $P_j$  (upper case  $P$ )
- ▶ When testing multiple genes, using the marginal  $P$ -value is inappropriate
- ▶ Why?

# Introduction: More Notation

- ▶ Suppose that gene  $j$  is not associated with the outcome of interest ( $H_j$  is true)
  - ▶ Then
    - ▶ Decision rule rejects  $\rightarrow$  False-Positive (FP)
    - ▶ Decision rule fails to reject  $\rightarrow$  True-Negative (TN)
- ▶ Suppose that gene  $j$  is associated with the outcome of interest ( $H_j$  is false)
  - ▶ Decision rule rejects  $\rightarrow$  True-Positive (TP)
  - ▶ Decision rule fails to reject  $\rightarrow$  False-Negative (FN)

# Introduction: Summarizing a Multiple Testing Procedure

- ▶ The results from any multiple testing procedure can be summarized as the following table

	Accept	Reject	Total
Truth Null	$A_0$	$R_0$	$m_0$
Alt.	$A_1$	$R_1$	$m_1$
	$A$	$R$	$m$

- ▶ Notation:
  - ▶  $m$ : Number of tests,  $m_0, m_1$  number of null/true genes
  - ▶  $R$ : Number of genes rejected according to the decision rule
  - ▶  $A$ : Number of genes accepted according to the decision rule
  - ▶  $R_0/R_1$  number of TN/FP
  - ▶  $A_0/A_1$  number of FN/TP

# Introduction: Example

- Results from an analysis based on  $m = 10$  genes:

##	gene	truth	pvalue
## 1	gene1	0	0.29070
## 2	gene2	1	0.61630
## 3	gene3	1	0.00320
## 4	gene4	0	0.01641
## 5	gene5	0	0.25150
## 6	gene6	0	0.58450
## 7	gene7	0	0.22890
## 8	gene8	1	0.12630
## 9	gene9	0	0.26080
## 10	gene10	0	0.04980

- Investigator decides to use following decision rule: Any gene with a corresponding unadjusted  $P$ -value of less than 0.05 will be rejected.
- Reject  $H_j$  if  $p_j < 0.05$  or accept  $H_j$  otherwise

## Exercise: Fill in the 2x2 table

	Accept	Reject	Total
Truth Null	$A_0 = ?$	$R_0 = ?$	$m_0 = ?$
Alt.	$A_1 = ?$	$R_1 = ?$	$m_1 = ?$
	$A = ?$	$R = ?$	$m = ?$



## Example: Fill in the 2x2 table

	Accept	Reject	Total
Truth Null	$A_0 = 5$	$R_0 = 2$	$m_0 = 7$
Alt.	$A_1 = 2$	$R_1 = 1$	$m_1 = 3$
	$A = 7$	$R = 3$	$m = 10$

- ▶  $m_0 = 7$  and  $m_1 = 3$
- ▶  $R = 3$  will be rejected based on the decision rule
- ▶ Consequently  $A = m - R = 7$  will be accepted
- ▶  $R_0 = 2, R_1 = 1, A_0 = 5$  and  $A_1 = 2$

# The Truth

- ▶ What know or observe is this

```
##      gene  pvalue
## 1  gene1 0.29070
## 2  gene2 0.61630
## 3  gene3 0.00320
## 4  gene4 0.01641
## 5  gene5 0.25150
## 6  gene6 0.58450
## 7  gene7 0.22890
## 8  gene8 0.12630
## 9  gene9 0.26080
## 10 gene10 0.04980
```

- ▶ and not (truth column is not known to us):

```
dat

##      gene truth  pvalue
## 1  gene1     0 0.29070
## 2  gene2     1 0.61630
## 3  gene3     1 0.00320
## 4  gene4     0 0.01641
## 5  gene5     0 0.25150
## 6  gene6     0 0.58450
## 7  gene7     0 0.22890
## 8  gene8     1 0.12630
## 9  gene9     0 0.26080
## 10 gene10    0 0.04980
```

## Example: Fill in the 2x2 table (based on what we observe)

- ▶ We can only fill in the bottom row of the table

	Accept	Reject	Total
Truth Null	$A_0$	$R_0$	$m_0$
Alt.	$A_1$	$R_1$	$m_1$
	$A = 7$	$R = 3$	$m = 10$

- ▶ The remaining quantities are fixed unknown quantities or unobservable random variables.

# Comments

	Accept	Reject	Total
Truth Null	$A_0$	$R_0$	$m_0$
Alt.	$A_1$	$R_1$	$m_1$
	$A$	$R$	$m$

# Comments

	Accept	Reject	Total
Truth Null	$A_0$	$R_0$	$m_0$
Alt.	$A_1$	$R_1$	$m_1$
	$A$	$R$	$m$

- $m$  is a known constant

# Comments

	Accept	Reject	Total
Truth Null	$A_0$	$R_0$	$m_0$
Alt.	$A_1$	$R_1$	$m_1$
	$A$	$R$	$m$

- ▶  $m$  is a known constant
- ▶  $m_0$  and  $m_1$  are unknown constants

# Comments

	Accept	Reject	Total
Truth Null	$A_0$	$R_0$	$m_0$
Alt.	$A_1$	$R_1$	$m_1$
	$A$	$R$	$m$

- ▶  $m$  is a known constant
- ▶  $m_0$  and  $m_1$  are unknown constants
- ▶  $R$  and  $A$  are determined on the basis of applying the decision rule to the data

# Comments

	Accept	Reject	Total
Truth Null	$A_0$	$R_0$	$m_0$
Alt.	$A_1$	$R_1$	$m_1$
	$A$	$R$	$m$

- ▶  $m$  is a known constant
- ▶  $m_0$  and  $m_1$  are unknown constants
- ▶  $R$  and  $A$  are determined on the basis of applying the decision rule to the data
- ▶ They are *observable* random quantities



# Comments

	Accept	Reject	Total
Truth Null	$A_0$	$R_0$	$m_0$
Alt.	$A_1$	$R_1$	$m_1$
	$A$	$R$	$m$

- ▶  $m$  is a known constant
- ▶  $m_0$  and  $m_1$  are unknown constants
- ▶  $R$  and  $A$  are determined on the basis of applying the decision rule to the data
- ▶ They are *observable* random quantities
- ▶ The true states of the genes of the genes are unknown

# Comments

	Accept	Reject	Total
Truth Null	$A_0$	$R_0$	$m_0$
Alt.	$A_1$	$R_1$	$m_1$
	$A$	$R$	$m$

- ▶  $m$  is a known constant
- ▶  $m_0$  and  $m_1$  are unknown constants
- ▶  $R$  and  $A$  are determined on the basis of applying the decision rule to the data
- ▶ They are *observable* random quantities
- ▶ The true states of the genes of the genes are unknown
- ▶  $A_0, A_1, R_0$  and  $R_1$  are *unobservable* random quantities

# Introduction: Multiple Testing Problem

- ▶ Control error rate(s) in multiple testing context

# Introduction: Multiple Testing Problem

- ▶ Control error rate(s) in multiple testing context
- ▶ Multiple testing methods are designed to control a particular error rate

# Introduction: Multiple Testing Problem

- ▶ Control error rate(s) in multiple testing context
- ▶ Multiple testing methods are designed to control a particular error rate
- ▶ Multiple error rates exist

# Introduction: Multiple Testing Problem

- ▶ Control error rate(s) in multiple testing context
- ▶ Multiple testing methods are designed to control a particular error rate
- ▶ Multiple error rates exist → need to choose error rate to control and then method to control it

# Introduction: Error Rates

- ▶ **Family-wise error rate (FWER)**: the probability of at least one type I error

# Introduction: Error Rates

- ▶ **Family-wise error rate (FWER)**: the probability of at least one type I error
- ▶ **False discovery rate (FDR)**: the expected proportion of type I errors among the rejected hypotheses.



# Family-wise Error Rate (FWER)

- ▶ Probability of committing at least one false-rejection (among  $m$ ) given that *all* genes are null

# Family-wise Error Rate (FWER)

- ▶ Probability of committing at least one false-rejection (among  $m$ ) given that *all* genes are null
- ▶  $\text{FWER} = P(R \geq 1 | m = m_0)$

# Family-wise Error Rate (FWER)

- ▶ Probability of committing at least one false-rejection (among  $m$ ) given that *all* genes are null
- ▶  $\text{FWER} = P(R \geq 1 | m = m_0)$
- ▶ Note that when  $m = 1$  (single gene), this definition is identical to the type I error we have previously considered

## Controlling FWER: Sidak's method

Recall that we showed that with  $m$  independent  $\alpha$ -level tests:

$$\text{FWER} = 1 - (1 - \alpha)^m$$

# Controlling FWER: Sidak's method

Recall that we showed that with  $m$  independent  $\alpha$ -level tests:

$$\text{FWER} = 1 - (1 - \alpha)^m$$

Therefore,

$$\begin{aligned} 1 - \text{FWER} &= (1 - \alpha)^m \\ (1 - \text{FWER})^{1/m} &= 1 - \alpha \\ 1 - (1 - \text{FWER})^{1/m} &= \alpha \end{aligned}$$

# Controlling FWER: Sidak's method

Recall that we showed that with  $m$  independent  $\alpha$ -level tests:

$$\text{FWER} = 1 - (1 - \alpha)^m$$

Therefore,

$$\begin{aligned} 1 - \text{FWER} &= (1 - \alpha)^m \\ (1 - \text{FWER})^{1/m} &= 1 - \alpha \\ 1 - (1 - \text{FWER})^{1/m} &= \alpha \end{aligned}$$

This suggests that we can control FWER by choosing  $\alpha$  for each individual test to be  $1 - (1 - \text{FWER})^{1/m}$

# Controlling FWER: Bonferroni

Note that  $(1 - x)^m \approx 1 - mx$  (for  $x$  close to zero)

# Controlling FWER: Bonferroni

Note that  $(1 - x)^m \approx 1 - mx$  (for  $x$  close to zero)



# Controlling FWER: Bonferroni

Note that  $(1 - x)^m \approx 1 - mx$  (for  $x$  close to zero)

Therefore

$$1 - \text{FWER} = (1 - \alpha)^m$$

$$1 - \text{FWER} \approx 1 - m\alpha$$

$$\text{FWER} \approx m\alpha$$

# Controlling FWER: Bonferroni

Note that  $(1 - x)^m \approx 1 - mx$  (for  $x$  close to zero)

Therefore

$$1 - \text{FWER} = (1 - \alpha)^m$$

$$1 - \text{FWER} \approx 1 - m\alpha$$

$$\text{FWER} \approx m\alpha$$

This suggests that we can control FWER by choosing  $\alpha$  for each individual test to be  $\frac{\text{FWER}}{m}$

# Controlling FWER: Bonferroni

- ▶ The Bonferroni adjusted *P-value* is defined as

$$P_j = m \times p_j$$

- ▶ Technical note:  $P_j$  as defined above could be larger than 1 so a more technically rigorous definition is

$$P_j = \min\{m \times p_j, 1\}$$

- ▶ In other words, if  $m \times p_j$  is larger than 1, then truncate  $P_j$  at 1.

# Controlling FWER: Holm's

- ▶ Sidak and Bonferroni are one-step approaches

# Controlling FWER: Holm's

- ▶ Sidak and Bonferroni are one-step approaches
- ▶ Some power can be gained by sequential methods

# Controlling FWER: Holm's

- ▶ Sidak and Bonferroni are one-step approaches
- ▶ Some power can be gained by sequential methods
- ▶ The simplest sequential method is Holm's method:

# Controlling FWER: Holm's

- ▶ Sidak and Bonferroni are one-step approaches
- ▶ Some power can be gained by sequential methods
- ▶ The simplest sequential method is Holm's method:
  - ▶ Order the unadjusted  $P$ -values such that  $p_1 \leq p_2 \leq \dots \leq p_m$

# Controlling FWER: Holm's

- ▶ Sidak and Bonferroni are one-step approaches
- ▶ Some power can be gained by sequential methods
- ▶ The simplest sequential method is Holm's method:
  - ▶ Order the unadjusted  $P$ -values such that  $p_1 \leq p_2 \leq \dots \leq p_m$
  - ▶ To control FWER, the step-down Holm adjusted  $P$ -values are

$$P_j = \min\{(m - j + 1) \times p_j, 1\}$$



# Controlling FWER: Holm's

- ▶ Sidak and Bonferroni are one-step approaches
- ▶ Some power can be gained by sequential methods
- ▶ The simplest sequential method is Holm's method:
  - ▶ Order the unadjusted  $P$ -values such that  $p_1 \leq p_2 \leq \dots \leq p_m$
  - ▶ To control FWER, the step-down Holm adjusted  $P$ -values are

$$P_j = \min\{(m - j + 1) \times p_j, 1\}$$

- ▶ Note that every unadjusted  $P$ -value is not multiplied by same factor

# Controlling FWER: Permutation

- ▶ These methods work well when tests are independent

# Controlling FWER: Permutation

- ▶ These methods work well when tests are independent (not generally the case in genomics experiments)

# Controlling FWER: Permutation

- ▶ These methods work well when tests are independent (not generally the case in genomics experiments)
- ▶ When tests are correlated, these methods are conservative

# Controlling FWER: Permutation

- ▶ These methods work well when tests are independent (not generally the case in genomics experiments)
- ▶ When tests are correlated, these methods are conservative
- ▶ Permutation approaches are useful in this context

# Controlling FWER: Permutation

Assume we are interested in assessing differential expression between 2 groups :

1. Compute minimum unadjusted  $P$ -value for all genes from the observed data (call it  $p_1$ )
2. Randomly permute the group labels
  - ▶ *Breaks* relationship between group and expression
  - ▶ Reflects sample from global null hypothesis
3. Compute minimum  $P$ -value from data set generated in 2 (call it  $p_1^1$ )
4. Repeat 2 and 3  $B$  times to get  $p_1^1, p_1^2, \dots, p_1^B$
5. Compute the proportion of  $p_1^1, p_1^2, \dots, p_1^B$  that are  $\leq p_1$
6. This proportion is the permutation adjusted  $P_1$

# Controlling FWER: Permutation

- ▶ Can get  $P_2, \dots, P_m$  in a similar way (a Holm's-like permutation procedure)

# Controlling FWER: Permutation

- ▶ Can get  $P_2, \dots, P_m$  in a similar way (a Holm's-like permutation procedure)
- ▶ Correlation among the genes is accounted for



# False Discovery Rate (FDR)

- ▶ Consider the quantity  $\frac{R_0}{R}$
- ▶ This is the proportion of false discoveries among the genes rejected
- ▶ This is an *unobservable* random quantity ( $R_0$  is not observable)
- ▶ In the FDR framework is based on controlling the *expected* value of this ratio
- ▶  $\text{FDR} \equiv E[\frac{R_0}{R}]$ 
  - ▶ Expectation is set to zero if  $R = 0$ ,  
therefore  $\text{FDR} = E[\frac{R_0}{R} | R > 0]Pr(R > 0)$
- ▶ Note that when  $m_0 = m$  (none of the genes are true),  
 $\text{FWER} = \text{FDR}$

# Controlling FDR: Benjamini and Hochberg

1. Order the unadjusted  $P$ -values  $p_1 \leq p_2 \leq \dots \leq p_m$

# Controlling FDR: Benjamini and Hochberg

1. Order the unadjusted  $P$ -values  $p_1 \leq p_2 \leq \dots \leq p_m$
2. Find the largest  $j$  such that  $p_j \leq j\alpha/m$

# Controlling FDR: Benjamini and Hochberg

1. Order the unadjusted  $P$ -values  $p_1 \leq p_2 \leq \dots \leq p_m$
2. Find the largest  $j$  such that  $p_j \leq j\alpha/m$
3. Reject null hypotheses affiliated with  $p_1, p_2, \dots, p_j$

# Controlling FDR: Benjamini and Hochberg

1. Order the unadjusted  $P$ -values  $p_1 \leq p_2 \leq \dots \leq p_m$
2. Find the largest  $j$  such that  $p_j \leq j\alpha/m$
3. Reject null hypotheses affiliated with  $p_1, p_2, \dots, p_j$

This procedure will control FDR at  $\alpha m_0/m$  when the tests are independent and continuous

# Controlling FDR: Benjamini and Hochberg

1. Order the unadjusted  $P$ -values  $p_1 \leq p_2 \leq \dots \leq p_m$
2. Find the largest  $j$  such that  $p_j \leq j\alpha/m$
3. Reject null hypotheses affiliated with  $p_1, p_2, \dots, p_j$

This procedure will control FDR at  $\alpha m_0/m$  when the tests are independent and continuous

Benjamini and Yekutieli (2001) proposed a modification of the BH procedure that always controls FDR (no larger than  $\alpha m_0/m$ )

# Controlling FDR: Benjamini and Hochberg

1. Order the unadjusted  $P$ -values  $p_1 \leq p_2 \leq \dots \leq p_m$
2. Find the largest  $j$  such that  $p_j \leq j\alpha/m$
3. Reject null hypotheses affiliated with  $p_1, p_2, \dots, p_j$

This procedure will control FDR at  $\alpha m_0/m$  when the tests are independent and continuous

Benjamini and Yekutieli (2001) proposed a modification of the BH procedure that always controls FDR (no larger than  $\alpha m_0/m$ )

← can be quite conservative

# Controlling FDR: Benjamini and Hochberg

1. Order the unadjusted  $P$ -values  $p_1 \leq p_2 \leq \dots \leq p_m$
2. Find the largest  $j$  such that  $p_j \leq j\alpha/m$
3. Reject null hypotheses affiliated with  $p_1, p_2, \dots, p_j$

This procedure will control FDR at  $\alpha m_0/m$  when the tests are independent and continuous

Benjamini and Yekutieli (2001) proposed a modification of the BH procedure that always controls FDR (no larger than  $\alpha m_0/m$ )

← can be quite conservative

Note that when  $m_0 = m$  (i.e., all hypotheses are null), these procedures maintain FWER at  $\alpha$



## Controlling pFDR: q-values

$$\text{FDR} = E\left[\frac{R_0}{R} \mid R > 0\right] Pr(R > 0)$$

## Controlling pFDR: q-values

$$\text{FDR} = E\left[\frac{R_0}{R} \mid R > 0\right] \Pr(R > 0)$$

$$\text{pFDR} = E\left[\frac{R_0}{R} \mid R > 0\right] \leftarrow \text{positive FDR}$$

# Controlling pFDR: q-values

$$\text{FDR} = E\left[\frac{R_0}{R} \mid R > 0\right] \Pr(R > 0)$$

$$\text{pFDR} = E\left[\frac{R_0}{R} \mid R > 0\right] \leftarrow \text{positive FDR}$$

- ▶ Since  $\Pr(R > 0)$  is often  $\sim 1$  in most genomics experiments, FDR and pFDR are ver similar

# Controlling pFDR: q-values

- ▶ q-value is the minimum pFDR for which the affiliated hypothesis is rejected

# Controlling pFDR: q-values

- ▶ q-value is the minimum pFDR for which the affiliated hypothesis is rejected
- ▶ q-value can be interpreted as the expected proportion of false positives incurred when calling that test significant

# Genome-wide Significance

- ▶ In GWAS papers,  $\alpha = 5 \times 10^{-8}$  is typically considered the threshold for genome-wide significance
- ▶ It is based on a Bonferroni correction: If you consider testing  $m = 1,000,000$  SNPs at the FWER level of 0.05, then each SNP should be tested at the

$$\alpha = \frac{0.05}{1,000,000} = 5 \times 10^{-8},$$

level

- ▶ Suppose that the unadjusted  $P$ -value for a SNP is  $5 \times 10^{-7}$
- ▶ Is this "reaching" genome-wide significance?
- ▶ The term "suggestive" is also used

## "Reaching" Genome-wide Significance

- ▶ Suppose that your  $m = 1,000,000$  SNPs are independent
- ▶ The adjusted  $P$ -value is

$$P = 5 \times 10^{-7} \times m = 5 \times 10^{-7} \times 10^6 = 0.5,$$

- ▶ This is off by an order of magnitude ( $0.5 = 0.05 \times 10$ )
- ▶ It is not "reaching"
- ▶ Note: Due to linkage disequilibrium among SNPs the adjusted  $P$ -value is likely to be smaller than 0.5
- ▶ The point is that while  $5 \times 10^{-7}$  is small number, it may not be small enough when testing a large number of hypotheses

# Conclusions

- ▶ Multiple testing *must* be accounted for when testing for associations in the context of high-dimensional data
- ▶ FWER and FDR are the two common frameworks for quantifying error
- ▶ Error rate estimates can be used to compute 'adjusted' p-values
- ▶ Resampling-based methods can increase power in controlling error when sample sizes are sufficient for their use.
- ▶ When large-scale patterns of differential expression are observed, it is important to consider if such effects are biologically reasonable, and if technical factors can be attributed to the variation.