CALIFORNIA STATE UNIVERSITY, NORTHRIDGE

PROTEIN FOLDING: PLANAR CONFIGURATION SPACES OF DISC ARRANGEMENTS AND HINGED POLYGONS: PROTEIN FOLDING IN FLATLAND

A thesis submitted in partial fulfillment of the requirements For the degree of Master of Science in Mathematics

By

Clinton Bowen

Spring 2014

The thesis of Clinton Bowen is approve	d:
Dr. John Dye	Date
Dr. Silvia Fernandez	 Date
Dr. Bernardo Abrego	Date
Dr. Csaba Toth, Chair	 Date

California State University, Northridge

DEDICATIONS

ACKNOWLEGDEMENTS

Table of Contents

Signa	ture pag	ge		•	•		•		•	•	•	•	•		•	•	•	•	•	•	•	•		•	•	•	•	•	ii
Dedic	cation.						•									•			•					•					iii
Ackn	owledge	ement .																											iv
																													vi
1		ction .																											1
2	_	ound .																											2
	2.1	Linkag																											2
	2.2	Circle	Packi	ng	•	٠	•	•	•	٠	•	•	٠	•	٠	٠	٠	•	٠	•	٠	٠	•	٠	٠	٠	٠	•	3
2.2.1	Circle	Packing	gs and	l Po	lyg	ona	l L	ink	ag	es																			
2.2.2	Hinge	d Polygo	ons																										
2.2.3	Hinge	d Hexag	gons c	f Fi	xed	l Si	ze																						
3	Configuration Spaces of Polygonal Chains															5													
	3.1	Polygo																											5
3.1.1																													
	3.2	Dissect	tions.																										6
	3.3	SAT P	robler	ns.															•						•	•			6
3.3.1	3-SAT Problems Problem																												
4	Probler	n																											6
	4.1	Proble	m Sta	tem	ent																								6
	4.2	Decida																											6
	4.3	Hexago																											6
5	Conclu																												7

ABSTRACT

PROTEIN FOLDING: PLANAR CONFIGURATION SPACES

OF DISC ARRANGEMENTS AND HINGED POLYGONS:

PROTEIN FOLDING IN FLATLAND

Ву

Clinton Bowen

Master of Science in Mathematics

Insert Abstract here

Abstract

We look into the decidability of whether a hinged configuration locks.

1 Introduction

We look into the decidability of continuity on planar configuration space using regular, unitary hexagonal polygons. These polygons can also represent unit disk configurations [4]




Figure 1: A locked 7 ball configuration

Motivation Protein folding, graphite, crystalline structures in metallurgy; disc packing; hexagonal configurations; Determine whether chemical structures are realizable.

Outline Section 2 covers the necessary mathematical concepts to understanding the problem. Section 3 explains the problem, Section 4 covers the results and findings about the problem. Section 5, the conclusion, offers final remarks on the problem.

2 Background

Here we review some of the necessary mathematics behind the problem. The definitions found in this chapter are those found in [9, 11, 8].

2.1 Linkages

Definition 2.1 (Graph). An ordered pair G = (V, E) comprising a set V of vertices or nodes together with a set E of edges or lines

Definition 2.2 (Linkage). A collection of fixed-length 1D segments joined at their endpoints to form a graph.

A linkage can be thought of as a type of path-connected graph, i.e. the segments of a linkage are the edges of a graph, and the endpoints of the segments are the vertices. For this paper, we restrict our self to linkages that are simple planar graphs, i.e. a linkage that:

- (i) does not have multiple edges between any pair of vertices,
- (ii) does not have edges that cross, or

(iii) have loops (i.e. $(v, v) \in E$).

Definition 2.3 (Cycle). A closed walk with no repetitions of vertices or edges allowed, other than the repetition of the starting and ending vertex

Definition 2.4 (Configuration). A specification of the location of all the link endpoints, link orientations and joint angles.[6]

Definition 2.5 (Configuration Space). The space of all configurations of a linkage.

A configurations space is said to be continuous if for any two configurations, \mathscr{A} and \mathscr{B} of a linkage L, \mathscr{A} can be continuously reconfigured to \mathscr{B} such that, the reconfigurations reside in the configuration domain, L remains rigid throughout reconfiguration (i.e. all links' lengths are preserved), and no violations of linkage intersection conditions.

Definition 2.6 (Pinned Joint). A vertex of a graph (or linkage) that is fixed to a position in a plane.

Definition 2.7 (Free Joint). A vertex of a graph (or linkage) that is not fixed to a position in a plane.




Figure 2: A linkage with joints.




Figure 3: The cross represents a free joint; the pinned joints are denoted as disks. The range of motion shown by the arc describes the continous configuration space of the linkage.

For illustrations in the remainder of this paper, free joints will be represented as crosses and pinned joints will be represented as disks.

2.2 Circle Packing

Definition 2.8 (Circle Packing). P of a planar graph G is a set of of circles with disjoint interiors $\{C_v\}_{v \in G}$ such that two circles are tangent if and only if the corresponding vertices form an edge. [2]

Theorem 2.1 (Circle Packing Theorem). For every connected simple planar graph G there is a circle packing in the plane whose intersection graph is (isomorphic to) G.

A proof of Theorem 2.1 is found in [11].




Figure 4: This figure is an example of a circle packing for the given simple planar graph.




Figure 5: A circumbscribed hexagon

2.2.1 Circle Packings and Polygonal Linkages

Given a circle of radius r and its center point, (x,y), we establish the isomorphism to a hexagon by circumscribing the vertices of the regular hexagon.

2.2.2 Hinged Polygons

Definition 2.9 (Polygonal Chain). A polygonal chain $P = (v_0, v_1, \dots, v_{n-1})$ is a sequence of consecutively joined segments (or edges) $e_i = v_i v_{i+1}$ of fixed lengths $l_i = |e_i|$, in a plane. [3]

A chain is said to be closed if $v_{n-1} = v_1$, otherwise it is said to be open. Hinged polygons have been researched for decades and related to linkage problems [3, 5].

Consider the locked configuration of figure 6. We can configure the hexagons to be locked by placing hinged points as follows:




Figure 6: A locked 7 hexagonal configuration. (needs to modify picture by placing red points for hing points.)

2.2.3 Hinged Hexagons of Fixed Size

The Shapes Figure 7 is a locking shape: Figure 7 shall reside in the boundary of a lattice and have a hinge




Figure 7: A locking shape in the lattice boundary's channel.

point at one vertex where the locking shape and boundary meet.

Junctions We define junctions to be the point three hexagons meet in a hexagonal lattice, e.g. Figure 8.




Figure 8: A portion of a hexagonal lattice.

Central Scaling

Junctions in Conjunctive Normal Form Explain the configurations we're interested in.

3 Configuration Spaces of Polygonal Chains

3.1 Polygonal Linkages




Figure 9: Not sure how to describe this graph with free joints, pinned joints. Do I need to define another joint(i.e. axis of rotation joint)?

3.1.1 Configurations and Locked Configurations

3.2 Dissections

Problem 3.1 (Polygonal Dissection). Given two polygons of equal area, P_1 and P_2 , partition P_1 into smaller pieces, $\{P_{1,i}\}_{i=1}^n$, rearrange the pieces to form P_2 . [8]

Theorem 3.1. Any finite collection of polygons of equal area has a common hinged dissection. [1]




Figure 10: An axample of two polygons of equal area that can be rearranged into the other by the given partition.[7]

3.3 SAT Problems

Problem 3.2 (Satisfiability Problem). Let $\{x_i\}_{i=1}^n$ be boolean variables, and $t_i \in \{x_i\}_{i=1}^n \cup \{\bar{x}_i\}_{i=1}^n$. A *clause* is is said to be a disjuction of distinct terms:

$$t_1 \vee \cdots \vee t_{j_k} = C_k$$

Then the *satisfiability problem* is the decidability of a conjuction of a set of clauses, i.e.:

$$\wedge_{i=1}^m C_i$$

[10]

3.3.1 3-SAT Problems

A 3-SAT problem is a SAT problem with all clauses having only three boolean variables.

4 Problem

4.1 Problem Statement

text

4.2 Decidability of Problem

test

4.3 Hexagonal Locked Configuration




Figure 11: 7 hexagonal configuration

5 Conclusion

We conclude...

References

- [1] Timothy G Abbott, Zachary Abel, David Charlton, Erik D Demaine, Martin L Demaine, and Scott Duke Kominers. Hinged dissections exist. *Discrete & Computational Geometry*, 47(1):150–186, 2012.
- [2] Omer Angel, Martin T. Barlow, Ori Gurel-Gurevich, and Asaf Nachmias. Boundaries of planar graphs, via circle packings. November 2013.
- [3] T. Biedl, E. Demaine, M. Demaine, S. Lazard, A. Lubiw, J. O'Rourke, M. Overmars, S. Robbins, I. Streinu, G. Toussaint, and S. Whitesides. Locked and Unlocked Polygonal Chains in 3D, 1999.
- [4] Heinz Breu and David G. Kirkpatrick. Unit disk graph recognition is NP-hard. *Computational Geometry*, 9(1–2):3–24, 1998. Special Issue on Geometric Representations of Graphs.
- [5] J. Canny. *The Complexity of Robot Motion Planning*. ACM doctoral dissertation award. MIT Press, 1988.
- [6] E.D. Demaine and J. O'Rourke. *Geometric Folding Algorithms: Linkages, Origami, Polyhedra*. Cambridge University Press, 2008.
- [7] David Eppstien. The Geometry Junkyard, November 2013.
- [8] G.N. Frederickson. Dissections: Plane and Fancy. Cambridge University Press, 1997.
- [9] J. Kleinberg and E. Tardos. Algorithm Design. Pearson Education, 2006.
- [10] S.S. Skiena. The Algorithm Design Manual. Springer, 2009.
- [11] K. Stephenson. *Introduction to Circle Packing: The Theory of Discrete Analytic Functions*. Cambridge University Press, 2005.