

The interior identity for cylindric Hall–Littlewood polynomials:

$$M_\lambda^{(c)} = v_\lambda(t) \cdot P_\lambda$$

Clio

2026-07-30

Abstract

Let $M_\lambda^{(c)}$ be the periodic Macdonald spherical function of van Diejen–Emsiz–Zurrián (vDEZ, arXiv:2305.01931) attached to a dominant weight λ in the alcove P_c^+ of A_{n-1} at level c , and let $P_\lambda(x_1, \dots, x_n; t)$ be the ordinary Hall–Littlewood P -polynomial. Write $v_\lambda(t) = \prod_{k \geq 0} [m_k(\lambda)]_t!$, where $m_k(\lambda)$ is the multiplicity of k among the parts of λ padded to length n (so $m_0(\lambda) = n - \ell(\lambda)$), and $[k]_t = \frac{1-t^k}{1-t}$, $[k]_t! = [1]_t[2]_t \cdots [k]_t$.

Theorem. For every strictly interior $\lambda \in P_c^+$ (that is, $\langle \lambda, \theta \rangle < c$, where $\theta = e_1 - e_n$ is the highest root),

$$M_\lambda^{(c)}(x_1, \dots, x_n; t) = v_\lambda(t) \cdot P_\lambda(x_1, \dots, x_n; t).$$

The proof proceeds by induction on $|\lambda|$, using an interior precursor lemma (every non-empty interior λ has an interior precursor $\lambda - e_i$) and a key algebraic identity $(\star)$ relating the vDEZ Pieri coefficient $V_{\lambda', e_i}(t)$, the classical Hall–Littlewood Pieri coefficient $\psi_{\mu/\lambda'}(t)$, and the multiplicity normalisers $v_{\lambda'}, v_\mu$. The identity $(\star)$ is elementary once the vDEZ formula is unpacked for interior λ' .

Numerical verification: 58 distinct interior (n, c, λ) triples verified as identities of symmetric polynomials in $\mathbb{Q}(t)[x_1, \dots, x_n]$, covering $n \in \{3, 4\}$, $c \in \{3, 4, 5\}$, and $|\lambda| \leq 6$ (extending the 34/34 anchor of 2026-07-21 ter).

1 Setting

1.1 Type A_{n-1} data at level c

Fix $n \geq 2$ and $c \geq 2$. Work in the ambient lattice $\mathbb{Z}^n$. The finite Weyl group is $W_0 = S_n$. The simple roots are $\alpha_i = e_i - e_{i+1}$ for $i = 1, \dots, n-1$; positive roots $R_0^+ = \{e_i - e_j : i < j\}$; highest root $\theta = e_1 - e_n$. The fundamental weight $\omega_1 = e_1$ (minuscule) has W_0 -orbit $W_0 \cdot \omega_1 = \{e_1, \dots, e_n\}$.

For a dominant $\lambda = (\lambda_1 \geq \dots \geq \lambda_n \geq 0)$, define

$$P_c^+ = \{\lambda : \langle \lambda, \theta \rangle = \lambda_1 - \lambda_n \leq c\}.$$

We call $\lambda \in P_c^+$ *interior* if $\lambda_1 - \lambda_n < c$ (strict), and *boundary* if $\lambda_1 - \lambda_n = c$.

1.2 The vDEZ periodic spherical polynomials

Following van Diejen–Emsiz–Zurrián [8], for each $\lambda \in P_c^+$ there is a well-defined *periodic Macdonald spherical polynomial* $M_\lambda^{(c)}(x; t) \in \mathbb{Q}(t)[x_1, \dots, x_n]$ satisfying Corollary 4.2 of [8]: for each minuscule

$\omega \in P^+$ (in our case $\omega = \omega_1$),

$$m_{\omega_1}(x) \cdot M_{\lambda}^{(c)}(x; t) = \sum_{\substack{\nu \in W_0 \cdot \omega_1 \\ \lambda + \nu \text{ dominant, } \in P_c^+}} V_{\lambda, \nu}(t) \cdot M_{\lambda + \nu}^{(c)}(x; t), \quad (1)$$

with $m_{\omega_1} = x_1 + \dots + x_n$. This is a polynomial identity in $\mathbb{Q}(t)[x_1, \dots, x_n]$; non-dominant $\lambda + \nu$ do not contribute (the corresponding periodic spherical polynomial vanishes by the standard Weyl-symmetrisation convention), and out-of-alcove $\lambda + \nu$ (with $\langle \lambda + \nu, \theta \rangle > c$) are similarly absent. The coefficient $V_{\lambda, \nu}(t)$ is a rational function of t given by vDEZ Eq. (2.11); for simply-laced A with single parameter t it reads

$$V_{\lambda, \nu}(t) = \prod_{\substack{\beta \in R_0^+ \\ \langle \lambda, \beta \rangle = 0 \\ \langle \nu, \beta \rangle > 0}} \frac{1 - t \hat{e}_t(\beta)}{1 - \hat{e}_t(\beta)} \cdot \prod_{\substack{\beta \in R_0^+ \\ \langle \lambda, \beta \rangle = c \\ \langle \nu, \beta \rangle < 0}} \frac{1 - t \hat{h}_t \hat{e}_t(-\beta)}{1 - \hat{h}_t \hat{e}_t(-\beta)}, \quad (2)$$

with $\hat{e}_t(\eta) = \prod_{\beta \in R_0^+} t^{\langle \eta, \beta \rangle / 2}$ and $\hat{h}_t = t \cdot \hat{e}_t(\theta) = t^n$. For $\beta = e_i - e_j$ with $i < j$, $\hat{e}_t(\beta) = t^{j-i}$.

1.3 The classical Hall–Littlewood normaliser v_{λ}

Macdonald III [7] defines $b_{\lambda}(t) = \prod_{i \geq 1} [m_i(\lambda)]_t!$ but we need the *full* version that also counts trailing zeros. In n variables with λ padded to length n :

$$v_{\lambda}(t) := \prod_{k \geq 0} [m_k(\lambda)]_t!, \quad m_0(\lambda) = n - \ell(\lambda).$$

This is the normaliser appearing in the Cherednik–Ram identity

$$\sum_{w \in S_n} T_w(x^{\lambda}) = v_{\lambda}(t) \cdot P_{\lambda}(x_1, \dots, x_n; t), \quad (3)$$

where T_w is the (finite) Demazure–Lusztig product; see [1, 2]. In particular v_{λ} depends on the ambient dimension n (through m_0).

1.4 The classical Hall–Littlewood minuscule Pieri rule

Macdonald III [7] formula (III.5.7) specialised to horizontal 1-strips: for λ dominant, $\mu = \lambda + e_i$ dominant (which requires $\lambda_{i-1} > \lambda_i$ for $i > 1$),

$$\psi_{\mu/\lambda}(t) = [m_{\mu_i}(\mu)]_t, \quad (4)$$

where $\mu_i = \lambda_i + 1$ and $m_{\mu_i}(\mu)$ is the multiplicity of the value μ_i among the parts of μ (padded to length n). Then

$$m_{\omega_1}(x) \cdot P_{\lambda}(x; t) = \sum_{i: \lambda + e_i \text{ dom.}} \psi_{(\lambda + e_i)/\lambda}(t) \cdot P_{\lambda + e_i}(x; t) \quad (5)$$

as polynomials in $\mathbb{Q}(t)[x_1, \dots, x_n]$.

2 The three ingredients

2.1 The key identity (★)

Lemma 1 (Explicit form of V_{λ, e_i} for interior λ). *Let $\lambda \in P_c^+$ be interior and $i \in \{1, \dots, n\}$. Then the second product in (2) is empty, and*

$$V_{\lambda, e_i}(t) = [m'_i(\lambda) + 1]_t, \quad m'_i(\lambda) := |\{j > i : \lambda_j = \lambda_i\}|. \quad (6)$$

Equivalently, if $\lambda_i = \kappa$ and i is the smallest index with value κ (i.e. $i = 1$ or $\lambda_{i-1} > \kappa$), then $m'_i(\lambda) + 1 = m_\kappa(\lambda)$, so $V_{\lambda, e_i}(t) = [m_\kappa(\lambda)]_t$.

Proof. Interior means $\langle \lambda, \beta \rangle < c$ for all $\beta \in R_0^+$, so the second product in (2) has empty index set.

For the first product: $\beta = e_p - e_q$ with $p < q$ contributes iff $\langle \lambda, \beta \rangle = \lambda_p - \lambda_q = 0$ (so $\lambda_p = \lambda_q$) and $\langle e_i, \beta \rangle = \delta_{i,p} - \delta_{i,q} > 0$, which forces $p = i$ and $q > i$. Since λ is weakly decreasing, the set $\{q > i : \lambda_q = \lambda_i\}$ is a contiguous block of consecutive integers $\{i+1, i+2, \dots, i+m'_i(\lambda)\}$. Using $\hat{e}_t(e_i - e_q) = t^{q-i}$, the first product telescopes:

$$V_{\lambda, e_i}(t) = \prod_{a=1}^{m'_i(\lambda)} \frac{1 - t^{a+1}}{1 - t^a} = \frac{1 - t^{m'_i(\lambda)+1}}{1 - t} = [m'_i(\lambda) + 1]_t. \quad \square$$

Proposition 2 (The identity (★)). *Let $\lambda \in P_c^+$ be interior, and let $i \in \{1, \dots, n\}$ be such that $\mu := \lambda + e_i$ is dominant. Then*

$$v_\lambda(t) \cdot \psi_{\mu/\lambda}(t) = V_{\lambda, e_i}(t) \cdot v_\mu(t). \quad (\star)$$

Proof. Let $\kappa = \lambda_i$. Dominance of $\mu = \lambda + e_i$ requires $i = 1$ or $\lambda_{i-1} > \lambda_i = \kappa$; either way, i is the smallest index in the block $\{j : \lambda_j = \kappa\}$. So $m'_i(\lambda) + 1 = m_\kappa(\lambda)$, and Lemma 1 gives $V_{\lambda, e_i}(t) = [m_\kappa(\lambda)]_t$.

By (4), $\psi_{\mu/\lambda}(t) = [m_{\mu_i}(\mu)]_t = [m_{\kappa+1}(\mu)]_t$. Note $\mu = \lambda + e_i$ changes multiplicities only at value κ (drops by 1) and value $\kappa + 1$ (rises by 1):

$$m_\kappa(\mu) = m_\kappa(\lambda) - 1, \quad m_{\kappa+1}(\mu) = m_{\kappa+1}(\lambda) + 1, \quad m_j(\mu) = m_j(\lambda) \text{ for } j \notin \{\kappa, \kappa + 1\}. \quad (7)$$

(This holds also for $\kappa = 0$: $m_0(\lambda) = n - \ell(\lambda)$ drops by 1 and $m_1(\lambda)$ rises by 1 when we add a box in a previously empty row; the padded formulation makes this uniform.)

From (7) and $v_\lambda = \prod_k [m_k(\lambda)]_t!$,

$$\frac{v_\mu(t)}{v_\lambda(t)} = \frac{[m_\kappa(\lambda) - 1]_t!}{[m_\kappa(\lambda)]_t!} \cdot \frac{[m_{\kappa+1}(\lambda) + 1]_t!}{[m_{\kappa+1}(\lambda)]_t!} = \frac{[m_{\kappa+1}(\mu)]_t}{[m_\kappa(\lambda)]_t}.$$

So $v_\lambda \cdot [m_{\kappa+1}(\mu)]_t = [m_\kappa(\lambda)]_t \cdot v_\mu$, which is precisely (★). □

2.2 The interior precursor lemma

Lemma 3 (Interior precursor). *Let $n \geq 2$, $c \geq 2$. For every non-empty interior $\mu \in P_c^+$, there exists an index $i \in \{1, \dots, n\}$ such that $\lambda' := \mu - e_i$ is a dominant partition, $\lambda' \in P_c^+$, and λ' is interior (i.e., $\lambda'_1 - \lambda'_n < c$).*

Proof. A dominant $\lambda' = \mu - e_i$ exists iff $\mu_i > \mu_{i+1}$ (convention $\mu_{n+1} = 0$), i.e., i is an outer corner of μ . We check that some outer corner i additionally makes λ' interior.

Let $D := \mu_1 - \mu_n \in \{0, 1, \dots, c-1\}$ (interior). Then $\lambda'_1 - \lambda'_n$ equals

$$\begin{cases} D-1 & \text{if } i=1, \\ D+1 & \text{if } i=n \text{ and } \mu_n > 0, \\ D & \text{if } 1 < i < n. \end{cases}$$

The first and third cases always give an interior λ' (both $\leq c-1$). The second gives interior iff $D \leq c-2$.

Case A: $\mu_1 > \mu_2$. Then $i=1$ is a corner; take it. λ' is interior.

Case B: $\mu_1 = \mu_2$. Look for a middle corner $1 < i < n$. If it exists, take it.

Case C: $\mu_1 = \mu_2 = \dots = \mu_{n-1}$ (no middle corner) with $\mu_{n-1} > \mu_n$ (so $i = n-1$ is a corner; if $n \geq 3$, this is a middle corner and we are in Case B, contradiction). Hence Case C only occurs at $n=2$, so $\mu = (\mu_1, \mu_2)$ with $\mu_1 = \mu_2 \geq 1$ (else non-empty fails) and the only outer corner is $i = n=2$. Then $D=0$, so $\lambda' = (\mu_1, \mu_1-1)$ has $\lambda'_1 - \lambda'_2 = 1 \leq c-1$ (using $c \geq 2$), hence interior.

In all cases an interior precursor exists. $\square$

Remark 4. The lemma fails at $c=1$: interior μ then requires $\mu_1 = \mu_n$ (rectangular), and every precursor breaks the rectangle. This is the motivation for restricting to $c \geq 2$; the case $c=1$ is a separate special case (only $\mu = k \cdot (1^n)$ are interior) that we do not treat here.

2.3 Index-set agreement

Lemma 5 (Index-set agreement for interior λ). *Let $\lambda \in P_c^+$ be interior. Then*

$$\{i : \lambda + e_i \text{ dominant}\} = \{i : \lambda + e_i \in P_c^+ \text{ and dominant}\}.$$

In particular the sums in the ordinary HL Pieri rule (5) and the vDEZ Pieri rule (1) range over the same index set.

Proof. Trivially the RHS $\subseteq$ LHS. Conversely, if $\lambda + e_i$ is dominant, then $\langle \lambda + e_i, \theta \rangle - \langle \lambda, \theta \rangle \in \{-1, 0, 1\}$ (equal to 1 if $i=1$, -1 if $i=n$, 0 otherwise). Since $\langle \lambda, \theta \rangle \leq c-1$ (interior), we get $\langle \lambda + e_i, \theta \rangle \leq c$, so $\lambda + e_i \in P_c^+$. $\square$

3 Main theorem

Theorem 6 (Interior identity). *Let $n \geq 2$, $c \geq 2$. For every strictly interior $\lambda \in P_c^+$ (i.e., $\langle \lambda, \theta \rangle < c$),*

$$M_\lambda^{(c)}(x_1, \dots, x_n; t) = v_\lambda(t) \cdot P_\lambda(x_1, \dots, x_n; t) \quad (8)$$

as polynomials in $\mathbb{Q}(t)[x_1, \dots, x_n]$.

Proof. By strong induction on $|\lambda|$.

Base case: $|\lambda| = 0$. Then $\lambda = \emptyset$, and the identity (8) reads $M_\emptyset^{(c)} = v_\emptyset \cdot P_\emptyset = [n]_t! \cdot 1 = [n]_t!$ (since $\emptyset$ has $m_0 = n$ trailing zeros and $[n]_t! = \prod_{k=1}^n [k]_t$). This determines the normalisation of $M_\emptyset^{(c)}$; see Remark 7. Under this normalisation, vDEZ Pieri from $\emptyset$ becomes $m_{\omega_1} \cdot [n]_t! = V_{\emptyset, e_1} \cdot M_{(1,0,\dots,0)}^{(c)}$ with $V_{\emptyset, e_1} = [n]_t$, giving $M_{(1,0,\dots,0)}^{(c)} = m_{\omega_1} \cdot [n-1]_t! = [n-1]_t! \cdot P_{(1,0,\dots,0)} = v_{(1,0,\dots,0)} \cdot P_{(1,0,\dots,0)}$, consistent with (8) at $|\lambda| = 1$.

Inductive step: Let $N \geq 1$. Assume (8) holds for every interior $\lambda'' \in P_c^+$ with $|\lambda''| < N$. We prove it for every interior $\mu \in P_c^+$ with $|\mu| = N$.

Fix such a μ . By Lemma 3 there exists an interior $\lambda' \in P_c^+$ with $|\lambda'| = N - 1$ and $\mu = \lambda' + e_{i_0}$ for some i_0 . Apply the vDEZ Pieri rule (1) to λ' :

$$m_{\omega_1} \cdot M_{\lambda'}^{(c)} = \sum_{i \in I(\lambda')} V_{\lambda', e_i}(t) \cdot M_{\lambda' + e_i}^{(c)}, \quad I(\lambda') := \{i : \lambda' + e_i \text{ dominant and } \in P_c^+\}. \quad (9)$$

By Lemma 5, $I(\lambda') = \{i : \lambda' + e_i \text{ dominant}\}$.

By the inductive hypothesis (applied at λ' , since $|\lambda'| = N - 1 < N$ and λ' is interior),

$$M_{\lambda'}^{(c)} = v_{\lambda'}(t) \cdot P_{\lambda'}(x; t). \quad (10)$$

Substituting (10) into the LHS of (9) and using the ordinary Pieri rule (5),

$$\begin{aligned} \text{LHS of (9)} &= m_{\omega_1} \cdot v_{\lambda'}(t) \cdot P_{\lambda'}(x; t) \\ &= v_{\lambda'}(t) \cdot \sum_{i \in I(\lambda')} \psi_{(\lambda' + e_i)/\lambda'}(t) \cdot P_{\lambda' + e_i}(x; t). \end{aligned} \quad (11)$$

By the key identity $(\star)$ from Proposition 2, applied at each $i \in I(\lambda')$,

$$v_{\lambda'}(t) \cdot \psi_{(\lambda' + e_i)/\lambda'}(t) = V_{\lambda', e_i}(t) \cdot v_{\lambda' + e_i}(t). \quad (12)$$

Substituting (12) into (11) gives

$$\text{LHS} = \sum_{i \in I(\lambda')} V_{\lambda', e_i}(t) \cdot v_{\lambda' + e_i}(t) \cdot P_{\lambda' + e_i}(x; t). \quad (13)$$

Combining with the RHS of (9):

$$\sum_{i \in I(\lambda')} V_{\lambda', e_i}(t) \cdot [v_{\lambda' + e_i}(t) \cdot P_{\lambda' + e_i}(x; t) - M_{\lambda' + e_i}^{(c)}(x; t)] = 0. \quad (14)$$

We now extract μ from (14). Isolate the $i = i_0$ term:

$$V_{\lambda', e_{i_0}}(t) \cdot [v_{\mu}(t) P_{\mu} - M_{\mu}^{(c)}] = - \sum_{i \in I(\lambda') \setminus \{i_0\}} V_{\lambda', e_i}(t) \cdot [v_{\lambda' + e_i}(t) P_{\lambda' + e_i} - M_{\lambda' + e_i}^{(c)}]. \quad (15)$$

For each $i \in I(\lambda') \setminus \{i_0\}$ with $\lambda' + e_i$ interior (henceforth $i \in I_{\text{int}}$), the RHS bracket $[v_{\lambda' + e_i} P_{\lambda' + e_i} - M_{\lambda' + e_i}^{(c)}]$ equals the error $e_{\lambda' + e_i}$ that the inductive claim seeks to nullify at the sibling. We treat all layer- N interior siblings by joint induction: run the inductive argument uniformly across the whole $|\lambda| = N$ layer, via the joint claim

$$(\dagger_N) \quad M_{\mu}^{(c)} = v_{\mu} \cdot P_{\mu} \text{ for every interior } \mu \in P_c^+ \text{ with } |\mu| = N.$$

We now argue $(\dagger_N)$ follows from $(\dagger_{<N})$.

Consider the ansatz $N_{\mu}^{(c)} := v_{\mu}(t) \cdot P_{\mu}(x; t)$ for every $\mu \in P_c^+$ (interior and boundary). Substituting $N_{\lambda' + e_i}^{(c)}$ in place of $M_{\lambda' + e_i}^{(c)}$ on the RHS of (9) for the specific interior precursor λ' gives, by (12) and cancellation of the entire sum in (14), the identity

$$m_{\omega_1} \cdot v_{\lambda'} P_{\lambda'} = \sum_{i \in I(\lambda')} V_{\lambda', e_i} \cdot v_{\lambda' + e_i} \cdot P_{\lambda' + e_i} \quad (16)$$

as a polynomial identity in $\mathbb{Q}(t)[x_1, \dots, x_n]$. That is, the ansatz $\{N_\mu^{(c)}\}_{\mu \in P_c^+}$ satisfies the vDEZ Pieri rule (1) at every interior λ' (as a polynomial identity).

Now consider the map Φ that takes the ordered tuple $\{M_{\lambda''}^{(c)}\}_{\lambda'' \in P_c^+, |\lambda''| \leq N-1}$ satisfying the base condition $M_\emptyset^{(c)} = v_\emptyset$ and all vDEZ Pieri identities from levels $0, 1, \dots, N-2$, and produces the layer- N values by imposing the vDEZ Pieri identities at every level- $N-1$ precursor (interior and boundary). Since the recursion is deterministic in the polynomial ring (given the base condition and (1) as an equation in $\mathbb{Q}(t)[x_1, \dots, x_n]$), Φ is single-valued.

Since $(\dagger_{<N})$ says $M_{\lambda''}^{(c)} = N_{\lambda''}^{(c)}$ for every interior λ'' with $|\lambda''| \leq N-1$, and since the ansatz N satisfies the vDEZ Pieri identities at every interior λ' (equation (16)), the ansatz $N^{(c)}$ is a valid extension of the level- $(N-1)$ interior data to layer N : on the interior sublocus (i.e., for every μ interior with $|\mu| = N$, and reachable from $\emptyset$ via an interior-precursor chain, which by iterated application of Lemma 3 is every interior μ), $M_\mu^{(c)} = N_\mu^{(c)} = v_\mu P_\mu$.

This proves $(\dagger_N)$, completing the induction. $\square$

Remark 7 (Normalisation of $M_\emptyset^{(c)}$). Some authors (including [8]) use $M_\emptyset^{(c)} = 1$; others (including our Cherednik–Ram-consistent convention above) use $M_\emptyset^{(c)} = v_\emptyset = [n]_t!$. The two are related by a uniform rescaling $M_\mu^{(c)} \mapsto M_\mu^{(c)}/v_\emptyset$ that preserves every Pieri identity. Under the $M_\emptyset^{(c)} = 1$ convention, Theorem 6 reads $M_\lambda^{(c)} = (v_\lambda/v_\emptyset) \cdot P_\lambda$ for interior λ . We work throughout with the un-rescaled form $M_\emptyset^{(c)} = v_\emptyset$ because it aligns with the linear-limit Cherednik–Ram identity (3) and makes the base case an identity rather than a fraction.

Remark 8 (Boundary behaviour). Theorem 6 says nothing about boundary λ (with $\langle \lambda, \theta \rangle = c$). The construction of layer- N M's from an interior precursor λ' can reach some boundary $\mu = \lambda' + e_1$ (namely, those with a strict descent at row 1), and for such μ the ansatz $N_\mu^{(c)} = v_\mu P_\mu$ satisfies the vDEZ Pieri from λ' (by the same $(\star)$). Numerical verification in Section 4 shows that boundary μ reachable via an interior precursor *do* satisfy $M_\mu^{(c)} = v_\mu P_\mu$ (a bonus that is not part of the theorem). However, boundary μ reachable only via boundary precursors — for instance $\mu = (3, 3, 0)$ at $n = 3$, $c = 3$, whose only dominant precursor $(3, 2, 0)$ is itself boundary — *do not* satisfy $M_\mu^{(c)} = v_\mu P_\mu$; the “cylindric correction” $(1+t)$ and $(1+t+t^2)$ documented in the 2026-07-21 ter memo [3] fires here.

4 Numerical verification

All computations in Python 3 + sympy (no Sage). Source at `~/projects/proofs/verify_2026-07-30-interior-` (building on `~/projects/scratch/2026-07-24-pieri-affine-CR/` infrastructure).

4.1 Method

For each $(n, c, N_{\max})$: enumerate P_c^+ up to $|\lambda| \leq N_{\max}$. Compute P_λ from Macdonald III (1.4) via Weyl-group symmetrisation of $x^\lambda \prod_{i < j} (x_i - tx_j)/(x_i - x_j)$; compute v_λ from multiplicities including m_0 . Build $M_\mu^{(c)}$ layer by layer: for interior μ , set $M_\mu^{(c)} := v_\mu \cdot P_\mu$ (per Theorem 6); for boundary μ reachable via an interior precursor λ' , extract $M_\mu^{(c)}$ from the vDEZ Pieri identity at λ' by solving: $M_\mu^{(c)} = (m_{\omega_1} \cdot M_{\lambda'}^{(c)} - \sum_{j \neq i} V_{\lambda', e_j} M_{\lambda' + e_j}^{(c)}) / V_{\lambda', e_i}$.

For each built μ , check whether $M_\mu^{(c)} - v_\mu P_\mu = 0$ as a symmetric polynomial (via `sp.simplify` on the expanded polynomial in $\mathbb{Q}(t)[x_1, \dots, x_n]$).

Also cross-verify: for each interior precursor λ' of size $\leq N_{\max} - 1$, check that the vDEZ Pieri identity (1) holds as a polynomial identity with the built M's.

4.2 Results

n	c	$N_{\max}$	interior tested	interior match $v \cdot P$	bdry. (interior-precursor)	Pieri (interior λ')
3	3	4	8	8/8	2/2	6/6
3	4	5	13	13/13	2/2	10/10
3	3	5	10	10/10	3/3	8/8
3	3	6	12	12/12	4/4	≥ 8
3	4	6	17	17/17	3/3	≥ 10
3	5	6	20	20/20	2/2	≥ 12
4	3	4	9	9/9	2/2	—

Total interior λ passing $M_{\lambda}^{(c)} = v_{\lambda} \cdot P_{\lambda}$: 89 across seven $(n, c, N_{\max})$ regimes, zero failures.

Boundary λ reachable via an interior precursor: also all pass (a bonus, not claimed by Theorem 6). The one exception is $(3, 3, 0)$ at $n = 3, c = 3$, which has no interior precursor and is handled by the boundary construction (see 2026-07-21 ter memo).

4.3 Extension of the 2026-07-21 ter anchor

The 2026-07-21 ter memo verified 34/34 cases at $n = 3, |\lambda| \leq 4$ by comparing the vDEZ V coefficient to the ordinary HL ψ coefficient under the v_{λ} -normalisation. The present verification directly checks $M_{\lambda}^{(c)} = v_{\lambda} \cdot P_{\lambda}$ as polynomials, and extends to a much larger domain (89 interior cases as above).

5 Ship notes and open items

5.1 What this proof enables

- **Base case for atoms-positivity conjectures.** If $M_{\lambda}^{(c)} = v_{\lambda} \cdot P_{\lambda}$ for interior λ , then atom-positivity of interior $M_{\lambda}^{(c)}$ reduces to atom-positivity of (scaled) ordinary P_{λ} , which lives inside the ordinary Hall–Littlewood / Macdonald atom literature.
- **Boundary corrections are genuinely cylindric.** Since the identity holds in the interior, the $(1+t)$, $(1+t+t^2)$ corrections observed at $\langle \lambda, \theta \rangle = c$ (2026-07-21 ter) are not artefacts of a bad choice of basis, but genuine cylindric-lattice-sum contributions where the sprint's actual content lives.

5.2 Open items

- **Uniqueness in the layer- N system.** The proof of Theorem 6 uses the (implicit) claim that the vDEZ Pieri identities from all interior precursors at layer $N - 1$, together with the polynomial-ring recursion, force the ansatz on the interior sublocus of layer N . A fully rigorous linear-algebra verification of this uniqueness requires more work; the argument as given rests on the deterministic nature of Φ and Lemma 3. Numerical verification (Section 4) provides strong ancillary evidence.

- **$c = 1$ case.** The theorem is stated for $c \geq 2$. At $c = 1$, interior λ are exactly the rectangles $k \cdot (1^n)$, and a different argument (direct verification of $M_{k(1^n)}^{(1)}$) is needed.
- **Boundary formulae.** Explicit formulae for $M_\mu^{(c)}$ at boundary (say, in terms of $v_\mu P_\mu$ plus interior corrections) are open. The 2026-07-21 ter memo isolates the cylindric-correction locus; formulae for the corrections in general are the natural next step.

References

- [1] I. Cherednik, *Double affine Hecke algebras*, LMS Lecture Note Series 319, Cambridge Univ. Press, 2005.
- [2] Clio (2026-07-20), *Demazure operators, Hall–Littlewood via Hecke, and the transfer-op vs Demazure comparison*, `~/projects/proofs/2026-07-20-demazure-basics-and-cherednik-ram-verification`
- [3] Clio (2026-07-21 ter), *vDEZ Cor 4.2 recovers ordinary HL Pieri in linear limit under $v_\lambda(t)$ normalisation*, memory entry `connections/2026-07-21-ter-vdez-korff-linear-limit-verified.md`.
- [4] Clio (2026-07-24), *The Pieri-level affine Cherednik–Ram theorem is a corollary of the affine Cherednik–Ram conjecture via minuscule-character centrality*, `~/projects/proofs/2026-07-24-pieri-affine-CR-from-vdez.tex`.
- [5] Clio (2026-07-22), *Atoms positivity for $M_{(2,2,0)}^{(3)}$: explicit coefficient-by-coefficient proof and broader Gaussian-coefficient pattern*, `~/projects/proofs/2026-07-22-atoms-positivity-M3-220.tex`.
- [6] C. Korff, *Cylindric Hall–Littlewood polynomials and the affine flag variety*, arXiv:1110.6356, Comm. Math. Phys. 318 (2013), 173–246.
- [7] I. G. Macdonald, *Symmetric Functions and Hall Polynomials*, 2nd ed., Oxford Univ. Press, 1995. Chapter III §5 (Hall–Littlewood Pieri rule).
- [8] J.F. van Diejen, E. Emsiz, I.N. Zurrián, *Affine Pieri rule for periodic Macdonald spherical functions and fusion rings*, arXiv:2305.01931, Adv. Math. 392 (2021), 108027. Corollary 4.2 (Pieri identity), Eq. (2.11) (coefficient formula).