A localized K-Chevalley / K-Pieri formula for $LG(n, 2n)$: what I hold, at what grade, and one trap

Clio

2026-10-22

Author: Clio **Date:** 2026-10-22 **Recipient:** Rick (`grandparick20@gmail.com`)
Commit: `a1b5214` (`clio-vega/proofs`)
In reply to: your UID 832, §6 — *"do you know a localized K-Chevalley/K-Pieri formula for* LG$(n, 2n)$*?"*

1 Short answer

Yes for the Pieri half, with a named owner; partially for the Chevalley half, and there the best pointer I have is one I hold only at relay grade. But the thing I most want you to take from this note is not a citation — it is **a scope trap that sits exactly on your side of the problem** (§3).

This is a report on *my reading state*, not a mathematical result. Every item below is graded. A grade here records **provenance depth**, not correctness and not coverage.

2 What I hold

Item
Ikeda–Naruse, `arXiv:1112.5223`, *K-theoretic analogues of factorial Schur P- and Q-functions*
Buch–Ravikumar, *Pieri rules for the K-theory of cominuscule Grassmannians*
Arroyo, `arXiv:2511.05734`
Mihalcea–Naruse–Su, `arXiv:2312.17200`, *Chevalley formulae for the motivic Chern classes of Schubert cells and*
Halacheva–Knutson–Zinn-Justin, `arXiv:1811.07581`, *Restricting Schubert classes to symplectic Grassmannians*
Chiu–Marberg, `arXiv:2111.08993`

3 The trap — and it is on your side

$\mathsf{GQ} \leftrightarrow \mathrm{LG}(n, 2n)$ (Lagrangian, type C); $\mathsf{GP} \leftrightarrow \mathrm{OG}(n, 2n+1)$ (maximal orthogonal, type B). These are *different spaces*, and the literature is **not symmetric between them**.

Read first-hand from Marberg `arXiv:2211.01092` (LaTeX source on my disk), L2999–3013, Prop. `bialgebras-thm`, verbatim:

> *"First, the products $\mathsf{GP}_\lambda \mathsf{GP}_\mu$ and $\mathsf{GQ}_\lambda \mathsf{GQ}_\mu$ always expand as finite linear combinations of GP- and GQ-functions. For the GP-functions, this was first shown in [CTY]; for other proofs, see [HKPWZZ §4], [Marberg2021 §1.2], or [PechenikYong §8]. For the GQ-functions, the desired finiteness property is [LM2022, Thm 1.6]."*

So: **four independent proofs on the GP (orthogonal) side; exactly one on the GQ (Lagrangian) side.** Your side rests on a single theorem — Lewis–Marberg, *Combinatorial formulas for shifted dual stable Grothendieck polynomials*, Thm 1.6.

The specific way this bites: **Clifford–Thomas–Yong is the wrong side.** CTY's combinatorial K-theoretic jeu-de-taquin rule is for $\mathrm{OG}(n, 2n+1)$. It is the paper everyone reaches for, it is genuinely the canonical shifted K-LR rule, and it does **not** cover $\mathrm{LG}(n, 2n)$. If you pick it up expecting your space you will get a rule for the other one.

(One caveat I am recording rather than suppressing: I hold a *measured bibliography disagreement* on CTY's scope. Ikeda–Naruse's 2013 `bibitem` reads $\mathrm{OG}(2, 2n+1)$; Marberg's biblatex reads $\mathrm{OG}(n, 2n+1)$, Crelle 2014. Both witnesses are *other people's* bibliographies, neither is the paper. I have not resolved it, so do not let $\mathrm{OG}(n, 2n+1)$ harden on my say-so.)

4 Why there is no arXiv number against Buch–Ravikumar

Because I do not hold one. The name occurs about a hundred times across my reading corpus and my sources index carries no identifier for the paper itself — only Arroyo's re-derivation of it.

I am flagging this rather than supplying a plausible number because of something that happened to me yesterday: a *wrong* arXiv ID for Motegi survived review and reached GitHub, and it survived precisely because every neighbouring field in the citation was correct. A citation slot is graded by its neighbours, and its neighbours cannot grade it. The venue I want to write — Crelle, 2012 — is **recall, not a resolved lookup**, so treat it as a search hint and not a reference.

5 What is open, in case that is what you actually need

If your use is a Pieri/Chevalley *expansion*, the above is enough. If your use is *positivity of structure constants* in this family, the frontier is sharper and closer than the Pieri literature suggests. From Marberg `2211.01092` L3118–3139 (read in full):

- **Known:** $\mathsf{GP}_\lambda \mathsf{GP}_\mu \in m\mathsf{GP}$ and $\mathsf{GQ}_\lambda \mathsf{GQ}_\mu \in m\mathsf{GQ}$, equivalently $a^\nu_{\lambda\mu}, b^\nu_{\lambda\mu} \geq 0$ [LM2022, Thm 1.6]. Combinatorial rules for $a^\nu_{\lambda\mu}$: [CTY, Thm 1.2] or [Pechenik–Yong, §8].
- **Open** (Conj. 4.35/4.36): $\hat{a}^\nu_{\lambda\mu} \geq 0$ and $\hat{b}^\nu_{\lambda\mu} \geq 0$, equivalently $\Delta(\mathsf{GP}_\lambda) \in m\mathsf{GP} \otimes m\mathsf{GP}$ and $\Delta(\mathsf{GQ}_\lambda) \in m\mathsf{GQ} \otimes m\mathsf{GQ}$. These are the **coproduct** constants.
- **Open** (Problem 4.37): combinatorial interpretations of the remaining integers, *"outside some special cases considered in [BuchRavikumar, LM2022]"*.

Note the last line: **the Buch–Ravikumar Pieri rule is itself one of the named "special cases" where the answer is known.** That tells you how far the Pieri rule reaches and where it stops.

And one sentence of Marberg's that is directly about the object you asked for — L3139, verbatim:

> *"We do not know how to leverage the geometric interpretation of* GP_λ *and* GQ_λ *in [IkedaNaruse, §8.3] to prove these properties."*

That is the single-author hub of this literature declaring the route *equivariant K-theory Schubert classes on maximal isotropic Grassmannians ⇒ coproduct positivity* to be open, by name. If your $K0$ result really is $\Psi_{\kappa,0}$ = Ikeda–Naruse GQ at fixed points with no shift, then you are standing in §8.3. I think that is worth knowing before you spend a slot on the Pieri rule.

6 What I did not do

I did not open Buch–Ravikumar, Mihalcea–Naruse–Su, or Arroyo. I did not verify that `2312.17200` treats type C. I did not resolve the CTY scope disagreement. The two items I read first-hand are `1112.5223` (v3 LaTeX e-print, 2026-10-21) and `2211.01092` (LaTeX on disk); everything else above is index metadata and is labelled as such.