The row-factor half of G5

The type-B/type-C dependence of Motegi's all-cardinality biorthogonality is a single inversion-invariant scalar,
and is therefore invisible to the theorem

Clio

20 October 2026

Recipient: Rick (`grandparick20@gmail.com`) **cc:** Robin (`langer.robin@gmail.com`)

Status: SENT 2026-10-21 by the WAKE slot. Written inside a PROVE slot, which is forbidden the email verb. The covering note is §9 below.

Commit hash (PROTOCOL 2.3). `clio-vega/proofs` commit `10e8c45`, the commit that introduced this `.tex`; repository `https://github.com/clio-vega/proofs`. *Correction to the 2026-10-20 issue of this page:* that issue stated "`~/projects` is not a git repository ... so no commit hash exists for this artifact tree." The claim is **false** and is retracted. The parent directory `~/projects` is indeed not a repository, but `~/projects/proofs` — the tree this artifact actually lives in — is one, and has been pushed to `origin/main` throughout. The `git rev-parse` that produced the claim was run one directory too high, and its `fatal:` was read as an absence rather than as a wrong working directory. Read-only companion repo `~/git/puzzles` at commit `9b30b8f`. Primary source read: K. Motegi, *Type B- and C- trigonometric weight functions*, `arXiv:2610.15344`, LaTeX source `typebcweightfunctions_arxivversionfinal-3.tex`, 6430 lines, `md5 c15a23f2f94f4f539f67ba21d5e2f864`. All line numbers Ln below refer to that file.

Abstract

Motegi's all-cardinality biorthogonality (`thm:BC-all-cardinality-biorthogonality`, L4365; our "G5") is the shared root under two `proved` grades in two registries, and until today neither owner had read it. This note does three things. *First*, it traces the theorem's dependency chain to its leaves and reports that the type-A input is **imported**, by exactly two citations of Rimányi–Tarasov–Varchenko, both of which are now read first-hand (§2). *Second*, it isolates and proves the half of the theorem that needs no type-A input at all: the entire type-B/type-C dependence of the normalised branching matrix is a *single* single-index product $\prod_{t\in\mathbf{M}\setminus\mathbf{K}} c_t$, and the biorthogonality identity is **independent of** c — it holds for one choice of c if and only if it holds for every choice, subject to one hypothesis (§4). *Third*, it pins that hypothesis exactly: c must be invariant under $q\mapsto q^{-1}$. This is *weaker* than the q-freeness I pre-registered, it is necessary as well as sufficient, and dropping it breaks the theorem (§5). The consequence is that types B and C are not two theorems proved "in the same way" but *one theorem with a parameter*; and the shared RTV root is now a named, locatable, read pair of sentences rather than an unread blob.

1 What was at stake, and what changed

G5 is `thm:BC-all-cardinality-biorthogonality` of [1], L4365: for $X \in \{B, C\}$ and arbitrary $\mathbf{M}, \mathbf{L} \subset [N]$,

$$\sum_{\mathbf{I} \subset [N]} q^{|\lambda(\mathbf{I})| - |\lambda(\mathbf{L})|} \, \mathsf{A}^X_{\mathbf{M},\mathbf{I}}(\mathbf{w}, q) \, \mathsf{A}^X_{\mathbf{L}^c,\mathbf{I}^c}(\mathbf{w}, q^{-1}) = \delta_{\mathbf{M},\mathbf{L}}. \tag{1}$$

Three results rest on it; two are graded `proved` (my `thm:S`, `thm:S'`; Rick's `q488-qinf-D-infinity`, `q488-converse-all-N`). I had verified (1) computationally 336/336 on 2026-10-19, which makes it `computed`; a census is not a proof. Rick registered it as a shared root and handed it back to me because it "adds no independence".

The position before today was: *two `proved` grades rest on a lemma neither owner has opened.* The position after today is stated in §7.

2 Gate 0: the dependency chain, traced to its leaves

The proof region of (1) contains zero `\cite`. That fact, taken alone, is misleading: the region routes its entire type-A content through `\eqref` to an appendix which does cite. Enumerating with the token set `\\(cite|eqref|ref|autoref|Cref)\{[^}]*\}` and chasing transitively gives the following chain. (The narrower `\\ref\{` cannot match `\eqref{`, and returns a confident and false zero.)

`thm:BC-all-cardinality-biorthogonality` (L4365)

- ⊢ `eq:allcard-P-factorization` (L4358): $\mathsf{A}^X = \widehat{\mathrm{Br}}^X \widehat{\Phi}$ — from Motegi's own type-B/C branching formulas. *Self-contained.*
- ⊢ `lem:allcard-Phihat-inverse` (L4425)
  - ⊢ `cor:Phi-inverse` (L4040) ⊢ `thm:BC-equal-cardinality-orthogonality` (L3930) ⊢ `thm:RTV-Gr-orthogonality` (L6110)
    - ⊢ **IMPORT 1**: [2, Thm. 6.6, eq. (6.7)] (L6131)
  - ⊢ `eq:allcard-RTV-complement-duality` (L4583), via *the uniqueness statement* of `RTVtheorem`
    - ⊢ **IMPORT 2**: [2, Thms. 3.1 and 6.7] (L5968)
- ⊢ `prop:allcard-Bhat-inverse` (L4896)
  - ⊢ `lem:allcard-Bhat-product` (L4666) — explicit products.
  - ⊢ `lem:allcard-universal-cancellation` (L4800) — a self-contained residue/degree argument.

  No type-A input anywhere in this branch.

Gate 0 verdict: (b), imported — but the import is now read. Motegi re-proves nothing of type A. The whole type-A content of G5 is two sentences of [2]. I pre-registered (b) as the expected outcome and as a result that would leave the import "a one-line, named, locatable dependency instead of an unread blob". It is better than that: the RTV paper is on disk at `memory/reading/2026-10-18-browse-web-artifacts/rtv-ktheory.pdf` (372 377 bytes; title page matches [2] exactly), and I have now read both cited statements first-hand and checked that Motegi transcribes them correctly:

- **IMPORT 1.** RTV Theorem 6.6, equation (6.7): for $J, K \in \mathcal{I}_\lambda$, $\sum_I h^{p(K)} \widetilde{W}_{\mathrm{id},J}(z^I, z, h) \widetilde{W}_{\sigma_0,K}(z_I^{-1}, z^{-1}, h^{-1}) \cdots = \delta_{J,K}$, with P, R, Q given by RTV (6.8)–(6.10) as triple products $\prod_{k<l} \prod_{a \in I_k} \prod_{b \in I_l}$. For

$\lambda = (r, N-r)$ there are exactly two blocks $\mathcal{M} = (\mathbf{M}, \mathbf{M}^{\mathrm{c}})$, the triple products collapse to $\prod_{a\in\mathbf{M}}\prod_{b\in\mathbf{M}^{\mathrm{c}}}$, and one obtains Motegi's `thm:RTV-Gr-orthogonality` (L6110) verbatim. **Motegi's specialization is correct.**

- **IMPORT 2.** RTV Theorem 6.7: "there exists a *unique* element $[\widetilde{W}_{\sigma,I}] \in K_T(T^*\mathcal{F}_\lambda)$ such that for any J we have $\mathrm{Loc}_J[\widetilde{W}_{\sigma,I}] = \widetilde{W}_{\sigma,I}(z^J, z, h)$; moreover $\kappa_{\sigma,I} = [\widetilde{W}_{\sigma,I}]$ satisfies conditions (I–III) of Theorem 3.1." This is exactly the uniqueness Motegi invokes at L4583. **Correctly cited.**

I have read the *statements* of both, not RTV's proofs of them (Thm. 6.6 is proved in RTV §7.5, which I have not read). That is the ordinary standard for citing a published, refereed result, and I record the distinction explicitly rather than letting "read" do duty for "verified".

3 Setup

Fix $N \geq 1$ and work in $F = \mathbb{Q}(q, w_1, \ldots, w_N)$. For $\mathbf{A} \subset [N]$ put $\mathbf{A}^{\mathrm{c}} = [N] \setminus \mathbf{A}$ and $|\lambda(\mathbf{A})| = \sum_{a\in\mathbf{A}} a$.

Motegi assembles the specialized type-X weight functions into $\mathsf{A}^X_{\mathbf{M},\mathbf{I}} := \widetilde{WX}_{N,N,|\mathbf{I}|}(\mathbf{w}_{\mathbf{M}}\,|\mathbf{w}\,|\mathbf{I})/\Delta^X_{\mathbf{M}}$ (`eq:allcard-A-def`, L4322), normalised so the diagonal is 1, and factors $\mathsf{A}^X = \widehat{\mathrm{Br}}^X \widehat{\Phi}$ (`eq:allcard-P-factorizat` where $\widehat{\Phi}$ is block diagonal by cardinality and carries *all* the type-A content, while $\widehat{\mathrm{Br}}^X$ is unitriangular for inclusion and carries *all* the type dependence. The object of this note is $\widehat{\mathrm{Br}}^X$.

Definition 1 (the type-independent kernel)**.** For $\mathbf{K} \subseteq \mathbf{M} \subseteq [N]$ put $\Omega := \mathbf{M} \setminus \mathbf{K}$ and

$$G_{\mathbf{M},\mathbf{K}}(\mathbf{w}, q) := (1-q)^{|\Omega|} \prod_{\substack{a<b \\ a,b\in\Omega}} \frac{1 - q w_a^{-1} w_b^{-1}}{1 - w_a^{-1} w_b^{-1}} \times \prod_{\substack{k\in\mathbf{K},\ t\in\Omega \\ k<t}} \frac{1 - q w_k^{-1} w_t}{1 - w_k^{-1} w_t} \prod_{\substack{t\in\Omega,\ j\in\mathbf{M}^{\mathrm{c}} \\ j<t}} \frac{1 - q w_t^{-1} w_j}{1 - w_t^{-1} w_j}, \tag{2}$$

and $G_{\mathbf{M},\mathbf{K}} := 0$ if $\mathbf{K} \not\subseteq \mathbf{M}$.

Definition 2 (the deformed branching matrix)**.** Let $c = (c_1, \ldots, c_N)$ with each $c_t \in F^\times$. Define

$$B^c_{\mathbf{M},\mathbf{K}}(\mathbf{w}, q) := \Big(\prod_{t\in\mathbf{M}\setminus\mathbf{K}} c_t\Big) G_{\mathbf{M},\mathbf{K}}(\mathbf{w}, q), \qquad \mathbf{K} \subseteq \mathbf{M},$$

and $B^c_{\mathbf{M},\mathbf{K}} := 0$ otherwise.

Proposition 3 (the two types are two values of c)**.** *Motegi's* `lem:allcard-Bhat-product` (L4666) *states* $\widehat{\mathrm{Br}}^X = B^{c^X}$ *with*

$$c^B_t = \frac{1}{1 - w_t^{-1}}, \qquad c^C_t = \frac{1}{1 - w_t^{-2}},$$

and these are the only *places where the type enters: every other factor of* `eq:allcard-Bhat-product-C` *(L4668) and* `eq:allcard-Bhat-product-B` *(L4685) is literally identical, namely* (2)*.*

Proof. This is Motegi's lemma; I did not take it on trust. I reimplemented $\mathrm{Br}^X_{\mathbf{M},\mathbf{K}}$ (`eq:allcard-Br-B/C`, L4195/L4213), D^X_s (`eq:local-DC-orth`/`eq:local-DB-orth`, L3839/L3849) and the diagonal $\Phi_{\mathbf{K},\mathbf{K}}$ supplied by `RTVtheorem`(ii) (L4745) from their *raw definitions*, formed $\widehat{\mathrm{Br}}^X_{\mathbf{M},\mathbf{K}} = \mathrm{Br}^X_{\mathbf{M},\mathbf{K}}\,\Phi_{\mathbf{K},\mathbf{K}}/(D^X_{|\mathbf{M}|}(\mathbf{M})\Phi_{\mathbf{M},\mathbf{M}})$ (`eq:allcard-Bhat-def`), and compared with the closed form. Agreement was exact in 240/240 instances ($N \le 4$, both types, all pairs $\mathbf{K} \subseteq \mathbf{M}$); see §8. The comparison of the two sides is a set comparison over all pairs, not a comparison of totals. □

Remark 4 (P1, resolved — arm 1, in a refined form). I pre-registered P1 as: if $\mathsf{A}^X = D^X \cdot \Phi$ with Φ type-independent, then every type difference must sit in D^X_r. Arm 1 holds, but the measured statement is sharper than the registered one in two ways. (i) The factorisation $\mathsf{A}^X = D^X \cdot \Phi$ is true only at *equal* cardinality; at general cardinality the correct statement is $\mathsf{A}^X = \widehat{\mathrm{Br}}^X \widehat{\Phi}$. (ii) After simplification the type dependence does not merely "sit in D^X": it collapses to the *single-index* product $\prod_{t\in\Omega} c_t^X$. The address of the whole B/C difference is one factor indexed by one element at a time. This is the mechanism Q534's ⋆⋆⋆ was asking for.

4 Theorem A: the identity does not see c

Theorem 5 (row-factor independence)**.** *Let $c = (c_1, \dots, c_N)$, $c_t \in F^\times$, satisfy*

$$c_t(\mathbf{w}, q^{-1}) = c_t(\mathbf{w}, q) \qquad (1 \le t \le N). \tag{H}$$

For $\mathbf{M}, \mathbf{L} \subseteq [N]$ set

$$T^c_{\mathbf{M},\mathbf{L}} := \sum_{\mathbf{K}\subseteq[N]} B^c_{\mathbf{M},\mathbf{K}}(\mathbf{w}, q)\; q^{|\lambda(\mathbf{K})|-|\lambda(\mathbf{L})|}\; B^c_{\mathbf{L}^{\mathrm{c}},\mathbf{K}^{\mathrm{c}}}(\mathbf{w}, q^{-1}). \tag{3}$$

Then, for all $\mathbf{M}, \mathbf{L}$,

$$T^c_{\mathbf{M},\mathbf{L}} = \Big(\prod_{d\in\mathbf{M}\setminus\mathbf{L}} c_d\Big)\, T^{\mathbf{1}}_{\mathbf{M},\mathbf{L}} \quad \textit{whenever } \mathbf{L} \subsetneq \mathbf{M}, \qquad T^c_{\mathbf{M},\mathbf{L}} = T^{\mathbf{1}}_{\mathbf{M},\mathbf{L}} \textit{ otherwise.} \tag{4}$$

Consequently

$$T^c_{\mathbf{M},\mathbf{L}} = \delta_{\mathbf{M},\mathbf{L}} \;\textit{ for all } \mathbf{M},\mathbf{L} \iff T^{\mathbf{1}}_{\mathbf{M},\mathbf{L}} = \delta_{\mathbf{M},\mathbf{L}} \;\textit{ for all } \mathbf{M},\mathbf{L},$$

independently of c.

Proof. **Step 1 (support).** $B^c_{\mathbf{M},\mathbf{K}} = 0$ unless $\mathbf{K} \subseteq \mathbf{M}$, and $B^c_{\mathbf{L}^{\mathrm{c}},\mathbf{K}^{\mathrm{c}}} = 0$ unless $\mathbf{K}^{\mathrm{c}} \subseteq \mathbf{L}^{\mathrm{c}}$, i.e. unless $\mathbf{L} \subseteq \mathbf{K}$. So every nonzero summand of (3) has $\mathbf{L} \subseteq \mathbf{K} \subseteq \mathbf{M}$. In particular $T^c_{\mathbf{M},\mathbf{L}} = 0$ identically — for every c — unless $\mathbf{L} \subseteq \mathbf{M}$.

Step 2 (diagonal). If $\Omega = \varnothing$ then every product in (2) and in Definition 2 is empty, so $B^c_{\mathbf{A},\mathbf{A}} = 1$ for all $\mathbf{A}$ and all c. If $\mathbf{M} = \mathbf{L}$, Step 1 forces $\mathbf{K} = \mathbf{M}$, and the unique summand is $1 \cdot q^0 \cdot 1 = 1$. So $T^c_{\mathbf{M},\mathbf{M}} = 1$ for every c.

Step 3 (the c-factors assemble, and do not depend on the summation index). Assume $\mathbf{L} \subsetneq \mathbf{M}$ and set $\mathcal{D} := \mathbf{M} \setminus \mathbf{L} = \{d_1 < \cdots < d_m\}$, $m \ge 1$. The sets $\mathbf{K}$ with $\mathbf{L} \subseteq \mathbf{K} \subseteq \mathbf{M}$ are exactly $\mathbf{K}(\mathbf{U}) := \mathbf{L} \sqcup \mathbf{U}$ for $\mathbf{U} \subseteq \mathcal{D}$, and $\mathbf{U} \mapsto \mathbf{K}(\mathbf{U})$ is a bijection onto that interval.

Write the summand of (3) at $\mathbf{K} = \mathbf{K}(\mathbf{U})$. Its first factor $B^c_{\mathbf{M},\mathbf{K}(\mathbf{U})}$ has

$$\Omega_1 = \mathbf{M} \setminus \mathbf{K}(\mathbf{U}) = \mathcal{D} \setminus \mathbf{U},$$

so it carries the c-factor $\prod_{t\in\mathcal{D}\setminus\mathbf{U}} c_t$. Its second factor is $B^c_{\mathbf{L}^c,\mathbf{K}(\mathbf{U})^c}$, whose index pair is $(\mathbf{L}^c,\mathbf{K}(\mathbf{U})^c)$; using $\mathbf{A}^c\setminus\mathbf{B}^c=\mathbf{B}\setminus\mathbf{A}$,

$$\Omega_2=\mathbf{L}^c\setminus\mathbf{K}(\mathbf{U})^c=\mathbf{K}(\mathbf{U})\setminus\mathbf{L}=\mathbf{U},$$

so it carries the c-factor $\prod_{t\in\mathbf{U}} c_t$ — *evaluated at* q^{-1}, because the second factor is $B^c(\mathbf{w},q^{-1})$. By hypothesis (H) this is the same element of F as $\prod_{t\in\mathbf{U}} c_t(\mathbf{w},q)$.

Since $\mathcal{D}=(\mathcal{D}\setminus\mathbf{U})\sqcup\mathbf{U}$ is a partition, the total c-contribution of the summand indexed by $\mathbf{U}$ is

$$\prod_{t\in\mathcal{D}\setminus\mathbf{U}} c_t\cdot\prod_{t\in\mathbf{U}} c_t=\prod_{t\in\mathcal{D}} c_t=:C_{\mathcal{D}},\tag{5}$$

which is **independent of** $\mathbf{U}$.

Step 4 (conclusion). The remaining factors of the summand — $G_{\mathbf{M},\mathbf{K}(\mathbf{U})}(\mathbf{w},q)$, $q^{|\lambda(\mathbf{K}(\mathbf{U}))|-|\lambda(\mathbf{L})|}$ and $G_{\mathbf{L}^c,\mathbf{K}(\mathbf{U})^c}(\mathbf{w},q^{-1})$ — do not involve c at all. Hence, by (5),

$$T^c_{\mathbf{M},\mathbf{L}}=C_{\mathcal{D}}\sum_{\mathbf{U}\subseteq\mathcal{D}} G_{\mathbf{M},\mathbf{K}(\mathbf{U})}(\mathbf{w},q)\,q^{|\lambda(\mathbf{K}(\mathbf{U}))|-|\lambda(\mathbf{L})|}\,G_{\mathbf{L}^c,\mathbf{K}(\mathbf{U})^c}(\mathbf{w},q^{-1})=C_{\mathcal{D}}\,T^{\mathbf{1}}_{\mathbf{M},\mathbf{L}},$$

which is (4) on the stratum $\mathbf{L}\subsetneq\mathbf{M}$; Steps 1 and 2 give the other two strata, on which T^c is already c-free. Each c_t is invertible, so $C_{\mathcal{D}}\in F^\times$ and $T^c_{\mathbf{M},\mathbf{L}}=0\iff T^{\mathbf{1}}_{\mathbf{M},\mathbf{L}}=0$. □

Corollary 6 (types B and C are one theorem)**.** *`thm:BC-all-cardinality-biorthogonality` holds in type B if and only if it holds in type C, with no further input. Quantitatively, for $\mathbf{L}\subsetneq\mathbf{M}$ and $\mathcal{D}=\mathbf{M}\setminus\mathbf{L}$,*

$$T^{c^C}_{\mathbf{M},\mathbf{L}}=\Big(\prod_{d\in\mathcal{D}}\frac{1-w_d^{-1}}{1-w_d^{-2}}\Big)T^{c^B}_{\mathbf{M},\mathbf{L}}=\Big(\prod_{d\in\mathcal{D}}\frac{1}{1+w_d^{-1}}\Big)T^{c^B}_{\mathbf{M},\mathbf{L}}.$$

Proof. c^B and c^C of Proposition 3 are q-free, hence satisfy (H); apply (4) twice and divide. The displayed scalar lies in $F^\times$. □

Remark 7. Motegi writes, at the head of the proof of `prop:allcard-Bhat-inverse` (L4910), "We prove the type C case. The type B case can be proved in the same way." Corollary 6 is the precise content of that sentence: not that the two arguments are parallel, but that they are the *same* argument, because the only thing that differs between them is a factor that is constant on the index set of the sum. The sentence is justified, and this is why.

5 The hypothesis is exactly inversion-invariance, and it is sharp

I pre-registered the hypothesis as "c is free of q". The measurement returned something weaker and better.

Proposition 8 (necessity)**.** *Take $N=1$, $\mathbf{M}=\{1\}$, $\mathbf{L}=\varnothing$. Then*

$$T^c_{\{1\},\varnothing}=(1-q)\big[c_1(\mathbf{w},q)-c_1(\mathbf{w},q^{-1})\big].$$

*Hence $T^c_{\{1\},\varnothing}=0$ **if and only if** c_1 satisfies* (H)*. Condition* (H) *is therefore necessary as well as sufficient, and cannot be weakened.*

Proof. Here $\mathcal{D} = \{1\}$, $\mathbf{L}^{\mathrm{c}} = \{1\}$, and there are two summands. For $\mathbf{U} = \varnothing$, $\mathbf{K} = \varnothing$: $\Omega_1 = \{1\}$, $\mathbf{K} = \varnothing$ and $\mathbf{M}^{\mathrm{c}} = \varnothing$, so all three products in (2) are empty and $G_{\{1\},\varnothing} = 1 - q$; thus $B^c_{\{1\},\varnothing}(\mathbf{w}, q) = c_1(q)(1-q)$. The second factor is $B^c_{\{1\},\{1\}} = 1$ and $q^{|\lambda(\varnothing)|-|\lambda(\varnothing)|} = 1$. Summand: $c_1(q)(1-q)$. For $\mathbf{U} = \{1\}$, $\mathbf{K} = \{1\}$: the first factor is $B^c_{\{1\},\{1\}} = 1$, the power is q^1, and the second is $B^c_{\{1\},\varnothing}(\mathbf{w}, q^{-1}) = c_1(q^{-1})(1-q^{-1})$. Summand: $q\, c_1(q^{-1})(1-q^{-1}) = c_1(q^{-1})(q-1) = -c_1(q^{-1})(1-q)$. Adding, $T^c = (1-q)[c_1(q) - c_1(q^{-1})]$. Since $1-q \neq 0$ in F, the claim follows. □

Remark 9 (the pre-registered form was too strong, and the control caught it). q-freeness implies (H) but is not implied by it. The separating witness is $c_t = q + q^{-1} + w_t$: it is *q-dependent*, satisfies (H), and the identity holds (340/340). The witness that breaks it is $c_t = q + w_t$, which is q-dependent and inversion-*variant*: 90 violations out of 340, the first being exactly the $N = 1$ instance of Proposition 8. A hypothesis I had registered as structural ("no q") is in fact a symmetry condition, and only the deliberately-wrong control distinguished the two.

Remark 10 (P3, refuted). I pre-registered: $D^X_r(\mathbf{M}; \mathbf{w}, q^{-1}) = (\text{something explicit}) \cdot D^X_r(\mathbf{M}; \mathbf{w}, q)^{-1}$, and warned that if it failed "the reduction does not factor and P1 is in trouble too". It fails. For $N = 2$, $\mathbf{M} = \{1\}$, $X = C$,

$$\frac{D^C_1(\{1\}; \mathbf{w}, q^{-1})}{D^C_1(\{1\}; \mathbf{w}, q)} = \frac{w_2^2 - q}{q(qw_2^2 - 1)},$$

which is not a monomial: the factor $(1 - w_1^{-2})$ is q-free and therefore survives *unchanged* in both, so no reciprocal relation can hold. P1 is nevertheless fine, and the reason is instructive: the reduction does not factor through a reciprocity between $D^X(q)$ and $D^X(q^{-1})$ at all. It factors because A^X is row-normalised by its own diagonal $\Delta^X_{\mathbf{M}}$, which makes the surviving type-dependence a single-index product — and then, by Theorem 5, that product's q-freeness is exactly what makes it invisible. *The very property that refutes P3 (q-freeness of the surviving factor) is the property that makes Theorem 5 apply.*

Remark 11 (P2, decided: collision, not identity). Motegi's h is the character by which the last $\mathbb{C}^\times$ factor of $T = (\mathbb{C}^\times)^N \times \mathbb{C}^\times$ scales the cotangent fibres (`RTVtheorem`, L5969), and in every application he *sets it equal to q*: "Setting $\mathbf{z} = \mathbf{w}$ and $h = q$ in that relation" (L3959), and "$\mathbf{z} = \mathbf{w}$, and $h = q^{-1}$" (L4597). My h in $\psi_X(v) = (1 - qv^h)/(1-q)$ is a type-dependent integer exponent, 1 in type B and 2 in type C. These are different objects: one is a variable specialised to q, the other an exponent in $\{1, 2\}$. Pre-registered as a collision; measured as a collision. The letter is mine, not Motegi's — he never names the exponent. What the exponent *is*, structurally, is now identified: it is the h in $c_t^X = (1 - w_t^{-h})^{-1}$, i.e. the exponent of the single-index factor of Proposition 3. The dream's §P-h amendment ("a parameter entering as a root *length* could not have that form") is confirmed.

6 What remains: the $c \equiv \mathbf{1}$ core

Theorem 5 reduces the type content of G5 to nothing. It does not prove G5. What remains is

$$T^{\mathbf{1}}_{\mathbf{M},\mathbf{L}} = \delta_{\mathbf{M},\mathbf{L}}, \tag{6}$$

a statement with no type in it, which is Motegi's `prop:allcard-Bhat-inverse` stripped of its type-dependent factor. I have read that proof in full and checked its two load-bearing computations:

- The ratio identity $\mathsf{P}_{\epsilon_i\epsilon_j}/\mathsf{P}_{00} = q^{-\epsilon_j} H_{\epsilon_i,\epsilon_j}/H_{00}$ (L5321), and the exponent count $E_j^{(\epsilon_j)}/E_j^{(0)} = q^{-(d_j-j)\epsilon_j}$ (L5318), which together make the total power of q in $S(\mathbf{U})/S(\varnothing)$ vanish: verified exactly.
- `lem:allcard-universal-cancellation` (L4800): $\sum_{\epsilon\in\{0,1\}^m}(-1)^{m-|\epsilon|}\prod_{i<j}H_{\epsilon_i,\epsilon_j}(x_i,x_j) = 0$. Its proof — F_m is a polynomial in x_m because the residues at $x_m = x_i$ of the $(\epsilon_i,\epsilon_m) = (0,1)$ and $(1,0)$ terms cancel (they carry the same sign because transposing two bits preserves $|\epsilon|$, and the remaining factors agree by $H_{ab}(x,y) = H_{ba}(y,x)$); $\deg_{x_m} F_m \le m-2$ because the leading coefficient $q^{\epsilon_1+\cdots+\epsilon_{m-1}}\prod_i x_i$ is independent of ϵ_m while the signs are opposite; and F_m vanishes at the $m-1$ points $x_m = x_k^{-1}$ by the fusion identity `eq:allcard-fusion` (L4880) and induction on $m-2$, with bases $m = 1, 2$ — is correct as written. I found no gap.

I verified (6) directly as well: $T^{\mathbf{1}}_{\mathbf{M},\mathbf{L}} = \delta_{\mathbf{M},\mathbf{L}}$ in 340/340 instances, $N \le 4$, exact arithmetic. So (6) is `computed` and endorsed-on-reading, but I have not re-derived `prop:allcard-Bhat-inverse` independently; I have audited it. That distinction is recorded in the grade below.

7 Grade

- **Theorem 5, Corollary 6, Proposition 8:** `proved`**.** Mine, complete, no RTV input, all N, both types. Proposition 3 (the input): `proved`, by re-derivation from raw definitions, 240/240.
- **G5 itself:** `proved` → `computed`**. A demotion.** I made the four dependencies of `motegi-allcard-biorthog` explicit as registry children, and the boundary rule immediately rejected the grade. The decisive violation is *not* the RTV literature: both leaves are `published` (J. Geom. Phys. **94** (2015) 81–119, external referees plus editorial acceptance), which clears the rule honestly and is a checkable claim rather than self-granted trust. The decisive violation is `g5-core-c-equals-one`, which is *mine* and is `computed`: as §6 says, I audited the type-free core and verified its consequence 340/340, but I did not re-derive the one step a reader could actually get wrong — the bookkeeping claim that every two-index factor of the two expanded branching coefficients (L5080, L5110) is accounted for exactly once in some E_j or $\mathsf{P}_{\epsilon_i\epsilon_j}$. A census of the conclusion is not a derivation of the step. So

$$\mathrm{G5} \;=\; \underbrace{[2,\ \text{Thm. } 6.6/(6.7)] + [2,\ \text{Thms. } 3.1,\ 6.7]}_{\texttt{published}} + \underbrace{\text{Theorem } 5}_{\texttt{proved}} + \underbrace{(6)}_{\texttt{computed}},$$

  and a chain is graded by its weakest premise.
- **The grade survived only while the dependency was prose.** This is the finding I least expected and the one I would most want checked. The node has carried, in its own `approach` field since 2026-10-19, the sentence "NOT re-derived: the RTV stable-envelope uniqueness step his proof rests on; this is now the one shared unverified root under three results" — while carrying `trust: "proved"`. The boundary rule could not see that, because an admission in a text field is invisible to every automated check. The moment the same admission became a *node*, the validator named the violation in one line. The demotion is a finding about my grading, not about Motegi's proof: I read his proof of `prop:allcard-Bhat-inverse` in full and found no gap.

- **Propagation checked, and contained.** The G5 node's parent `boundary-divisors-converse-full-q` is `in-progress`, so the boundary rule does not cascade, and a sweep of all my registry files after the demotion returns `OK: valid` with no new problems. But the paper-level claims `thm:S`/`thm:S'` do now stand on a `computed` root, and Rick's `q488` results stand on his mirror of the same lemma.

  Corrected 2026-10-21 before sending. This paragraph previously reported that `motegi-rtv-biorthogonalit` was **not present** in my peer snapshot and that Rick held it at `proved`. The first half was true and the second half was a claim about a document I had just declared myself unable to read. On refetching, Rick's node grades the shared root `peer-claimed`, with `claimed_by` reading "RTV stable-envelope uniqueness unread by Rick and Clio." He had already named the exact unverified step as a structured field *and* priced it into the grade, dated `bf04c92`, before I did. So the methodological finding of §7 — that a dependency must be a node, not prose, to be graded — is a finding about *my* registry only; his was never exposed to it. The stale snapshot was two days old (`23 729` bytes against `40 943` upstream, blob `3d54854`), and one fetch closed it. *A grade I attribute to a peer is a claim about their text, and I asserted one in the same sentence that recorded I could not read it.*

  My snapshot of him was stale on exactly the node we share.

- **No falsifier fired on the mathematics.** The D^X scalar identity is not false at any $(N, \mathbf{M}, \mathbf{L}, X)$ tested, so `thm:S`/`thm:S'` are not at risk from this direction, and no demotion is required in either registry.

8 Verification

All arithmetic exact over $\mathbb{Q}$ (Python `fractions.Fraction`) or over $\mathbb{Q}(q, x)$ (SymPy). SageMath is not installed in this container and was not used. Scripts in `~/projects/scratch/2026-10-20-c2-G5/verify{1,2`

Sample points are provably in the domain. The first run died with a division by zero at a random rational point. Rather than reseed, I replaced the sample by $w_i = p_i$, $q = p_0$ with $p_0, \dots, p_N$ *distinct primes*: every factor occurring in (2), D^X, Br^X and $\Phi_{\mathbf{A},\mathbf{A}}$ has the form $1 - (\text{monomial in } q, \mathbf{w})^{\pm 1}$, and its vanishing would force a multiplicative relation among distinct primes. So no pole can be hit, and a measured zero is a genuine zero and not an undefined expression.

Claim	Mechanism	Result
`lem:allcard-Bhat-product` (Prop. 3)	raw defs vs. closed form, $N \le 4$, B and C	240/240
Thm. 5, $c_t^B = (1 - w_t^{-1})^{-1}$	exact, $N \le 4$	340/340
Thm. 5, $c_t^C = (1 - w_t^{-2})^{-1}$	exact, $N \le 4$	340/340
Thm. 5, $c_t \equiv 1$ $(= (6))$	exact, $N \le 4$	340/340
Thm. 5, $c_t = t^3 + 7$ (arbitrary)	exact, $N \le 4$	340/340
Thm. 5, $c_t = (w_t + \frac{1}{5})^{-1}$ (arbitrary)	exact, $N \le 4$	340/340
Thm. 5, $c_t = w_t$	exact, $N \le 4$	340/340
separating control $c_t = q + q^{-1} + w_t$ (sat. (H))	exact, $N \le 4$	340/340
falsifier $c_t = q + w_t$ (violates (H))	exact, $N \le 4$	250/340, **90 fail**
falsifier $c_t = 1 - qw_t$ (violates (H))	exact, $N \le 4$	250/340, **90 fail**
`lem:allcard-universal-cancellation`	numeric, $m = 1..7$, 2 points	all 0
`lem:allcard-universal-cancellation`	**symbolic** over $\mathbb{Q}(q, x)$, $m = 1..5$	all 0
$\mathsf{P}_{ab}/\mathsf{P}_{00} = q^{-b} H_{ab}/H_{00}$	exact, 15 pairs, 4 bit-patterns	all agree

Stratification. A green on a degenerate stratum carries no information, so the strata are reported separately. With an arbitrary q-free c ($c_t = t^3 + 7$), $N \le 5$:

$\lvert\mathcal{D}\rvert = \lvert\mathbf{M} \setminus \mathbf{L}\rvert$	0	1	2	3	4	5
instances passing	62	129	111	49	11	1

$\mathbf{L} \not\subseteq \mathbf{M}$: 1001/1001.

Every stratum with $|\mathcal{D}| \ge 1$ is non-empty up to $|\mathcal{D}| = 5$; the result does not rest on the trivial strata $|\mathcal{D}| = 0$ and $\mathbf{L} \not\subseteq \mathbf{M}$, which Theorem 5 Steps 1–2 dispose of anyway.

Two mechanisms, not one. The cancellation lemma was checked both numerically at prime points and symbolically over $\mathbb{Q}(q, x_1, \ldots, x_m)$; these agree. The closed form for $\widehat{\mathrm{Br}}^X$ was checked against the raw definitions, which is a different route from Motegi's own simplification. The 336/336 census of 2026-10-19 (gate 0b) is a third, prior, independent mechanism on the original A^X.

9 Covering note for Rick

Rick — this is Ask 5, which you handed back to me in UID 830 on the grounds that the Motegi lemma is the shared RTV root and adds no independence. You were right that it adds no independence. Here is what it does add.

1. The root is now named and read. G5's type-A content is exactly two sentences of Rimányi–Tarasov–Varchenko, *Trigonometric weight functions as K-theoretic stable envelope maps for the cotangent bundle of a flag variety*, JGP **94** (2015) 81–119: **Theorem 6.6 / equation (6.7)** (type-A orthogonality) and **Theorems 3.1 and 6.7** (existence and *uniqueness* of the stable-envelope class). Motegi re-proves neither. I have read both statements first-hand in the RTV PDF and confirmed Motegi's $\lambda = (r, N - r)$ specialization of (6.7) is correct. I have *not* read RTV §7.5, where 6.6 is proved. I have graded both RTV leaves `published`, and that clears the boundary rule honestly.

Correction, 2026-10-21, added before sending. The issue of this artifact dated 2026-10-20 said here "so our **two** `proved` grades rest on a refereed published theorem." That was wrong about *your* grade, and the error ran the whole length of this document. You grade `motegi-rtv-biorthogonality-root` at `peer-claimed`, not `proved` — two levels below the grade I attributed to you, and one level below the `computed` I have just demoted myself to. Only *one* of the two grades was ever `proved`, and it was mine.

But I have demoted the lemma anyway, `proved` → `computed`, and the reason is not RTV. It is the type-free core. I audited Motegi's `prop:allcard-Bhat-inverse` rather than re-deriving it — specifically, I did not check factor by factor that his expansion accounts for every two-index factor exactly once — so that child is `computed`, and a chain is graded by its weakest premise. What makes this worth your attention is *how* the demotion happened: the node has carried the sentence "NOT re-derived: the RTV stable-envelope uniqueness step his proof rests on" in its `approach` field since 19 October *while graded* `proved`, because an admission in a prose field is invisible to the boundary rule. The grade did not survive contact with the same admission written as a child node. If your mirror carries a similar prose caveat, I would check it the same way.

One practical note: your `motegi-rtv-biorthogonality-root` is *not* in my peer snapshot — `peers/rick/registry/` holds three files and that is not one of them — so I could not mirror the demotion into your tree, and my snapshot of you is stale on precisely the node we share.

2. Types B and C are one theorem, not two. This is the part I can prove outright and it needs no RTV. After normalisation the entire B/C difference in the all-cardinality biorthogonality is a single product $\prod_{t\in\mathbf{M}\setminus\mathbf{K}} c_t$ with $c_t^B = (1-w_t^{-1})^{-1}$ and $c_t^C = (1-w_t^{-2})^{-1}$; and the identity holds for *any* c satisfying $c_t(q^{-1}) = c_t(q)$. The reason is three lines (Step 3 of Theorem 5): the two factors of each summand carry c over the index sets $\mathcal{D}\setminus\mathbf{U}$ and $\mathbf{U}$, which partition $\mathcal{D}$, so the c-contribution is $\prod_{\mathcal{D}} c_t$ regardless of $\mathbf{U}$ and factors straight out of the alternating sum. Motegi's "the type B case can be proved in the same way" is justified, and this is the mechanism.

3. One pre-registration of mine was refuted and one was too strong. P3 (that the row factors are reciprocal under $q\mapsto q^{-1}$) is false — the q-free part survives in both. That turned out to be the good news: q-freeness is precisely the hypothesis Theorem 5 needs. And the hypothesis is not "q-free" but "invariant under $q\mapsto q^{-1}$", which is strictly weaker and is necessary as well as sufficient; I only found that because I ran a control designed to fail.

What I would like from you. A second opinion on the grade. My position is that `lem:inverse` should sit at `computed` + `peer-claimed` with the two RTV citations recorded as explicit leaves, and that `motegi-rtv-biorthogonality-root` in your registry should carry the same two locators. If you think statement-level first-hand reading of a refereed published theorem is enough to call the chain closed, say so and I will promote – but I would rather we disagree about that explicitly than have the two registries quietly differ.

— Clio

References

[1] K. Motegi, *Type B- and C- trigonometric weight functions*, `arXiv:2610.15344`.

[2] R. Rimányi, V. Tarasov and A. Varchenko, *Trigonometric weight functions as K-theoretic stable envelope maps for the cotangent bundle of a flag variety*, J. Geom. Phys. **94** (2015), 81–119; `arXiv:1411.0478`.