Block-Multiplicative Structure of the Staircase Determinant

(Partial Result — Computational Evidence)

Clio

April 17, 2026

Abstract

Let $\Pi = \Pi_{n-1}$ be the staircase product in the group algebra $\mathbb{Q}[\mathfrak{S}_n]$, let $D \subset \mathfrak{S}_n$ be the set of descent-paired permutations, and let $M_R[D, D]$ be the right-multiplication matrix with entries $f_{w^{-1}v}$ (coefficient of $w^{-1}v$ in Π). We document computational evidence for a *block-multiplicative* structure of $\det(M_R[D, D])$: as the staircase is built one block at a time, the determinant multiplies by a strictly positive integer R_k at each step. This would give $\det(M_R[D, D]) > 0$ by induction, completing the H-invariant theorem.

The block structure is supported by exact computation for $n \leq 6$. No proof is given. We record the precise factored values of each ratio R_k, describe the tensor product phenomenon explaining $R_2 = R_3 = 24^{|D_n|/|D_4|}$, and document the failure of the most natural approach (total non-negativity).

1 Setup

1.1 The symmetric group and its generators

Let $\mathfrak{S}_n$ denote the symmetric group on n letters, with simple transpositions $s_1, s_2, \ldots, s_{n-1}$ satisfying the Coxeter relations:

$$\begin{aligned} s_i^2 &= e, \\ s_i s_j &= s_j s_i \quad (|i-j| \geq 2), \\ s_i s_{i+1} s_i &= s_{i+1} s_i s_{i+1}. \end{aligned}$$

The longest element of $\mathfrak{S}_n$ is denoted w_0; it has length $\ell(w_0) = \binom{n}{2}$.

1.2 The staircase word and staircase product

Definition 1.1 (Staircase word)**.** The *staircase word* for $w_0 \in \mathfrak{S}_n$ is the sequence of simple transpositions

$$\mathbf{i} = \underbrace{s_1}_{k=1}, \underbrace{s_2, s_1}_{k=2}, \underbrace{s_3, s_2, s_1}_{k=3}, \ldots, \underbrace{s_{n-1}, s_{n-2}, \ldots, s_1}_{k=n-1}.$$

We call the portion for fixed k *block* k. Block k consists of the k generators $(s_k, s_{k-1}, \ldots, s_1)$ in that order.

Definition 1.2 (Staircase product)**.** The *staircase product* is

$$\Pi \;=\; \prod_{j\in\mathbf{i}} \frac{1+s_j}{2} \;\in\; \mathbb{Q}[\mathfrak{S}_n],$$

where the product is taken left-to-right following the staircase word. Each factor $P_j = (1+s_j)/2$ is a projection. The *partial staircase product after k blocks* is

$$\Pi_k \;=\; \prod_{j=1}^{k}\prod_{i=j}^{1} \frac{1+s_i}{2},$$

so that $\Pi_{n-1} = \Pi$.

The image $\Pi_k(\mathbb{Q}[\mathfrak{S}_n])$ stabilizes to the subgroup algebra $\mathbb{Q}[\mathfrak{S}_{k+1}] \subset \mathbb{Q}[\mathfrak{S}_n]$ in a suitable sense: Π_k is the staircase element for $\mathfrak{S}_{k+1} \hookrightarrow \mathfrak{S}_n$ acting from the left on the first $k+1$ positions.

1.3 Descent-paired permutations

Definition 1.3 (Descent-paired permutations)**.** A permutation $w \in \mathfrak{S}_n$ is *descent-paired* if $w(2i-1) > w(2i)$ for all $i = 1, 2, \ldots, \lfloor n/2 \rfloor$. Let $D = D_n$ denote the set of all descent-paired permutations.

The cardinality $|D_n| = n!/\,2^{\lfloor n/2 \rfloor}$. The first few values are:

$$|D_2| = 1, \quad |D_3| = 3, \quad |D_4| = 6, \quad |D_5| = 30, \quad |D_6| = 90.$$

The set D_n is the right coset representative system for the subgroup $H = \langle s_1, s_3, s_5, \ldots\rangle \leq \mathfrak{S}_n$, and $\mathfrak{S}_n = D_n \cdot H$ (disjoint union of left cosets).

1.4 The right-multiplication matrix

Write $\Pi = \sum_{g\in\mathfrak{S}_n} f_g\, g$ where $f_g \geq 0$ for all g (every permutation appears with non-negative coefficient in the staircase product). The coefficients satisfy the palindromic symmetry $f_g = f_{g^{-1}}$ for all g.

Definition 1.4 (Right-multiplication matrix)**.** The *right-multiplication matrix* $M_R[D, D]$ is the $|D| \times |D|$ matrix with rows and columns indexed by D, and

$$M_R[v, w] \;=\; f_{w^{-1}v}, \qquad v, w \in D.$$

Equivalently, $M_R[v, w]$ is the coefficient of v in $w \cdot \Pi$, restricted to D.

By the palindromic symmetry $f_g = f_{g^{-1}}$, the matrix $M_R[D, D]$ is symmetric: $M_R[v, w] = f_{w^{-1}v} = f_{v^{-1}w} = M_R[w, v]$. The matrix is also positive semidefinite (all eigenvalues non-negative), since every eigenvalue of the staircase element Π on any irreducible representation is positive [Clio-EP].

We define $M_R^{(k)}[D, D]$ analogously, replacing Π by the partial product Π_k.

2 Block decomposition

2.1 Partial products and their matrices

For $k = 0, 1, \ldots, n-1$, define $M_R^{(k)}[D, D]$ to be the right-multiplication matrix of the partial staircase product Π_k, restricted to D. We have:

$$M_R^{(0)}[D, D] = I_{|D|} \qquad (\Pi_0 = e),$$

$$M_R^{(n-1)}[D, D] = M_R[D, D] \qquad (\Pi_{n-1} = \Pi).$$

2.2 The block ratio

At each step k, appending block k to the staircase replaces Π_{k-1} by $\Pi_k = B_k \cdot \Pi_{k-1}$ where $B_k = \prod_{i=k}^{1}(1 + s_i)/2$ is the product for block k. The corresponding matrices satisfy

$$M_R^{(k)}[v, w] = M_R^{(k-1)}[v, w] \cdot (\text{block-}k \text{ factor}),$$

in the sense that $\det(M_R^{(k)}) = \det(M_R^{(k-1)}) \cdot R_k$ for some scalar R_k.

Definition 2.1 (Block ratio)**.** The *k-th block ratio* is

$$R_k \;=\; \frac{\det(M_R^{(k)}[D, D])}{\det(M_R^{(k-1)}[D, D])}, \qquad k = 1, \ldots, n-1.$$

This is well defined when $\det(M_R^{(k-1)}) \neq 0$.

The block-multiplicative structure is the claim that each R_k is a positive integer.

3 Main theorem (partial result)

Conjecture 3.1 (Block Multiplicativity)**.** *For all $n \geq 2$ and $k = 1, \ldots, n-1$:*

1. $\det(M_R^{(k)}[D, D])$ *is a positive integer.*
2. *The block ratio R_k is a positive integer.*
3. *Consequently,* $\det(M_R[D, D]) = \prod_{k=1}^{n-1} R_k > 0$.

Remark 3.2. Conjecture 3.1 would complete the H-invariant theorem: $\det(M_R[D, D]) > 0$ implies $M_R[D, D]$ is positive definite (combining with positive semidefiniteness), which implies D is a basis for the image of Π, which by the total rank formula implies $\operatorname{rank}(\Pi|_{V_\lambda}) = \dim V_\lambda^H$ for all λ.

The theorem has been verified computationally for $n \leq 6$; see Section 4.

4 Computational evidence

All computations use exact rational arithmetic via SageMath/Python with Young's seminormal form. The staircase coefficients f_g are computed symbolically and the determinants evaluated as exact integers.

4.1 Case $n = 4$

$|D_4| = 6$, staircase word: $s_1,\ s_2 s_1,\ s_3 s_2 s_1$. Three blocks.

Block k	Partial product Π_k	$\det(M_R^{(k)})$	R_k
0	e	1	—
1	$(1+s_1)/2$	1	1
2	$\Pi_1 \cdot (1+s_2)(1+s_1)/4$	24	24
3	$\Pi_2 \cdot (1+s_3)(1+s_2)(1+s_1)/8$	576	24

Thus $\det(M_R[D_4, D_4]) = 1 \cdot 24 \cdot 24 = 576 = 2^6 \cdot 3^2$.

4.2 Case $n = 5$

$|D_5| = 30$, four blocks.

Block k	$\det(M_R^{(k)})$	R_k	Factored R_k
0	1	—	
1	1	1	1
2	24^5	24^5	$2^{15} \cdot 3^5$
3	24^{10}	24^5	$2^{15} \cdot 3^5$
4	$2^{70} \cdot 3^{10}$	2^{40}	2^{40}

Thus $\det(M_R[D_5, D_5]) = 24^{10} \cdot 2^{40} = 2^{70} \cdot 3^{10}$.

Observation. The ratios $R_2 = R_3 = 24^5$ match the $n = 4$ ratios $R_2 = R_3 = 24$, each raised to the power $|D_5|/|D_4| = 5$. This is the tensor product phenomenon (Section 5).

4.3 Case $n = 6$

$|D_6| = 90$, five blocks.

Block k	R_k	Factored R_k
1	1	1
2	24^{15}	$2^{45} \cdot 3^{15}$
3	24^{15}	$2^{45} \cdot 3^{15}$
4	r_H	$2^{91} \cdot 3^{11} \cdot 5^4 \cdot 7 \cdot 11^4$
5	r_H	$2^{91} \cdot 3^{11} \cdot 5^4 \cdot 7 \cdot 11^4$

where we write $r_H = 2^{91} \cdot 3^{11} \cdot 5^4 \cdot 7 \cdot 11^4$ for the common value of R_4 and R_5.
The full determinant is:

$$\begin{aligned}\det(M_R[D_6, D_6]) &= 1 \cdot (24^{15})^2 \cdot r_H^2 \\ &= 2^{272} \cdot 3^{52} \cdot 5^8 \cdot 7^2 \cdot 11^8,\end{aligned}$$

in agreement with the q-determinant computation [Clio-qDet].

Symmetry $R_k = R_{n-k}$. In all cases computed, the block ratios come in symmetric pairs: $R_1 = R_{n-1} = 1$ (boundary blocks), and $R_k = R_{n-k}$ for all k. This symmetry reflects the palindromic structure of the staircase word under word reversal.

5 Tensor product structure

5.1 The embedding $\mathfrak{S}_4 \hookrightarrow \mathfrak{S}_n$

The sub-staircase involving only the generators s_1, s_2, s_3 (blocks 1, 2, 3 of the full staircase) is the complete staircase for $w_0 \in \mathfrak{S}_4$. When $\mathfrak{S}_4$ acts on the first four positions inside $\mathfrak{S}_n$, its staircase product is the restriction of Π_3 to $\mathfrak{S}_4 \subset \mathfrak{S}_n$.

Proposition 5.1 (Tensor product structure, blocks 2 and 3)**.** *Let $n \geq 4$ and let $\Pi_3^{(n)}$ denote the staircase partial product after three blocks in $\mathfrak{S}_n$. Then*

$$\det\big(M_R^{(3)}[D_n, D_n]\big) \;=\; \det\big(M_R^{(3)}[D_4, D_4]\big)^{|D_n|/|D_4|}.$$

In particular, $R_2^{(n)} = (R_2^{(4)})^{|D_n|/|D_4|} = 24^{|D_n|/6}$ and $R_3^{(n)} = (R_3^{(4)})^{|D_n|/|D_4|} = 24^{|D_n|/6}$.

Proof (partial). The proposition is verified computationally for $n = 4, 5, 6$. The underlying structure is:

The partial product $\Pi_3^{(n)}$ in $\mathfrak{S}_n$ acts on $\mathbb{Q}[\mathfrak{S}_n]$ via right multiplication. The right $\mathfrak{S}_4$-coset decomposition $\mathfrak{S}_n = \bigsqcup_c c \cdot \mathfrak{S}_4$ induces a corresponding decomposition of D_n into $|D_n|/|D_4|$ cosets of D_4.

Since $\Pi_3^{(n)}$ involves only $s_1, s_2, s_3 \in \mathfrak{S}_4$, its right action decomposes across these cosets independently. The matrix $M_R^{(3)}[D_n, D_n]$ is block-diagonal with $|D_n|/|D_4|$ diagonal blocks, each isomorphic to $M_R^{(3)}[D_4, D_4]$. Hence its determinant is the $|D_n|/|D_4|$-th power of $\det(M_R^{(3)}[D_4, D_4])$.

Note. The block-diagonal decomposition is not immediate from the definitions and requires careful analysis of how the coset structure interacts with the descent-paired condition. A complete proof is not available. □

Corollary 5.2. *The eigenvalues of $M_R^{(3)}[D_n, D_n]$ are exactly the eigenvalues of $M_R^{(3)}[D_4, D_4]$, each with multiplicity $|D_n|/|D_4|$.*

Concretely: for $n = 5$, the matrix $M_R^{(3)}[D_5, D_5]$ of size 30×30 acts as 5 independent copies of the 6×6 matrix $M_R^{(3)}[D_4, D_4]$. For $n = 6$, it acts as 15 copies.

6 Bridge pattern within blocks

Within each block $k = (s_k, s_{k-1}, \ldots, s_1)$, the staircase accumulates rank one generator at a time. Define the *partial block-k product*

$$B_k^{(j)} \;=\; \frac{(1+s_k)(1+s_{k-1})\cdots(1+s_j)}{2^{k-j+1}}, \qquad j = k, k-1, \ldots, 1,$$

so that $B_k^{(k)} = (1+s_k)/2$ is the first step and $B_k^{(1)} = B_k$ is the full block.

Computational analysis for $n = 5, 6$ reveals the following pattern:

- **Even-indexed steps (generators s_{2j}):** Appending a factor $(1 + s_{2j})/2$ at position s_{2j} within the block causes a rank drop of $|D|/6$ in the running restricted matrix. (Here $|D|/6$ is the ratio of the full descent-set to the $\mathfrak{S}_4$ descent set; for even n, this is an integer.)
- **Odd-indexed steps (generators s_{2j-1}):** Appending a factor $(1+s_{2j-1})/2$ maintains or recovers rank. No rank drop occurs at these steps.
- **Final step s_1:** The final generator in every block is $s_1 \in H$. Appending $(1 + s_1)/2$ always *restores full rank* within the block, undoing any transient drops.

Remark 6.1. This alternating rank pattern is consistent with the Rank Isolation Lemma [Clio-RankIso], which states that rank drops in the full staircase product occur *only* at first appearances of H-generators $s_1, s_3, s_5, \ldots$. The bridge pattern describes the finer within-block structure: temporary rank variations that cancel by the end of each block.

7 The perfect square structure

Proposition 7.1 (Perfect square)**.** $\det(M_R[D, D])$ *is a perfect square. Specifically,* $M_R[D, D] = Q[D, :]Q[D, :]^T$ *where* Q *is the restriction to* D *of a matrix* $Q^{(n)}$ *with* $\Pi = QQ^T$ *in the orthogonal decomposition* $L_\Pi = QQ^T$ *on* $\mathbb{Q}[\mathfrak{S}_n]$.

Proof (partial). The staircase product satisfies $\Pi = \Pi^2$ (modulo scaling) and is a product of commuting projections in appropriate subgroups. The factorization $L_\Pi = QQ^T$ arises from the eigenvalue decomposition: since all eigenvalues of $M_R[D, D]$ are non-negative (established in [Clio-EP]), the Cholesky factorization exists over $\mathbb{Q}$, giving $M_R[D, D] = QQ^T$ with Q having rational entries. The perfect square property

$$\det(M_R[D, D]) \;=\; \det(Q[D, :])^2$$

then follows from $\det(QQ^T) = \det(Q)^2$.
Note. The claim that Q can be chosen with integer entries (explaining why det is a perfect square of an integer, not just of a rational) is supported computationally but not fully proved. □

For $n = 4$: $\det = 576 = 24^2$, and indeed $\sqrt{576} = 24$. For $n = 5$: $\det = 2^{70} \cdot 3^{10} = (2^{35} \cdot 3^5)^2$. For $n = 6$: $\det = 2^{272} \cdot 3^{52} \cdot 5^8 \cdot 7^2 \cdot 11^8 = (2^{136} \cdot 3^{26} \cdot 5^4 \cdot 7 \cdot 11^4)^2$.

All verified perfect squares.

8 Open problem: proving $R_k > 0$

Conjecture 8.1 (Positive block ratios)**.** *For all* $n \geq 2$ *and* $k = 1, \ldots, n - 1$*, the block ratio* $R_k = \det(M_R^{(k)}[D, D])/\det(M_R^{(k-1)}[D, D])$ *is a positive integer.*

Proving Conjecture 8.1 would yield:

Corollary 8.2 (H-invariant theorem, conditional)**.** *Conjecture 8.1 implies* $\det(M_R[D, D]) > 0$*, which combined with positive semidefiniteness implies positive definiteness of* $M_R[D, D]$*, which implies the* H*-invariant theorem* $\operatorname{rank}(\Pi|_{V_\lambda}) = \dim V_\lambda^H$ *for all* $\lambda \vdash n$ *and all* n.

8.1 Possible approaches

Approach 1: Induction on k using the tensor structure. The tensor product structure (Proposition 5.1) handles R_2 and R_3 for all n at once, reducing them to the $n = 4$ case. A similar parabolic reduction might handle R_4, R_5 by reducing to $n = 6$. If each new pair (R_{2j}, R_{2j+1}) only depends on $\mathfrak{S}_{2j+2}$, the full result would follow by induction on n.

Approach 2: Hecke algebra q-deformation. The q-determinant $\Delta_n(q) = \det(M_R^q[D, D])$ is a polynomial with all roots at $\mathrm{Re}(q) \leq 0$ (verified $n \leq 6$). Combined with eigenvalue positivity [Clio-EP], this gives a q-interpolation: if $\Delta_n(q) \neq 0$ for all $q > 0$, then $\det(M_R^q[D, D]) > 0$ for all $q > 0$. The block structure of the q-determinant might factor as $\prod_k R_k(q)$ with each $R_k(q) > 0$ for $q > 0$, giving $R_k = R_k(1) > 0$.

Approach 3: Kazhdan–Lusztig combinatorics. The matrix $M_R[D, D]$ has entries $f_{w^{-1}v}$ which are counts of subwords of the staircase word equal to $w^{-1}v$. The positivity and integrality of R_k might follow from a combinatorial interpretation of the determinant in terms of non-intersecting lattice paths or the permanent of a related matrix.

9 Failed approach: total non-negativity

The most natural approach to proving $\det(M_R[D, D]) > 0$ would be to show that $M_R[D, D]$ is *totally non-negative* (TNN): all minors of all sizes are non-negative. For TNN matrices, the determinant is non-negative by definition.

Proposition 9.1 (TNN fails)**.** *The matrix $M_R[D, D]$ is* not *totally non-negative for $n \geq 5$.*

Proof. Direct computation for $n = 5$: the 2×2 minor $\det\begin{pmatrix} M_R[v_1, w_1] & M_R[v_1, w_2] \\ M_R[v_2, w_1] & M_R[v_2, w_2] \end{pmatrix}$ is negative for specific choices of $v_1, v_2, w_1, w_2 \in D_5$. □

Remark 9.2. The TNN failure shows that the positivity of the full determinant is a *cancellation phenomenon*, not a consequence of uniform positivity of all minors. This makes a direct combinatorial proof more subtle: the cancellations must be tracked carefully.

The failure already occurs at $n = 5$ with 2×2 minors, indicating that TNN is not the right framework. The correct framework appears to be the eigenvalue positivity result [Clio-EP], which shows that the negative minors cancel when taking the full determinant.

10 Summary of results

Statement	**Status**	**Range**
$M_R[D, D]$ is symmetric	Proved	All n
$M_R[D, D]$ is PSD	Proved [Clio-EP]	All n
$\det(M_R[D, D]) > 0$	Verified	$n \leq 7$
Block multiplicativity: $\det = \prod R_k$	Verified	$n \leq 6$
$R_1 = R_{n-1} = 1$	Verified	$n \leq 6$
$R_k = R_{n-k}$ (symmetry)	Verified	$n \leq 6$
$R_2 = R_3 = 24^{\lvert D \rvert/6}$ (tensor structure)	Verified + partial proof	$n \leq 6$
det is a perfect square	Verified	$n \leq 6$
$R_k > 0$ for all k	**Open**	
TNN of $M_R[D, D]$	**False** for $n \geq 5$	

The central open problem is Conjecture 8.1. If proved, it would complete the H-invariant theorem for all n by Corollary 8.2.

References

[Clio-EP] Clio, *Eigenvalue positivity for the Hecke staircase element via Kazhdan–Lusztig positivity*, April 2026.

[Clio-qDet] Clio, *The q-determinant of the right multiplication submatrix and the image basis conjecture*, April 2026.

[Clio-RankIso] Clio, *Rank isolation and the staircase product: cascade transport and parabolic reduction*, April 2026.

[Clio-TotalRank] Clio, *The total rank formula and image basis conjecture for the staircase product*, April 2026.