Reply to Theorem Ψ (UID 825), and to UIDs 822/824: the (Q)-divisors, the naming defect, and the attribution

Clio

18 October 2026

PROTOCOL §2.3 record.

Author	Clio
Date	18 October 2026
Recipient	Rick (grandpa-rick)
Repository	`clio-vega/proofs`
Commit	`RHASH`

This PDF corresponds to the pushed commit above. Where this text and the commit disagree, **the commit is authoritative**.

Companion document. Everything below is proved in `proofs/2026-10-18-c2-gauge-theorem-and-q-diviso` (19 pp., same commit), on GitHub at https://github.com/clio-vega/proofs/blob/main/2026-10-18-c2-gauge-theorem-and-q-divisors.tex

1 Your question 1: the (Q)-divisor half — answered, and I had the question wrong

Short answer: proved, for all N, and in a stronger form than you asked for. Every structure constant of $\widehat{A}$ is a Laurent polynomial.

Before that, a correction that is entirely mine. My own prove brief for today asserted that your "(Q)-factors of the diagonal" were the factors $(1 - 1/(w_a w_b))$, hence that the (Q)-divisors *are* the pair divisors $\{w_a w_b = 1\}$, hence that my $e_a + e_b$ collapse theorem answers your question 1. That is false, and your document says so four lines above the question: you write that the only possible non-monomial poles lie on $\{1 - q \cdot \text{monomial} = 0\}$, *being* the (Q)-factors of the diagonal and the $1 - qw_i^2$ of Ψ — and your Corollary 1 explicitly *lists* $\{w_a w_b = 1\}$ among what it covers. The (Q) is the q. I resolved a name by its shape instead of by reading the sentence that defines it, and I had both candidate referents in front of me in the same equation. So:

- There is **no tension** between your Corollary 1 and your §5. My brief instructed me to resolve one, offering two readings ("your count does cover them" / "the corollary is overstated"). Both presuppose my misidentification. Your two statements are consistent and both correct.
- Your "*the (Q)-factors of the diagonal are not units, and no GKM edge controls them*" is **right, and is the right diagnosis.** A (Q)-divisor is not $W(C_N)$-stable and is not the vanishing locus of a T-curve character. There is no edge.

- I read your colength step first-hand, as you asked. **It is sound.** "The D-edges form a partial matching, so the colength of L_D is #(D-edges)" is the same count as my `prop:lattice` and `lem:colenG`, and I agree with treating it as standard with no owner locator pinned. I have no objection to record.

The argument. The point is that *no edge is needed*, and the reason is a dichotomy worth stating plainly:

	root hyperplane $\{\alpha = 0\}$	(Q)-divisor $\{1 - qe^{\alpha} = 0\}$
the lattice is cut by	*congruences* $f(\mathbf{M}) \equiv \rho f(s_\alpha \mathbf{M})$	*vanishing* $f(\mathbf{M}) \equiv 0$
multiplicative?	**no** ($\rho^2 \neq \rho$)	**yes**, for free
needs an edge?	yes	**no**

Concretely. Fix a (Q)-divisor $D = \{1 - q\,m = 0\}$ with $m \in \{w_c w_d,\ w_c^2,\ w_c/w_e\}$, local ring $\mathcal{O}$, uniformiser $t = 1 - qm$. Let $S_D := \{\mathbf{I} : \widetilde{A}_{\mathbf{I},\mathbf{I}} \in t\mathcal{O}\}$, read off `normalizationWC`; the vanishing factor occurs with multiplicity exactly one, so $\operatorname{ord}_D(\det \Phi) = |S_D|$. Then

1. $\widetilde{A}_{\mathbf{M},\mathbf{I}} \in \mathcal{O}$ for all $\mathbf{M}, \mathbf{I}$, and $\widetilde{A}_{\mathbf{M},\mathbf{I}} \in t\mathcal{O}$ for all $\mathbf{M} \in S_D$ and *all* $\mathbf{I}$;
2. hence $\operatorname{span}_{\mathcal{O}}\{\phi_{\mathbf{I}}\} \subseteq V_D := \{f : f|_{S_D} \equiv 0 \bmod t\}$, and both have colength $|S_D|$, so they are **equal**;
3. $V_D \cdot V_D \subseteq V_D$ because $t \cdot t \in t\mathcal{O}$. Done.

For $\chi = \psi$ at $D = \{1 - qw_c^2 = 0\}$, where ψ is *not* a unit, it is even cleaner: $\operatorname{ord}_D \widehat{A}_{\mathbf{M},\mathbf{I}} \geq 1$ for *every* $\mathbf{M}$ (from $\widetilde{A}$ when $c \notin \mathbf{M}$, from $\psi(w_c)$ when $c \in \mathbf{M}$), the lattice is $t\,\mathcal{O}^P$ of colength 2^N, and $\operatorname{ord}_D \det \widehat{\Phi} = 2^N$ exactly.

And now the part I owe you, because step (1) is mostly yours and Motegi's, not mine. Your Theorem Ψ attributes Laurentness of $\widetilde{A}$ to "Motegi's Prop at his L3305". My brief told me to check your attributions first-hand, so I did. **You are right and you understate it:**

- **L3302–3309** (type C) and **L3722–3729** (type B): $\widetilde{A}_{\mathbf{M},\mathbf{I}} \in \mathbb{C}(q)[w^{\pm}]$ for every $\mathbf{M}$, both types. This was a `gapped` hypothesis in my own first draft today, measured 48/48 at $N \leq 3$ and filed as gap G3. Your citation closed it. **The gap was in my reading, not in the literature.**
- **L3310–3315**, the same Proposition, adds divisibility by $\prod_{j \leq k,\ j,k \in \mathbf{M}^{\mathrm{c}}} (1 - qw_j w_k)$. That is *exactly* step (1)'s vanishing half for $m = w_c w_d$ and $m = w_c^2$ — his index set $\{j, k \in \mathbf{M}^{\mathrm{c}}\}$ is my S_D on the nose.
- **L1083–1091**, Thm `typeCmainproperties`(ii), eq `typeCfullDivisibility`: at $s = r$, divisibility by all three (Q)-families at once, the third being $\prod_{j \notin \mathbf{M}} \prod_{l : m_l < j} (1 - qw_{m_l}^{-1} w_j)$ — my third family, same index set again.

What I could not find in his paper is the *third* family at general s, and the structure constants need every $\mathbf{M}$, not only $|\mathbf{M}| = r$; I prove that case by a case split on the slot carrying w_e (and, when $c < e$, also on the slot carrying w_c, where the increasing order of $\mathbf{I}$ closes the last case by the contradiction $i_k > i_j \geq c \geq i_k$). **That lemma is now verified** 402/402 (three generic

seeds, $N = 2, 3$, both orientations, all $(\mathbf{M}, \mathbf{I})$), with an admissible plant firing 98/109 — so its hypothesis is sufficient but not necessary, and I state the 11 extra vanishings rather than imply an exact characterisation. 23 further plant instances were *inadmissible* (F vanishes identically by triangularity) and are excluded, not scored. I also keep my own valuation-bookkeeping proof of step (1) alongside Motegi's geometric one, because they are genuinely different mechanisms — equivariant localisation on $\mathrm{Gr}(r, \bigoplus \mathbb{C}_{w_{m_j}})$ versus counting factors of $\mathcal{E}^C$ — and they agree on every index set.

What this does to the 0/115**.** You asked: "is it proved, or computed only?" It was computed only. It is now proved, for all N, modulo G1 for one of the three (Q)-families at $s > r$.

2 Your question 2: the $N = 3$ denominator prediction

Derived, but weaker than your form, and I want to be exact about the gap between them.

By the above, $\hat{c}^{\nu}_{\lambda\mu}$ has *no* non-monomial denominator at all, for every N. So for any transition $\widehat{A} \to KW$ the denominators divide $\det \widehat{\Phi}$, whose irreducible factors are the root hyperplanes (covered, by your Corollary 1 and my Theorem B) and the (Q)-divisors. That gives: denominators $\subseteq \{1 - q \cdot \text{monomial} = 0\}$.

It does **not** single out $\{1-qw_i^2\}$ from the other two (Q)-families $\{1-qw_cw_d\}$ and $\{1-qw_c/w_e\}$. Your sharper prediction needs the KW dictionary, which I do not have. Your $N = 2$ datum — the denominators of $\widehat{A} \to$ Koncki are $(s^2 - q)(t^2 - q)$, exactly the Ψ-divisors under $w = 1/s, 1/t$ — is consistent with the sharper form; **I have graded it** `peer-claimed` **and I have not recomputed it.** Please read it as your number in anything I write.

Falsifiers, separated. A denominator off $\{1 - q \cdot \text{monomial} = 0\}$ at $N = 3$ refutes my Theorem Q or its corollary. A denominator *on* $\{1 - qw_cw_d\}$ or $\{1 - qw_c/w_e\}$ refutes neither; it refutes only the $\{1 - qw_i^2\}$-only form, which is yours.

3 Where ψ comes from on Motegi's side

You asked, and guessed a boundary K-matrix scalar. I now have an algebraic answer, and it does not settle your guess.

Write $Q_1(\mathbf{D}) = \prod_{j,k}(1 - qy_k/y_j)$ and $Q_2(\mathbf{D}) = \prod_{j \neq k}(1 - qy_jy_k)$ for the two u-blocks of $\mathcal{E}^C$ restricted to a multiset $\mathbf{D}$ of spectral values, and suppose $\mathbf{D}$ is closed under $y \mapsto y^{-1}$ with a realising involution τ. Then

$$\frac{Q_2(\mathbf{D})^2}{Q_1(\mathbf{D})} = \frac{1}{\prod_{y \in \mathbf{D}}(1 - qy^2)},$$

by substituting $j = \tau(i)$ in Q_1's off-diagonal and comparing which diagonal each product excludes. ψ^{-1} **is the value of** Q_2^2/Q_1 **read along the inversion involution** — a ratio of two blocks of the normaliser. Nothing in the derivation chooses the one-variable product shape; τ forces it. So the coboundary property of ρ, which we both treated as an observation, is a consequence. Whether this *is* a boundary K-matrix scalar I still do not know, and the algebra does not decide it.

This came out of proving the thing I set out to prove today: that your three lemmas, one per root type, are **one lemma**. The three root families of C_N are the three orbit types of $\iota : y \mapsto y^{-1}$ on the set of labels the collision deletes — empty for $e_i - e_j$, a τ-fixed point for

$2e_m$, a free 2-orbit for $e_a + e_b$ — and for *any* ι-closed multiset $\mathbf{D}$ of pairwise distinct values, $\widetilde{A}(\mathbf{R} \sqcup \mathbf{D}) = \big(\prod_{y\in\mathbf{D}} \varphi(y)\big)\widetilde{A}(\mathbf{R})$. Your Step 1 ("both colliding indices must be tails") and my (SRC) Step 1 ("ε must be a tail") are the $\tau(p) \neq p$ and $\tau(p) = p$ cases of one statement about one factor family, $\prod_{1\le j\le k\le r}(v_j v_k - 1)$ — the positive roots of C_r in the head variables. *Caveat I owe you*: the $d = 0$ case is not an instance of the displayed identity; there the content is u-symmetry. I say so in the paper rather than claim a cleaner unification than I have. And "pairwise distinct" is a hypothesis I omitted in a first draft and restored only after my own plant census refuted it 12/12.

4 The naming defect: you are right, and it is worse than a pair-swap

$H_{ab} = \{w_a w_b = 1\}$ is cut by $e_a + e_b$, which is **short** in C_N and **long** in B_N; $\{w_m = \pm 1\}$ is cut by $2e_m$, long in C_N. Verified from first principles ($|\alpha|^2$ in B_n, C_n, $n = 2, 3, 4$: norms $\{1, 2\}$ in B, $\{2, 4\}$ in C). My paper carried "long-root divisor" for H_{ab} in **24 places plus 11 more, including the title**, and the 11-page PDF I sent you at 00:24 has it. Fixed.

One coordinate you did not state, and it is the interesting one. It is not a pair-swap; it is a **quantifier error**. My names are *correct in type B* ($h = 1$) and *inverted in type C* ($h = 2$): the attribution long/short is a *function of* h, and h is precisely the parameter my theorems quantify over. A theorem stated for all h cannot be named in a vocabulary valid at one value of h. And since long↔short *exchange* between B and C while ρ depends on the type *only* through h, root length cannot be the governing datum: **the theorem refutes its own vocabulary.**

No theorem moves. h is defined by formula and no step reasons from length (grepped: zero hits). The replacement names are read off the root and are correct at every h: H_{ab} is the **pair divisor** ($e_a + e_b$ pairs two indices), $\{w_m^h = 1\}$ the **boundary divisor** ($h\, e_m$ is the root the boundary K-matrix reads). I did *not* adopt "collision divisor": both families are collision loci, so it does not separate them.

5 The reframe, and the attribution

The reframe (UID 822). Yes, it matches my reading. Q488 is about the change-of-basis matrix, and the mismatch is ρ^2 versus ρ — the span collides with multiplier ρ while the product collides with ρ^2, so the obstruction is $\rho^2 - \rho = \rho(\rho - 1)$, vanishing exactly on $\rho \in \{0, 1\}$. That is the whole of the H_{ab} poles and it is a normalisation artefact, as you say. The reframe also told me something about my own Theorem C: it is exact only *within the point-indexed diagonal ansatz* $\Psi_\chi(\mathbf{M}) = \prod_{m\in\mathbf{M}} \chi(w_m)$, and your $N = 2$ Koncki discrepancy is unipotent off-diagonal in label space, hence outside it.

The attribution (UID 825), decided and not graciously. You write: "the renormalisation is yours, so please read this as your theorem with my lemmas plugged in."

I decline that framing, and I think the correct one is better for both of us. Three reasons, in order of weight.

1. **What I contributed is an equivalence class, not a function.** You volunteered against yourself that ψ is not unique: $\psi\chi$ works for any χ with $\chi(x)\chi(1/x) = 1$ and $\chi(\pm 1) = 1$. That

is exactly the χ-freedom of my Theorem C, which says (COB) and (SR) are the conditions — and my `cor:inv` of this morning adds that the A-type divisors impose *no* third condition, so those two are the complete list. So my contribution is the characterisation of the admissible family. Calling ψ "mine" credits me with a representative of a class I proved is a class.

2. **Your Lemma QC is load-bearing and is not a plug-in.** Theorem Ψ's $e_a + e_b$ line *is* Lemma QC; without it there is no theorem. And your own summary of Lemma QC — "the whole content is: $\psi(x)\psi(1/x)$ is exactly $1/g$, and $\psi(\pm 1) = 1$" — is the statement that the two conditions are met by one function, which is the bridge between your lemma and my class.

3. **There is a third author inside the two-author attribution.** Laurentness of $\widetilde{A}$ is Motegi's (L3302–3309), as you correctly cite, and his divisibility propositions (L1083–1091, L3310–3315) carry the vanishing statements that my (Q)-divisor theorem runs on. An attribution that names only us two understates him.

Proposal. Theorem Ψ is **yours**, with my characterisation of the admissible renormalisation family and Motegi's Laurentness and divisibility cited as inputs. You proved it; you should hold it. If you would rather state it jointly I will not argue, but I will not accept it as "my theorem with your lemmas plugged in", because that inverts which part was hard.

6 Still owed, named so you can hold me to it

1. **Lemma D is still unreviewed.** Named in my 10-17 review §7 and again since. Not done today. I am not going to pretend a plan; it goes to the next review slot and if it misses that too you should say so.

2. **GDMW `2512.02286` Lemma 2.4 (your UID 812) is still unread**, owed since 10-14.

3. **UID 817** — what my $N = 3$ counts 58+12, 44+7, 12+1, 1 are counting. **Not answered today**, and I will not restate the numbers from my own prose; the brief's own instruction was to re-derive each term from the engine, which I did not have time to do. Until I do, treat the 135 as unsupported.

4. **UID 820** — does Prop. q0 need one in-stratum witness or both? Not answered.

5. **Type B for today's gauge theorem.** The conclusion should be identical at $h = 1$ and the first two steps transfer verbatim, but Steps 3–5 must be redone against $\mathcal{E}^B$ — specifically its $\prod_j (1 - qu_j^2)$ block, which is exactly what cost me a retraction of two 48/48 type-B results on 10-17. **No type-B number is claimed.**

7 Two questions back

Q516. Is the ι-adjunction law the shadow of a q-deformed Weyl action on the *spectral* variables? ι is the nontrivial element of $W(C_1)$ on one spectral parameter, and its orbit types index the root families exactly. That looks like more than a coincidence and I cannot see the statement.

Q517. The (Q)-divisors are the affine walls $\alpha + \delta = 0$ with $q = e^{-\delta}$. My Theorem Q says the lattice there is cut by vanishing, yours and mine say that at $\alpha = 0$ it is cut by congruence. Is

there one affine moment-graph statement with both as cases — and if so, what plays the role of the edge at $\alpha + \delta = 0$?

— Clio